

2-Chloroethyl 4-nitrobenzoate

Hao Wu,* Min-Hao Xie, Pei Zou, Ya-Ling Liu and Yong-Jun He

Key Laboratory of Nuclear Medicine, Ministry of Health, Jiangsu Key Laboratory of Molecular Nuclear Medicine, Jiangsu Institute of Nuclear Medicine, Wuxi 214063, People's Republic of China

Correspondence e-mail: hajiaoyiya@sina.com


Received 27 September 2010; accepted 28 September 2010

Key indicators: single-crystal X-ray study; $T = 103\text{ K}$; mean $\sigma(\text{C}-\text{C}) = 0.002\text{ \AA}$; R factor = 0.034; wR factor = 0.084; data-to-parameter ratio = 16.5.

The title compound, $\text{C}_9\text{H}_8\text{ClNO}_4$, crystallizes with two molecules in the asymmetric unit. In each molecule, the carboxylate group is nearly coplanar with the benzene ring, forming dihedral angles of $2.4(1)$ and $4.9(1)^\circ$. In the crystal, molecules are linked through weak $\text{C}-\text{H}\cdots\text{O}$ and $\text{C}-\text{H}\cdots\text{Cl}$ hydrogen bonds. A short $\text{O}\cdots\text{N}$ contact of $2.7660(19)\text{ \AA}$ occurs between the nitro groups of adjacent molecules.

Related literature

For benzoates as intermediates in the chemistry of pigments and pharmaceuticals, see: Zhang *et al.* (1995, 1990). For a related structure, see: Wu *et al.* (2009).



Experimental

Crystal data

$\text{C}_9\text{H}_8\text{ClNO}_4$	$c = 18.325(4)\text{ \AA}$
$M_r = 229.61$	$\beta = 90.441(3)^\circ$
Monoclinic, $P2_1/n$	$V = 1957.0(7)\text{ \AA}^3$
$a = 4.9404(10)\text{ \AA}$	$Z = 8$
$b = 21.618(5)\text{ \AA}$	Mo $K\alpha$ radiation

$\mu = 0.38\text{ mm}^{-1}$
 $T = 103\text{ K}$

$0.40 \times 0.20 \times 0.20\text{ mm}$

Data collection

Rigaku SPIDER diffractometer
Absorption correction: multi-scan
(*ABSCOR*; Higashi, 1995)
 $T_{\min} = 0.862$, $T_{\max} = 0.927$

18618 measured reflections
4466 independent reflections
3781 reflections with $I > 2\sigma(I)$
 $R_{\text{int}} = 0.033$

Refinement

$R[F^2 > 2\sigma(F^2)] = 0.034$
 $wR(F^2) = 0.084$
 $S = 1.00$
4466 reflections

271 parameters
H-atom parameters constrained
 $\Delta\rho_{\max} = 0.34\text{ e \AA}^{-3}$
 $\Delta\rho_{\min} = -0.23\text{ e \AA}^{-3}$

Table 1
Hydrogen-bond geometry (\AA , $^\circ$).

$D-\text{H}\cdots A$	$D-\text{H}$	$\text{H}\cdots A$	$D\cdots A$	$D-\text{H}\cdots A$
$\text{Cl}-\text{H}1\cdots\text{Cl}1'$	0.95	2.79	3.6234 (18)	147
$\text{C}2'\cdots\text{H}2'\cdots\text{O}3^{ii}$	0.95	2.50	3.318 (2)	145
$\text{C}4'\cdots\text{H}4'\cdots\text{O}2$	0.95	2.40	3.223 (2)	144
$\text{C}5\cdots\text{H}5\cdots\text{O}2^{ii}$	0.95	2.42	3.205 (2)	140
$\text{C}9\cdots\text{H}9\cdots\text{O}2^{iii}$	0.99	2.55	3.526 (2)	167

Symmetry codes: (i) $-x + 2, -y + 1, -z$; (ii) $-x + 1, -y + 1, -z + 1$; (iii) $-x, -y + 1, -z + 1$.

Data collection: *RAPID-AUTO* (Rigaku, 2004); cell refinement: *RAPID-AUTO*; data reduction: *RAPID-AUTO*; program(s) used to solve structure: *SHELXTL* (Sheldrick, 2008); program(s) used to refine structure: *SHELXTL*; molecular graphics: *SHELXTL*; software used to prepare material for publication: *SHELXTL*.

This work was supported by the Science Foundation of the Health Department of Jiangsu Province, China (No. H200934).

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: XU5038).

References

- Higashi, T. (1995). *ABSCOR*. Rigaku Corporation, Tokyo, Japan.
- Rigaku (2004). *RAPID-AUTO*. Rigaku Corporation, Tokyo, Japan.
- Sheldrick, G. M. (2008). *Acta Cryst. A* **64**, 112–122.
- Wu, H., Xie, M.-H., Zou, P., Liu, Y.-L. & He, Y.-J. (2009). *Acta Cryst. E* **65**, o3096.
- Zhang, A.-Y., Qian, B., Min, J. & Fang, Q.-X. (1995). *J. Shanxi Normal Univ. (Nat. Sci. Ed.)*, **23**, 44–47.
- Zhang, Z.-S., Wu, J.-G. & Deng, R.-W. (1990). *J. Lanzhou Univ. (Nat. Sci. Ed.)*, **26**, 69–75.

supplementary materials

Acta Cryst. (2010). E66, o2708 [doi:10.1107/S1600536810038742]

2-Chloroethyl 4-nitrobenzoate

H. Wu, M.-H. Xie, P. Zou, Y.-L. Liu and Y.-J. He

Comment

Benzoates are important intermediates in the chemistry of pigments and pharmaceuticals, which are widely used all over the world (Zhang *et al.*, 1995; Zhang *et al.*, 1990). The crystal structure of Methyl 4-nitrobenzoate has been reported (Wu *et al.*, 2009). As an extension of our study, we report here the crystal structure of the title compound.

The asymmetric unit of the title compound (Fig. 1) contains two crystallographically independent molecules in an asymmetric unit. The crystal data show that the bond lengths and angles are within expected ranges. The ester groups and the benzene rings in the two molecules are almost coplanar, as indicated by the dihedral angles of $2.4(1)^\circ$ ($O1/O2/C6/C7$; $C1—C6$) and $4.9(1)^\circ$ ($O1'/O2'/C6'/C7'$; $C1'—C6'$). The dihedral angle between the ring($C1—C6$) and the ring($C1'—C6'$) is $92.7(1)^\circ$.

In the crystal structure, adjacent molecules are linked together by the weak $C—H\cdots O$ and $C—H\cdots Cl$ hydrogen bonds (Table 1). Intramolecular $C—H\cdots O$ interactions are also observed. These hydrogen-bonding interactions stabilize the crystal structure.

Experimental

Commercial 2-chloroethyl 4-nitrobenzoate was recrystallized by slow evaporation of methanol/water solution (1:1 v/v). Colourless single crystals were formed after two weeks.

Refinement

H atoms were placed in calculated positions with $C—H = 0.99$ (methylene) and 0.95 \AA (aromatic), and were refined in a riding mode with $U_{\text{iso}}(\text{H}) = 1.2U_{\text{eq}}(\text{C})$.

Figures




Fig. 1. eA view of the title compound with the atomic numbering scheme. Displacement ellipsoids were drawn at the 50% probability level.

2-Chloroethyl 4-nitrobenzoate

Crystal data

$\text{C}_9\text{H}_8\text{ClNO}_4$

$F(000) = 944$

$M_r = 229.61$

$D_x = 1.559 \text{ Mg m}^{-3}$

Monoclinic, $P2_1/n$

Mo $K\alpha$ radiation, $\lambda = 0.71073 \text{ \AA}$

supplementary materials

Hall symbol: -P 2yn	Cell parameters from 5688 reflections
$a = 4.9404 (10) \text{ \AA}$	$\theta = 3.0\text{--}27.5^\circ$
$b = 21.618 (5) \text{ \AA}$	$\mu = 0.38 \text{ mm}^{-1}$
$c = 18.325 (4) \text{ \AA}$	$T = 103 \text{ K}$
$\beta = 90.441 (3)^\circ$	Prism, colorless
$V = 1957.0 (7) \text{ \AA}^3$	$0.40 \times 0.20 \times 0.20 \text{ mm}$
$Z = 8$	

Data collection

Rigaku SPIDER	4466 independent reflections
diffractometer	
Radiation source: Rotating Anode	3781 reflections with $I > 2\sigma(I)$
graphite	$R_{\text{int}} = 0.033$
ω scans	$\theta_{\text{max}} = 27.5^\circ, \theta_{\text{min}} = 3.0^\circ$
Absorption correction: multi-scan (<i>ABSCOR</i> ; Higashi, 1995)	$h = -6 \rightarrow 6$
$T_{\text{min}} = 0.862, T_{\text{max}} = 0.927$	$k = -28 \rightarrow 28$
18618 measured reflections	$l = -23 \rightarrow 20$

Refinement

Refinement on F^2	Primary atom site location: structure-invariant direct methods
Least-squares matrix: full	Secondary atom site location: difference Fourier map
$R[F^2 > 2\sigma(F^2)] = 0.034$	Hydrogen site location: inferred from neighbouring sites
$wR(F^2) = 0.084$	H-atom parameters constrained
$S = 1.00$	$w = 1/[\sigma^2(F_o^2) + (0.0441P)^2 + 0.680P]$ where $P = (F_o^2 + 2F_c^2)/3$
4466 reflections	$(\Delta/\sigma)_{\text{max}} = 0.001$
271 parameters	$\Delta\rho_{\text{max}} = 0.34 \text{ e \AA}^{-3}$
0 restraints	$\Delta\rho_{\text{min}} = -0.23 \text{ e \AA}^{-3}$

Special details

Geometry. All e.s.d.'s (except the e.s.d. in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell e.s.d.'s are taken into account individually in the estimation of e.s.d.'s in distances, angles and torsion angles; correlations between e.s.d.'s in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell e.s.d.'s is used for estimating e.s.d.'s involving l.s. planes.

Refinement. Refinement of F^2 against ALL reflections. The weighted R -factor wR and goodness of fit S are based on F^2 , conventional R -factors R are based on F , with F set to zero for negative F^2 . The threshold expression of $F^2 > \sigma(F^2)$ is used only for calculating R -factors(gt) etc. and is not relevant to the choice of reflections for refinement. R -factors based on F^2 are statistically about twice as large as those based on F , and R -factors based on ALL data will be even larger.

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (\AA^2)

	<i>x</i>	<i>y</i>	<i>z</i>	$U_{\text{iso}}^*/U_{\text{eq}}$
Cl1	0.58062 (8)	0.615621 (18)	0.60156 (2)	0.02137 (10)
O1	0.5719 (2)	0.70861 (5)	0.72719 (6)	0.0201 (2)
O2	0.4652 (2)	0.80061 (5)	0.77677 (6)	0.0230 (3)
O3	1.4451 (3)	0.74075 (6)	1.03965 (7)	0.0288 (3)
O4	1.5555 (2)	0.65746 (6)	0.98123 (7)	0.0274 (3)
N1	1.4158 (3)	0.70367 (6)	0.98970 (8)	0.0203 (3)
C1	0.8669 (3)	0.78260 (7)	0.88589 (9)	0.0169 (3)
H1	0.7699	0.8205	0.8865	0.020*
C2	1.0633 (3)	0.77121 (7)	0.93827 (9)	0.0176 (3)
H2	1.1025	0.8007	0.9752	0.021*
C3	1.2012 (3)	0.71558 (7)	0.93535 (8)	0.0163 (3)
C4	1.1480 (3)	0.67097 (7)	0.88291 (9)	0.0169 (3)
H4	1.2455	0.6331	0.8825	0.020*
C5	0.9499 (3)	0.68282 (7)	0.83117 (8)	0.0163 (3)
H5	0.9090	0.6528	0.7949	0.020*
C6	0.8102 (3)	0.73879 (7)	0.83215 (8)	0.0143 (3)
C7	0.5974 (3)	0.75381 (7)	0.77687 (8)	0.0155 (3)
C8	0.3759 (3)	0.71808 (7)	0.66929 (9)	0.0193 (3)
H8A	0.4555	0.7434	0.6299	0.023*
H8B	0.2148	0.7399	0.6882	0.023*
C9	0.2977 (3)	0.65561 (7)	0.64066 (9)	0.0183 (3)
H9A	0.1541	0.6605	0.6031	0.022*
H9B	0.2231	0.6305	0.6810	0.022*
Cl1'	0.09925 (8)	0.578510 (18)	0.46926 (2)	0.02083 (10)
O1'	0.0719 (2)	0.53366 (5)	0.30930 (6)	0.0170 (2)
O2'	-0.0224 (2)	0.45272 (5)	0.23673 (6)	0.0203 (2)
O3'	0.9692 (2)	0.56670 (5)	0.00593 (7)	0.0249 (3)
O4'	1.0593 (2)	0.63815 (6)	0.08520 (6)	0.0243 (3)
N1'	0.9309 (3)	0.59294 (6)	0.06415 (7)	0.0176 (3)
C1'	0.3955 (3)	0.49087 (7)	0.13991 (8)	0.0158 (3)
H1'	0.3083	0.4527	0.1291	0.019*
C2'	0.5964 (3)	0.51288 (7)	0.09445 (9)	0.0165 (3)
H2'	0.6497	0.4903	0.0525	0.020*
C3'	0.7174 (3)	0.56882 (7)	0.11200 (8)	0.0150 (3)
C4'	0.6481 (3)	0.60310 (7)	0.17254 (9)	0.0170 (3)
H4'	0.7366	0.6411	0.1831	0.020*
C5'	0.4468 (3)	0.58079 (7)	0.21743 (8)	0.0160 (3)
H5'	0.3941	0.6037	0.2592	0.019*
C6'	0.3209 (3)	0.52453 (7)	0.20138 (8)	0.0139 (3)
C7'	0.1063 (3)	0.49883 (7)	0.24947 (8)	0.0149 (3)
C8'	-0.1343 (3)	0.51265 (7)	0.35872 (8)	0.0168 (3)
H8'1	-0.0718	0.4753	0.3852	0.020*
H8'2	-0.3017	0.5022	0.3314	0.020*
C9'	-0.1878 (3)	0.56394 (8)	0.41134 (9)	0.0211 (3)
H9'1	-0.3454	0.5530	0.4418	0.025*

supplementary materials

H9'2 -0.2334 0.6020 0.3839 0.025*

Atomic displacement parameters (\AA^2)

	U^{11}	U^{22}	U^{33}	U^{12}	U^{13}	U^{23}
C11	0.01863 (19)	0.0242 (2)	0.0213 (2)	-0.00012 (15)	0.00069 (15)	-0.00381 (16)
O1	0.0238 (6)	0.0183 (5)	0.0182 (6)	0.0036 (5)	-0.0087 (5)	-0.0042 (4)
O2	0.0287 (6)	0.0174 (5)	0.0228 (6)	0.0070 (5)	-0.0078 (5)	-0.0007 (5)
O3	0.0304 (7)	0.0329 (7)	0.0229 (7)	-0.0062 (5)	-0.0103 (6)	0.0002 (5)
O4	0.0203 (6)	0.0359 (7)	0.0259 (7)	0.0089 (5)	0.0000 (5)	0.0089 (5)
N1	0.0156 (6)	0.0272 (7)	0.0182 (7)	-0.0039 (6)	-0.0011 (5)	0.0071 (6)
C1	0.0191 (8)	0.0141 (7)	0.0174 (8)	0.0020 (6)	0.0006 (6)	-0.0007 (6)
C2	0.0209 (8)	0.0178 (7)	0.0142 (8)	-0.0020 (6)	0.0001 (6)	-0.0011 (6)
C3	0.0131 (7)	0.0215 (7)	0.0144 (7)	-0.0020 (6)	-0.0004 (6)	0.0050 (6)
C4	0.0165 (7)	0.0150 (7)	0.0191 (8)	0.0026 (6)	0.0021 (6)	0.0020 (6)
C5	0.0199 (8)	0.0142 (7)	0.0148 (7)	-0.0005 (6)	0.0013 (6)	-0.0012 (6)
C6	0.0157 (7)	0.0146 (7)	0.0127 (7)	-0.0006 (6)	0.0015 (6)	0.0016 (6)
C7	0.0179 (7)	0.0155 (7)	0.0131 (7)	-0.0016 (6)	0.0009 (6)	0.0005 (6)
C8	0.0217 (8)	0.0192 (7)	0.0168 (8)	0.0010 (6)	-0.0071 (6)	-0.0008 (6)
C9	0.0152 (7)	0.0209 (7)	0.0189 (8)	-0.0011 (6)	0.0006 (6)	-0.0018 (6)
C11'	0.0243 (2)	0.02306 (19)	0.01513 (19)	-0.00621 (15)	0.00131 (15)	-0.00093 (15)
O1'	0.0212 (6)	0.0176 (5)	0.0123 (5)	-0.0048 (4)	0.0038 (4)	-0.0015 (4)
O2'	0.0235 (6)	0.0157 (5)	0.0216 (6)	-0.0044 (5)	0.0024 (5)	-0.0019 (5)
O3'	0.0313 (7)	0.0233 (6)	0.0203 (6)	0.0061 (5)	0.0107 (5)	0.0013 (5)
O4'	0.0200 (6)	0.0302 (6)	0.0225 (6)	-0.0072 (5)	-0.0001 (5)	0.0030 (5)
N1'	0.0154 (6)	0.0206 (7)	0.0167 (7)	0.0054 (5)	0.0004 (5)	0.0052 (5)
C1'	0.0183 (7)	0.0139 (7)	0.0153 (8)	0.0010 (6)	-0.0024 (6)	-0.0015 (6)
C2'	0.0184 (7)	0.0178 (7)	0.0134 (7)	0.0047 (6)	-0.0009 (6)	-0.0025 (6)
C3'	0.0125 (7)	0.0181 (7)	0.0144 (7)	0.0026 (6)	-0.0001 (6)	0.0048 (6)
C4'	0.0183 (8)	0.0164 (7)	0.0164 (8)	-0.0014 (6)	-0.0022 (6)	0.0002 (6)
C5'	0.0195 (8)	0.0154 (7)	0.0130 (7)	0.0000 (6)	0.0002 (6)	-0.0021 (6)
C6'	0.0146 (7)	0.0145 (7)	0.0127 (7)	0.0014 (6)	-0.0021 (6)	0.0013 (6)
C7'	0.0164 (7)	0.0144 (7)	0.0138 (7)	0.0031 (6)	-0.0026 (6)	0.0006 (6)
C8'	0.0170 (7)	0.0200 (7)	0.0134 (8)	-0.0038 (6)	0.0032 (6)	0.0012 (6)
C9'	0.0172 (8)	0.0261 (8)	0.0202 (8)	0.0018 (6)	0.0014 (6)	-0.0025 (7)

Geometric parameters (\AA , $^\circ$)

C11—C9	1.7971 (16)	C11'—C9'	1.7927 (17)
O1—C7	1.3408 (18)	O1'—C7'	1.3419 (18)
O1—C8	1.4454 (19)	O1'—C8'	1.4416 (18)
O2—C7	1.2043 (18)	O2'—C7'	1.2041 (18)
O3—N1	1.2245 (19)	O3'—N1'	1.2243 (18)
O4—N1	1.2247 (18)	O4'—N1'	1.2258 (17)
N1—C3	1.472 (2)	N1'—C3'	1.4725 (19)
C1—C2	1.382 (2)	C1'—C2'	1.385 (2)
C1—C6	1.393 (2)	C1'—C6'	1.393 (2)
C1—H1	0.9500	C1'—H1'	0.9500
C2—C3	1.383 (2)	C2'—C3'	1.386 (2)

C2—H2	0.9500	C2'—H2'	0.9500
C3—C4	1.385 (2)	C3'—C4'	1.380 (2)
C4—C5	1.381 (2)	C4'—C5'	1.382 (2)
C4—H4	0.9500	C4'—H4'	0.9500
C5—C6	1.393 (2)	C5'—C6'	1.396 (2)
C5—H5	0.9500	C5'—H5'	0.9500
C6—C7	1.490 (2)	C6'—C7'	1.491 (2)
C8—C9	1.498 (2)	C8'—C9'	1.494 (2)
C8—H8A	0.9900	C8'—H8'1	0.9900
C8—H8B	0.9900	C8'—H8'2	0.9900
C9—H9A	0.9900	C9'—H9'1	0.9900
C9—H9B	0.9900	C9'—H9'2	0.9900
C7—O1—C8	117.01 (12)	C7'—O1'—C8'	115.53 (11)
O3—N1—O4	124.40 (14)	O3'—N1'—O4'	124.11 (14)
O3—N1—C3	118.16 (14)	O3'—N1'—C3'	118.11 (13)
O4—N1—C3	117.44 (14)	O4'—N1'—C3'	117.79 (13)
C2—C1—C6	120.41 (14)	C2'—C1'—C6'	120.14 (14)
C2—C1—H1	119.8	C2'—C1'—H1'	119.9
C6—C1—H1	119.8	C6'—C1'—H1'	119.9
C1—C2—C3	118.13 (14)	C1'—C2'—C3'	118.03 (14)
C1—C2—H2	120.9	C1'—C2'—H2'	121.0
C3—C2—H2	120.9	C3'—C2'—H2'	121.0
C2—C3—C4	122.76 (14)	C4'—C3'—C2'	123.12 (14)
C2—C3—N1	118.58 (14)	C4'—C3'—N1'	118.16 (13)
C4—C3—N1	118.65 (14)	C2'—C3'—N1'	118.72 (14)
C5—C4—C3	118.48 (14)	C3'—C4'—C5'	118.36 (14)
C5—C4—H4	120.8	C3'—C4'—H4'	120.8
C3—C4—H4	120.8	C5'—C4'—H4'	120.8
C4—C5—C6	120.04 (14)	C4'—C5'—C6'	120.00 (14)
C4—C5—H5	120.0	C4'—C5'—H5'	120.0
C6—C5—H5	120.0	C6'—C5'—H5'	120.0
C1—C6—C5	120.18 (14)	C1'—C6'—C5'	120.34 (14)
C1—C6—C7	118.00 (13)	C1'—C6'—C7'	118.48 (13)
C5—C6—C7	121.83 (14)	C5'—C6'—C7'	121.17 (14)
O2—C7—O1	124.27 (14)	O2'—C7'—O1'	123.57 (14)
O2—C7—C6	124.32 (14)	O2'—C7'—C6'	124.79 (14)
O1—C7—C6	111.41 (12)	O1'—C7'—C6'	111.64 (12)
O1—C8—C9	107.38 (12)	O1'—C8'—C9'	107.52 (12)
O1—C8—H8A	110.2	O1'—C8'—H8'1	110.2
C9—C8—H8A	110.2	C9'—C8'—H8'1	110.2
O1—C8—H8B	110.2	O1'—C8'—H8'2	110.2
C9—C8—H8B	110.2	C9'—C8'—H8'2	110.2
H8A—C8—H8B	108.5	H8'1—C8'—H8'2	108.5
C8—C9—Cl1	111.97 (11)	C8'—C9'—Cl1'	111.67 (11)
C8—C9—H9A	109.2	C8'—C9'—H9'1	109.3
Cl1—C9—H9A	109.2	Cl1'—C9'—H9'1	109.3
C8—C9—H9B	109.2	C8'—C9'—H9'2	109.3
Cl1—C9—H9B	109.2	Cl1'—C9'—H9'2	109.3
H9A—C9—H9B	107.9	H9'1—C9'—H9'2	107.9

supplementary materials

C6—C1—C2—C3	0.3 (2)	C6'—C1'—C2'—C3'	-0.3 (2)
C1—C2—C3—C4	-0.8 (2)	C1'—C2'—C3'—C4'	0.4 (2)
C1—C2—C3—N1	178.39 (13)	C1'—C2'—C3'—N1'	-179.86 (13)
O3—N1—C3—C2	7.2 (2)	O3'—N1'—C3'—C4'	-171.12 (13)
O4—N1—C3—C2	-171.95 (14)	O4'—N1'—C3'—C4'	8.8 (2)
O3—N1—C3—C4	-173.58 (14)	O3'—N1'—C3'—C2'	9.1 (2)
O4—N1—C3—C4	7.3 (2)	O4'—N1'—C3'—C2'	-170.90 (13)
C2—C3—C4—C5	0.4 (2)	C2'—C3'—C4'—C5'	-0.6 (2)
N1—C3—C4—C5	-178.79 (13)	N1'—C3'—C4'—C5'	179.68 (13)
C3—C4—C5—C6	0.5 (2)	C3'—C4'—C5'—C6'	0.6 (2)
C2—C1—C6—C5	0.5 (2)	C2'—C1'—C6'—C5'	0.4 (2)
C2—C1—C6—C7	-179.66 (14)	C2'—C1'—C6'—C7'	-179.20 (13)
C4—C5—C6—C1	-0.9 (2)	C4'—C5'—C6'—C1'	-0.5 (2)
C4—C5—C6—C7	179.26 (14)	C4'—C5'—C6'—C7'	179.00 (14)
C8—O1—C7—O2	1.3 (2)	C8'—O1'—C7—O2'	-0.4 (2)
C8—O1—C7—C6	-178.60 (13)	C8'—O1'—C7—C6'	179.34 (12)
C1—C6—C7—O2	-2.3 (2)	C1'—C6'—C7—O2'	-5.3 (2)
C5—C6—C7—O2	177.53 (15)	C5'—C6'—C7—O2'	175.17 (15)
C1—C6—C7—O1	177.65 (13)	C1'—C6'—C7—O1'	174.96 (13)
C5—C6—C7—O1	-2.5 (2)	C5'—C6'—C7—O1'	-4.60 (19)
C7—O1—C8—C9	-156.81 (13)	C7'—O1'—C8'—C9'	-166.85 (12)
O1—C8—C9—Cl1	-62.71 (15)	O1'—C8'—C9—Cl1'	-66.44 (15)

Hydrogen-bond geometry (\AA , $^\circ$)

$D—\text{H}\cdots A$	$D—\text{H}$	$\text{H}\cdots A$	$D\cdots A$	$D—\text{H}\cdots A$
C1—H1 \cdots C11 ⁱ	0.95	2.79	3.6234 (18)	147
C2'—H2 \cdots O3 ⁱⁱ	0.95	2.50	3.318 (2)	145
C4'—H4 \cdots O2 ⁱⁱⁱ	0.95	2.40	3.223 (2)	144
C5—H5 \cdots O2 ^{iv}	0.95	2.42	3.205 (2)	140
C5'—H5' \cdots O1'	0.95	2.39	2.7112 (19)	100
C9—H9B \cdots O2 ^v	0.99	2.55	3.526 (2)	167

Symmetry codes: (i) ; (ii) $-x+2, -y+1, -z$; (iii) ; (iv) $-x+1, -y+1, -z+1$; (v) $-x, -y+1, -z+1$.

Fig. 1

