

3-(12-Bromododecyl)-1,5-dimethyl-1*H*-1,5-benzodiazepine-2,4(3*H*,5*H*)-dione

Rchida Dardouri,^a Fouad Ouazzani Chahdi,^a Natalie Saffon,^b El Mokhtar Essassi^a and Seik Weng Ng^{c*}

^aLaboratoire de Chimie Organique Hétérocyclique, Pôle de Compétences Pharmacochimie, Université Mohammed V-Agdal, BP 1014 Avenue Ibn Batout, Rabat, Morocco, ^bService Commun Rayons-X FR2599, Université Paul Sabatier, Bâtiment 2R1, 118 route de Narbonne, Toulouse, France, and ^cDepartment of Chemistry, University of Malaya, 50603 Kuala Lumpur, Malaysia
Correspondence e-mail: seikweng@um.edu.my


Received 6 October 2010; accepted 7 October 2010

Key indicators: single-crystal X-ray study; $T = 293\text{ K}$; mean $\sigma(\text{C}-\text{C}) = 0.003\text{ \AA}$; R factor = 0.039; wR factor = 0.099; data-to-parameter ratio = 26.0.

The seven-membered ring in the title compound, $\text{C}_{23}\text{H}_{35}\text{BrN}_2\text{O}_2$, adopts a boat-shaped conformation (with the C atoms of the fused-ring as the stern and the methine C atom as the prow). The bromododecyl substituent occupies an equatorial position, with the dodecyl chain exhibiting an extended conformation. Weak intermolecular $\text{C}-\text{H}\cdots\text{O}$ hydrogen bonding is present in the crystal structure.


Related literature

For the crystal structure of 1,5-dimethyl-1,5-benzodiazepin-2,4-dione, see: Mondieig *et al.* (2005). For the structure of a similar compound, 3-(6-bromoheptyl)-1,5-dimethyl-1,5-benzodiazepine-2,4-dione, see: Dardouri *et al.* (2010).



Experimental

Crystal data



Monoclinic, $P2_1/n$	$Z = 4$
$a = 7.5971 (1)\text{ \AA}$	Mo $K\alpha$ radiation
$b = 10.5032 (2)\text{ \AA}$	$\mu = 1.82\text{ mm}^{-1}$
$c = 28.7129 (5)\text{ \AA}$	$T = 293\text{ K}$
$\beta = 95.213 (1)^\circ$	$0.30 \times 0.20 \times 0.10\text{ mm}$
$V = 2281.64 (7)\text{ \AA}^3$	

Data collection

Bruker X8 APEXII diffractometer	24581 measured reflections
Absorption correction: multi-scan (<i>SADABS</i> ; Sheldrick, 1996)	6634 independent reflections
$(SADABS$; Sheldrick, 1996)	4370 reflections with $I > 2\sigma(I)$
$T_{\min} = 0.611$, $T_{\max} = 0.839$	$R_{\text{int}} = 0.037$

Refinement

$R[F^2 > 2\sigma(F^2)] = 0.039$	255 parameters
$wR(F^2) = 0.099$	H-atom parameters constrained
$S = 0.99$	$\Delta\rho_{\max} = 0.61\text{ e \AA}^{-3}$
6634 reflections	$\Delta\rho_{\min} = -0.23\text{ e \AA}^{-3}$

Table 1
Hydrogen-bond geometry (\AA , $^\circ$).

$D-\text{H}\cdots A$	$D-\text{H}$	$\text{H}\cdots A$	$D\cdots A$	$D-\text{H}\cdots A$
C2—H2 \cdots O1 ⁱ	0.93	2.52	3.424 (2)	164
C7—H7B \cdots O2 ⁱⁱ	0.96	2.40	3.340 (2)	166
C11—H11C \cdots O1 ⁱⁱ	0.96	2.48	3.407 (2)	164

Symmetry codes: (i) $-x + \frac{1}{2}, y - \frac{1}{2}, -z + \frac{1}{2}$; (ii) $-x - \frac{1}{2}, y - \frac{1}{2}, -z + \frac{1}{2}$.

Data collection: *APEX2* (Bruker, 2008); cell refinement: *SAINT* (Bruker, 2008); data reduction: *SAINT*; program(s) used to solve structure: *SHELXS97* (Sheldrick, 2008); program(s) used to refine structure: *SHELXL97* (Sheldrick, 2008); molecular graphics: *X-SEED* (Barbour, 2001); software used to prepare material for publication: *pubLCIF* (Westrip, 2010).

We thank Université Mohammed V-Agdal and the University of Malaya for supporting this study.

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: XU5050).

References

- Barbour, L. J. (2001). *J. Supramol. Chem.* **1**, 189–191.
- Bruker (2008). *APEX2* and *SAINT*. Bruker AXS Inc., Madison, Wisconsin, USA.
- Dardouri, R., Ouazzani Chahdi, F., Saffon, N., Essassi, E. M. & Ng, S. W. (2010). *Acta Cryst. E66*, o2804.
- Mondieig, M., Negrer, Ph., Léger, J. M., Benali, B., Lazar, Z., Elassyry, A., Jarmouni, C., Lahrissi, B. & Massoui, M. (2005). *Anal. Sci. X-Ray Struct. Anal. Online*, **21**, x145–x146.
- Sheldrick, G. M. (1996). *SADABS*. University of Göttingen, Germany.
- Sheldrick, G. M. (2008). *Acta Cryst. A64*, 112–122.
- Westrip, S. P. (2010). *J. Appl. Cryst.* **43**, 920–925.

supplementary materials

Acta Cryst. (2010). E66, o2805 [doi:10.1107/S1600536810040134]

3-(12-Bromododecyl)-1,5-dimethyl-1*H*-1,5-benzodiazepine-2,4(3*H*,5*H*)-dione

R. Dardouri, F. Ouazzani Chahdi, N. Saffon, E. M. Essassi and S. W. Ng

Comment

The methylene part of 1,5-dimethyl-1,5-benzodiazepine-2,4-dione is relatively acidic, and one proton can be abstracted by using potassium *t*-butoxide; the resulting carbanion can undergo a nucleophilic substitution with a dibromoalkane to form 3-substituted derivatives. In this study, the compound is reacted with 1,12-dibromododecane the title compound (Scheme I, Fig. 1).

Experimental

To a solution of the potassium *t*-butoxide (0.42 g, 3.6 mmol) in DMF (15 ml) was added 1,5-dimethyl-1,5-benzodiazepine-2,4-dione (0.50 g, 2.4 mmol) and 1,12-dibromododecane (0.94 g, 2.88 mmol). Stirring was continued for 24 h. The reaction was monitored by thin layer chromatography. The mixture was filtered and the solution evaporated to give colorless crystals.

Refinement

Carbon-bound H-atoms were placed in calculated positions (C—H 0.93–0.97 Å) and were included in the refinement in the riding model approximation, with $U(\text{H})$ set to 1.2–1.5 $U_{\text{eq}}(\text{C})$.

Figures




Fig. 1. Thermal ellipsoid plot (Barbour, 2001) of $\text{C}_{23}\text{H}_{35}\text{BrN}_2\text{O}_2$ at the 50% probability level; hydrogen atoms are drawn as arbitrary radius.

3-(12-Bromododecyl)-1,5-dimethyl-1*H*-1,5-benzodiazepine- 2,4(3*H*,5*H*)-dione

Crystal data

$\text{C}_{23}\text{H}_{35}\text{BrN}_2\text{O}_2$	$F(000) = 952$
$M_r = 451.44$	$D_x = 1.314 \text{ Mg m}^{-3}$
Monoclinic, $P2_1/n$	Mo $K\alpha$ radiation, $\lambda = 0.71073 \text{ \AA}$
Hall symbol: -P 2yn	Cell parameters from 5655 reflections
$a = 7.5971 (1) \text{ \AA}$	$\theta = 2.4\text{--}25.4^\circ$
$b = 10.5032 (2) \text{ \AA}$	$\mu = 1.82 \text{ mm}^{-1}$
$c = 28.7129 (5) \text{ \AA}$	$T = 293 \text{ K}$
$\beta = 95.213 (1)^\circ$	Prism, colorless

supplementary materials

$V = 2281.64 (7) \text{ \AA}^3$

$Z = 4$

$0.30 \times 0.20 \times 0.10 \text{ mm}$

Data collection

Bruker X8 APEXII diffractometer	6634 independent reflections
Radiation source: fine-focus sealed tube graphite	4370 reflections with $I > 2\sigma(I)$
φ and ω scans	$R_{\text{int}} = 0.037$
Absorption correction: multi-scan (<i>SADABS</i> ; Sheldrick, 1996)	$\theta_{\text{max}} = 30.0^\circ, \theta_{\text{min}} = 2.7^\circ$
$T_{\text{min}} = 0.611, T_{\text{max}} = 0.839$	$h = -10 \rightarrow 10$
24581 measured reflections	$k = -14 \rightarrow 12$
	$l = -40 \rightarrow 40$

Refinement

Refinement on F^2	Primary atom site location: structure-invariant direct methods
Least-squares matrix: full	Secondary atom site location: difference Fourier map
$R[F^2 > 2\sigma(F^2)] = 0.039$	Hydrogen site location: inferred from neighbouring sites
$wR(F^2) = 0.099$	H-atom parameters constrained
$S = 0.99$	$w = 1/[\sigma^2(F_o^2) + (0.0518P)^2]$ where $P = (F_o^2 + 2F_c^2)/3$
6634 reflections	$(\Delta/\sigma)_{\text{max}} = 0.001$
255 parameters	$\Delta\rho_{\text{max}} = 0.61 \text{ e \AA}^{-3}$
0 restraints	$\Delta\rho_{\text{min}} = -0.23 \text{ e \AA}^{-3}$

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (\AA^2)

	x	y	z	$U_{\text{iso}}^*/U_{\text{eq}}$
Br1	1.65169 (3)	1.558652 (19)	0.586300 (7)	0.04634 (9)
N1	0.0090 (2)	0.57047 (12)	0.26467 (5)	0.0294 (3)
N2	-0.32138 (19)	0.57806 (12)	0.30691 (5)	0.0291 (3)
O1	0.08035 (17)	0.77902 (11)	0.26153 (4)	0.0368 (3)
O2	-0.35919 (17)	0.78733 (11)	0.32234 (5)	0.0411 (3)
C1	-0.0532 (2)	0.46396 (15)	0.28941 (6)	0.0276 (4)
C2	0.0475 (2)	0.35221 (16)	0.29190 (7)	0.0350 (4)
H2	0.1551	0.3497	0.2788	0.042*
C3	-0.0126 (3)	0.24597 (17)	0.31369 (7)	0.0411 (5)
H3	0.0549	0.1720	0.3154	0.049*
C4	-0.1723 (3)	0.24865 (17)	0.33297 (7)	0.0407 (5)
H4	-0.2131	0.1762	0.3472	0.049*
C5	-0.2713 (3)	0.35867 (16)	0.33120 (7)	0.0347 (4)
H5	-0.3781	0.3605	0.3447	0.042*
C6	-0.2126 (2)	0.46761 (15)	0.30930 (6)	0.0266 (4)
C7	0.0742 (3)	0.54706 (17)	0.21882 (6)	0.0355 (4)

H7A	0.0659	0.6240	0.2007	0.053*
H7B	0.0041	0.4819	0.2027	0.053*
H7C	0.1953	0.5200	0.2231	0.053*
C8	0.0180 (2)	0.69084 (15)	0.28235 (6)	0.0285 (4)
C9	-0.0598 (2)	0.70942 (15)	0.32913 (6)	0.0273 (4)
H9	-0.0165	0.6412	0.3505	0.033*
C10	-0.2600 (2)	0.69690 (16)	0.31965 (6)	0.0292 (4)
C11	-0.5119 (2)	0.56279 (17)	0.29472 (7)	0.0350 (4)
H11A	-0.5592	0.6397	0.2805	0.053*
H11B	-0.5690	0.5454	0.3225	0.053*
H11C	-0.5320	0.4933	0.2732	0.053*
C12	-0.0055 (2)	0.83761 (16)	0.35126 (7)	0.0327 (4)
H12A	-0.0200	0.9037	0.3276	0.039*
H12B	-0.0823	0.8576	0.3755	0.039*
C13	0.1849 (2)	0.83595 (15)	0.37232 (6)	0.0329 (4)
H13A	0.2597	0.8092	0.3485	0.039*
H13B	0.1965	0.7732	0.3972	0.039*
C14	0.2511 (2)	0.96386 (16)	0.39196 (7)	0.0328 (4)
H14A	0.2437	1.0262	0.3669	0.039*
H14B	0.1745	0.9921	0.4152	0.039*
C15	0.4398 (3)	0.95858 (15)	0.41418 (7)	0.0335 (4)
H15A	0.5142	0.9229	0.3918	0.040*
H15B	0.4446	0.9015	0.4408	0.040*
C16	0.5151 (3)	1.08722 (16)	0.43040 (7)	0.0348 (4)
H16A	0.5133	1.1439	0.4037	0.042*
H16B	0.4398	1.1239	0.4524	0.042*
C17	0.7028 (3)	1.07876 (15)	0.45343 (7)	0.0354 (4)
H17A	0.7779	1.0431	0.4312	0.043*
H17B	0.7045	1.0206	0.4797	0.043*
C18	0.7796 (3)	1.20563 (16)	0.47064 (7)	0.0365 (4)
H18A	0.7819	1.2629	0.4442	0.044*
H18B	0.7025	1.2427	0.4921	0.044*
C19	0.9648 (3)	1.19565 (16)	0.49503 (7)	0.0383 (4)
H19A	1.0418	1.1593	0.4734	0.046*
H19B	0.9625	1.1373	0.5212	0.046*
C20	1.0436 (3)	1.32136 (16)	0.51308 (7)	0.0362 (4)
H20A	1.0439	1.3808	0.4872	0.043*
H20B	0.9696	1.3568	0.5357	0.043*
C21	1.2304 (3)	1.30689 (16)	0.53584 (7)	0.0370 (4)
H21A	1.3013	1.2651	0.5139	0.044*
H21B	1.2278	1.2516	0.5628	0.044*
C22	1.3211 (3)	1.43160 (15)	0.55161 (7)	0.0365 (4)
H22A	1.3216	1.4894	0.5253	0.044*
H22B	1.2569	1.4718	0.5753	0.044*
C23	1.5070 (3)	1.40468 (18)	0.57104 (7)	0.0409 (5)
H23A	1.5643	1.3542	0.5485	0.049*
H23B	1.5038	1.3539	0.5992	0.049*

supplementary materials

Atomic displacement parameters (\AA^2)

	U^{11}	U^{22}	U^{33}	U^{12}	U^{13}	U^{23}
Br1	0.05160 (15)	0.04100 (12)	0.04792 (14)	-0.01138 (9)	0.01260 (10)	-0.00678 (9)
N1	0.0297 (8)	0.0275 (7)	0.0323 (8)	-0.0030 (6)	0.0092 (6)	-0.0026 (6)
N2	0.0229 (7)	0.0252 (7)	0.0392 (8)	0.0003 (5)	0.0030 (6)	-0.0004 (6)
O1	0.0387 (7)	0.0305 (6)	0.0421 (7)	-0.0073 (5)	0.0089 (6)	0.0033 (5)
O2	0.0327 (7)	0.0307 (7)	0.0598 (9)	0.0082 (6)	0.0037 (6)	-0.0064 (6)
C1	0.0281 (9)	0.0249 (8)	0.0300 (9)	-0.0010 (7)	0.0030 (7)	-0.0014 (6)
C2	0.0304 (10)	0.0325 (9)	0.0428 (11)	0.0046 (7)	0.0072 (8)	-0.0037 (8)
C3	0.0466 (12)	0.0276 (9)	0.0492 (12)	0.0100 (8)	0.0057 (9)	0.0017 (8)
C4	0.0507 (12)	0.0294 (9)	0.0433 (11)	-0.0005 (8)	0.0114 (9)	0.0078 (8)
C5	0.0340 (10)	0.0311 (9)	0.0405 (10)	-0.0021 (7)	0.0116 (8)	0.0013 (7)
C6	0.0257 (9)	0.0238 (8)	0.0305 (9)	0.0006 (6)	0.0029 (7)	-0.0015 (6)
C7	0.0343 (10)	0.0403 (10)	0.0331 (10)	-0.0048 (8)	0.0097 (8)	-0.0036 (8)
C8	0.0230 (9)	0.0274 (8)	0.0348 (9)	-0.0007 (7)	0.0002 (7)	-0.0007 (7)
C9	0.0255 (9)	0.0239 (8)	0.0323 (9)	0.0008 (6)	0.0015 (7)	-0.0024 (6)
C10	0.0286 (9)	0.0275 (8)	0.0317 (9)	0.0023 (7)	0.0041 (7)	-0.0009 (7)
C11	0.0254 (9)	0.0361 (9)	0.0431 (11)	-0.0003 (8)	0.0002 (8)	-0.0007 (8)
C12	0.0322 (10)	0.0257 (8)	0.0402 (10)	0.0003 (7)	0.0031 (8)	-0.0054 (7)
C13	0.0360 (10)	0.0253 (8)	0.0371 (10)	-0.0006 (7)	0.0017 (8)	-0.0047 (7)
C14	0.0375 (10)	0.0258 (8)	0.0351 (10)	-0.0005 (7)	0.0034 (8)	-0.0030 (7)
C15	0.0396 (11)	0.0257 (9)	0.0345 (10)	-0.0019 (7)	0.0000 (8)	-0.0036 (7)
C16	0.0414 (11)	0.0252 (9)	0.0378 (10)	-0.0010 (7)	0.0038 (8)	-0.0046 (7)
C17	0.0426 (11)	0.0263 (9)	0.0366 (10)	-0.0045 (7)	-0.0009 (8)	-0.0050 (7)
C18	0.0429 (11)	0.0252 (9)	0.0412 (10)	-0.0027 (8)	0.0024 (9)	-0.0049 (7)
C19	0.0449 (11)	0.0287 (9)	0.0400 (11)	-0.0014 (8)	-0.0025 (9)	-0.0040 (8)
C20	0.0419 (11)	0.0281 (9)	0.0384 (10)	-0.0033 (8)	0.0028 (8)	-0.0039 (7)
C21	0.0454 (11)	0.0271 (9)	0.0377 (10)	-0.0014 (8)	-0.0006 (9)	-0.0021 (7)
C22	0.0435 (11)	0.0272 (9)	0.0384 (10)	-0.0023 (8)	0.0023 (9)	-0.0027 (7)
C23	0.0442 (12)	0.0322 (9)	0.0457 (11)	-0.0041 (8)	0.0003 (9)	-0.0067 (8)

Geometric parameters (\AA , $^\circ$)

Br1—C23	1.9818 (19)	C13—C14	1.525 (2)
N1—C8	1.362 (2)	C13—H13A	0.9700
N1—C1	1.428 (2)	C13—H13B	0.9700
N1—C7	1.469 (2)	C14—C15	1.516 (3)
N2—C10	1.371 (2)	C14—H14A	0.9700
N2—C6	1.423 (2)	C14—H14B	0.9700
N2—C11	1.466 (2)	C15—C16	1.523 (2)
O1—C8	1.221 (2)	C15—H15A	0.9700
O2—C10	1.219 (2)	C15—H15B	0.9700
C1—C6	1.386 (2)	C16—C17	1.519 (3)
C1—C2	1.399 (2)	C16—H16A	0.9700
C2—C3	1.377 (3)	C16—H16B	0.9700
C2—H2	0.9300	C17—C18	1.519 (2)
C3—C4	1.378 (3)	C17—H17A	0.9700

C3—H3	0.9300	C17—H17B	0.9700
C4—C5	1.377 (3)	C18—C19	1.517 (3)
C4—H4	0.9300	C18—H18A	0.9700
C5—C6	1.398 (2)	C18—H18B	0.9700
C5—H5	0.9300	C19—C20	1.521 (2)
C7—H7A	0.9600	C19—H19A	0.9700
C7—H7B	0.9600	C19—H19B	0.9700
C7—H7C	0.9600	C20—C21	1.515 (3)
C8—C9	1.528 (2)	C20—H20A	0.9700
C9—C10	1.526 (2)	C20—H20B	0.9700
C9—C12	1.529 (2)	C21—C22	1.529 (2)
C9—H9	0.9800	C21—H21A	0.9700
C11—H11A	0.9600	C21—H21B	0.9700
C11—H11B	0.9600	C22—C23	1.497 (3)
C11—H11C	0.9600	C22—H22A	0.9700
C12—C13	1.516 (3)	C22—H22B	0.9700
C12—H12A	0.9700	C23—H23A	0.9700
C12—H12B	0.9700	C23—H23B	0.9700
C8—N1—C1	123.32 (14)	C15—C14—C13	113.01 (14)
C8—N1—C7	118.69 (14)	C15—C14—H14A	109.0
C1—N1—C7	117.93 (13)	C13—C14—H14A	109.0
C10—N2—C6	123.20 (14)	C15—C14—H14B	109.0
C10—N2—C11	117.90 (14)	C13—C14—H14B	109.0
C6—N2—C11	118.61 (13)	H14A—C14—H14B	107.8
C6—C1—C2	119.76 (15)	C14—C15—C16	114.23 (14)
C6—C1—N1	121.61 (15)	C14—C15—H15A	108.7
C2—C1—N1	118.59 (15)	C16—C15—H15A	108.7
C3—C2—C1	120.02 (17)	C14—C15—H15B	108.7
C3—C2—H2	120.0	C16—C15—H15B	108.7
C1—C2—H2	120.0	H15A—C15—H15B	107.6
C2—C3—C4	120.41 (17)	C17—C16—C15	113.12 (14)
C2—C3—H3	119.8	C17—C16—H16A	109.0
C4—C3—H3	119.8	C15—C16—H16A	109.0
C5—C4—C3	120.00 (17)	C17—C16—H16B	109.0
C5—C4—H4	120.0	C15—C16—H16B	109.0
C3—C4—H4	120.0	H16A—C16—H16B	107.8
C4—C5—C6	120.50 (17)	C16—C17—C18	114.08 (14)
C4—C5—H5	119.7	C16—C17—H17A	108.7
C6—C5—H5	119.7	C18—C17—H17A	108.7
C1—C6—C5	119.30 (15)	C16—C17—H17B	108.7
C1—C6—N2	121.91 (14)	C18—C17—H17B	108.7
C5—C6—N2	118.76 (15)	H17A—C17—H17B	107.6
N1—C7—H7A	109.5	C19—C18—C17	113.72 (15)
N1—C7—H7B	109.5	C19—C18—H18A	108.8
H7A—C7—H7B	109.5	C17—C18—H18A	108.8
N1—C7—H7C	109.5	C19—C18—H18B	108.8
H7A—C7—H7C	109.5	C17—C18—H18B	108.8
H7B—C7—H7C	109.5	H18A—C18—H18B	107.7
O1—C8—N1	122.01 (16)	C18—C19—C20	114.64 (15)

supplementary materials

O1—C8—C9	122.12 (15)	C18—C19—H19A	108.6
N1—C8—C9	115.85 (14)	C20—C19—H19A	108.6
C10—C9—C8	106.92 (14)	C18—C19—H19B	108.6
C10—C9—C12	112.15 (14)	C20—C19—H19B	108.6
C8—C9—C12	111.47 (14)	H19A—C19—H19B	107.6
C10—C9—H9	108.7	C21—C20—C19	112.58 (15)
C8—C9—H9	108.7	C21—C20—H20A	109.1
C12—C9—H9	108.7	C19—C20—H20A	109.1
O2—C10—N2	121.92 (16)	C21—C20—H20B	109.1
O2—C10—C9	122.23 (15)	C19—C20—H20B	109.1
N2—C10—C9	115.83 (14)	H20A—C20—H20B	107.8
N2—C11—H11A	109.5	C20—C21—C22	114.90 (15)
N2—C11—H11B	109.5	C20—C21—H21A	108.5
H11A—C11—H11B	109.5	C22—C21—H21A	108.5
N2—C11—H11C	109.5	C20—C21—H21B	108.5
H11A—C11—H11C	109.5	C22—C21—H21B	108.5
H11B—C11—H11C	109.5	H21A—C21—H21B	107.5
C13—C12—C9	111.55 (14)	C23—C22—C21	109.44 (14)
C13—C12—H12A	109.3	C23—C22—H22A	109.8
C9—C12—H12A	109.3	C21—C22—H22A	109.8
C13—C12—H12B	109.3	C23—C22—H22B	109.8
C9—C12—H12B	109.3	C21—C22—H22B	109.8
H12A—C12—H12B	108.0	H22A—C22—H22B	108.2
C12—C13—C14	114.09 (14)	C22—C23—Br1	114.43 (12)
C12—C13—H13A	108.7	C22—C23—H23A	108.7
C14—C13—H13A	108.7	Br1—C23—H23A	108.7
C12—C13—H13B	108.7	C22—C23—H23B	108.7
C14—C13—H13B	108.7	Br1—C23—H23B	108.7
H13A—C13—H13B	107.6	H23A—C23—H23B	107.6
C8—N1—C1—C6	50.6 (2)	N1—C8—C9—C10	-70.41 (18)
C7—N1—C1—C6	-132.37 (17)	O1—C8—C9—C12	-15.1 (2)
C8—N1—C1—C2	-131.72 (18)	N1—C8—C9—C12	166.69 (15)
C7—N1—C1—C2	45.3 (2)	C6—N2—C10—O2	-171.56 (16)
C6—C1—C2—C3	0.6 (3)	C11—N2—C10—O2	2.2 (3)
N1—C1—C2—C3	-177.09 (17)	C6—N2—C10—C9	9.8 (2)
C1—C2—C3—C4	0.3 (3)	C11—N2—C10—C9	-176.45 (15)
C2—C3—C4—C5	-1.1 (3)	C8—C9—C10—O2	-111.04 (18)
C3—C4—C5—C6	1.0 (3)	C12—C9—C10—O2	11.4 (2)
C2—C1—C6—C5	-0.7 (3)	C8—C9—C10—N2	67.61 (18)
N1—C1—C6—C5	176.96 (16)	C12—C9—C10—N2	-169.93 (15)
C2—C1—C6—N2	-178.75 (16)	C10—C9—C12—C13	164.64 (15)
N1—C1—C6—N2	-1.1 (3)	C8—C9—C12—C13	-75.51 (19)
C4—C5—C6—C1	-0.1 (3)	C9—C12—C13—C14	175.98 (15)
C4—C5—C6—N2	178.02 (17)	C12—C13—C14—C15	178.16 (16)
C10—N2—C6—C1	-51.4 (2)	C13—C14—C15—C16	175.04 (15)
C11—N2—C6—C1	134.90 (17)	C14—C15—C16—C17	178.80 (16)
C10—N2—C6—C5	130.52 (18)	C15—C16—C17—C18	-179.03 (16)
C11—N2—C6—C5	-43.2 (2)	C16—C17—C18—C19	178.06 (17)
C1—N1—C8—O1	176.05 (16)	C17—C18—C19—C20	-179.30 (17)

supplementary materials

C7—N1—C8—O1	−0.9 (3)	C18—C19—C20—C21	−178.30 (16)
C1—N1—C8—C9	−5.7 (2)	C19—C20—C21—C22	175.93 (17)
C7—N1—C8—C9	177.33 (15)	C20—C21—C22—C23	−177.00 (16)
O1—C8—C9—C10	107.82 (18)	C21—C22—C23—Br1	173.24 (13)

Hydrogen-bond geometry (Å, °)

D—H···A	D—H	H···A	D···A	D—H···A
C2—H2···O1 ⁱ	0.93	2.52	3.424 (2)	164
C7—H7B···O2 ⁱⁱ	0.96	2.40	3.340 (2)	166
C11—H11C···O1 ⁱⁱ	0.96	2.48	3.407 (2)	164

Symmetry codes: (i) $-x+1/2, y-1/2, -z+1/2$; (ii) $-x-1/2, y-1/2, -z+1/2$.

supplementary materials

Fig. 1

