

Acta Crystallographica Section E

Structure Reports

Online

ISSN 1600-5368

Ethyl 4-[(3,5-di-*tert*-butyl-2-hydroxy-benzylidene)amino]benzoate

Raied Mustafa Shakir, Azhar Ariffin and Seik Weng Ng*

Department of Chemistry, University of Malaya, 50603 Kuala Lumpur, Malaysia
Correspondence e-mail: seikweng@um.edu.my


Received 23 September 2010; accepted 14 October 2010

Key indicators: single-crystal X-ray study; $T = 100$ K; mean $\sigma(\text{C}-\text{C}) = 0.003$ Å;
R factor = 0.053; wR factor = 0.142; data-to-parameter ratio = 18.9.

The title compound, a Schiff base, $\text{C}_{24}\text{H}_{31}\text{NO}_3$, has a substituted aromatic ring at both ends of the azomethine linkage and these make a dihedral angle of $24.9(1)^\circ$. There is an intramolecular hydrogen bond between the hydroxy group (donor) and the N atom of the azomethine linkage.

Related literature

For the use of the methyl ester analog of the title compound as a second-harmonic generation material, see: Sliwa *et al.* (2008).



Experimental

Crystal data

 $\text{C}_{24}\text{H}_{31}\text{NO}_3$ $M_r = 381.50$

Monoclinic, $P2_1/c$
 $a = 18.4789(18)$ Å
 $b = 10.7194(11)$ Å
 $c = 10.7768(10)$ Å
 $\beta = 97.437(2)^\circ$
 $V = 2116.7(4)$ Å³

$Z = 4$
Mo $K\alpha$ radiation
 $\mu = 0.08$ mm⁻¹
 $T = 100$ K
 $0.30 \times 0.05 \times 0.05$ mm

Data collection

Bruker SMART APEX
diffractometer
19941 measured reflections

4855 independent reflections
3123 reflections with $I > 2\sigma(I)$
 $R_{\text{int}} = 0.065$

Refinement

$R[F^2 > 2\sigma(F^2)] = 0.053$
 $wR(F^2) = 0.142$
 $S = 1.01$
4855 reflections
257 parameters
1 restraint

H atoms treated by a mixture of
independent and constrained
refinement
 $\Delta\rho_{\text{max}} = 0.64$ e Å⁻³
 $\Delta\rho_{\text{min}} = -0.26$ e Å⁻³

Table 1

Hydrogen-bond geometry (Å, °).

$D-H\cdots A$	$D-H$	$H\cdots A$	$D\cdots A$	$D-H\cdots A$
$\text{O1}-\text{H1}\cdots\text{N1}$	0.87 (1)	1.80 (2)	2.609 (2)	154 (3)

Data collection: *APEX2* (Bruker, 2009); cell refinement: *SAINTE* (Bruker, 2009); data reduction: *SAINTE*; program(s) used to solve structure: *SHELXS97* (Sheldrick, 2008); program(s) used to refine structure: *SHELXL97* (Sheldrick, 2008); molecular graphics: *X-SEED* (Barbour, 2001); software used to prepare material for publication: *publCIF* (Westrip, 2010).

We thank the University of Malaya for supporting this study.

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: FL2320).

References

- Barbour, L. J. (2001). *J. Supramol. Chem.* **1**, 189–191.
Bruker (2009). *APEX2* and *SAINTE*. Bruker AXS Inc., Madison, Wisconsin, USA.
Sheldrick, G. M. (2008). *Acta Cryst.* **A64**, 112–122.
Sliwa, M., Spangenberg, A., Metivier, R., Letard, S., Nakatani, K. & Yu, P. (2008). *Res. Chem. Intermed.* **34**, 181–190.
Westrip, S. P. (2010). *J. Appl. Cryst.* **43**, 920–925.

supplementary materials

Acta Cryst. (2010). E66, o2915 [doi:10.1107/S1600536810041383]

Ethyl 4-[(3,5-di-*tert*-butyl-2-hydroxybenzylidene)amino]benzoate

R. M. Shakir, A. Ariffin and S. W. Ng

Comment

The Schiff base, methyl 4-(di-3,5-*tert*-butyl-2-hydroxybenzylideneamino)benzoate, is a material suitable for second-harmonic generation as it has electron-donating and electron-withdrawing components that are critical for the manifestation of a permanent dipole (Sliwa *et al.*, 2008). Replacing the methyl group with an ethyl moiety leads to (I), an intensely orange-colored compound (Scheme I, Fig. 1) that crystallizes in a centric space group and is, therefore, not suitable as an SHG material. The azomethine bond has an *E*-configuration; the two aromatic rings are aligned at 24.9 (1)°. The compound is neutral as the hydroxy group bears a hydrogen atom which is a donor in an intra-molecular H bond to the azomethine nitrogen atom (Table 1). There are no important intermolecular contacts; on the other hand, the compound appears to pack in such a way as to accommodate the bulky *t*-butyl groups as far as possible (Fig. 2).

Experimental

Ethyl 4-aminobenzoate (0.35 g) dissolved in ethanol (5 ml) was added to 3,5-di-*tert*-butyl-2-hydroxybenzaldehyde (0.5 g) dissolved in ethanol (20 ml). Several drops of acetic acid were added. The solution was heated for 3 h. The solvent was evaporated and the product recrystallized from ethanol to yield orange prisms in 80% yield that were suitable for data collection.

Refinement

Carbon-bound H-atoms were placed in calculated positions (C—H 0.95–0.98 Å) and were included in the refinement in the riding model approximation, with $U_{\text{iso}}(\text{H})$ set to 1.2–1.5 $U_{\text{eq}}(\text{C})$.

The hydroxy H-atom was located in a difference Fourier map, and was refined with the O—H distance restrained to 0.84±0.01 Å; its temperature factor was refined.

In the final difference Fourier map, the largest peak was in the vicinity of an aromatic H-atom.

Figures




Fig. 1. Thermal ellipsoid plot (Barbour, 2001) of the $\text{C}_{24}\text{H}_{31}\text{NO}_3$ at the 70% probability level; hydrogen atoms are drawn as spheres of arbitrary radius.




Fig. 2. Van der Waals packing of two adjacent molecules.

Ethyl 4-[(di-3,5-*tert*-butyl-2-hydroxybenzylidene)amino]benzoate

Crystal data

$C_{24}H_{31}NO_3$	$F(000) = 824$
$M_r = 381.50$	$D_x = 1.197 \text{ Mg m}^{-3}$
Monoclinic, $P2_1/c$	Mo $K\alpha$ radiation, $\lambda = 0.71073 \text{ \AA}$
$a = 18.4789 (18) \text{ \AA}$	Cell parameters from 2280 reflections
$b = 10.7194 (11) \text{ \AA}$	$\theta = 2.2\text{--}23.8^\circ$
$c = 10.7768 (10) \text{ \AA}$	$\mu = 0.08 \text{ mm}^{-1}$
$\beta = 97.437 (2)^\circ$	$T = 100 \text{ K}$
$V = 2116.7 (4) \text{ \AA}^3$	Prism, orange
$Z = 4$	$0.30 \times 0.05 \times 0.05 \text{ mm}$

Data collection

Bruker SMART APEX diffractometer	3123 reflections with $I > 2\sigma(I)$
Radiation source: fine-focus sealed tube	$R_{\text{int}} = 0.065$
graphite	$\theta_{\text{max}} = 27.5^\circ$, $\theta_{\text{min}} = 2.2^\circ$
ω scans	$h = -24 \rightarrow 23$
19941 measured reflections	$k = -13 \rightarrow 13$
4855 independent reflections	$l = -13 \rightarrow 14$

Refinement

Refinement on F^2	Primary atom site location: structure-invariant direct methods
Least-squares matrix: full	Secondary atom site location: difference Fourier map
$R[F^2 > 2\sigma(F^2)] = 0.053$	Hydrogen site location: inferred from neighbouring sites
$wR(F^2) = 0.142$	H atoms treated by a mixture of independent and constrained refinement
$S = 1.01$	$w = 1/[\sigma^2(F_o^2) + (0.0558P)^2 + 1.1473P]$
4855 reflections	where $P = (F_o^2 + 2F_c^2)/3$
257 parameters	$(\Delta/\sigma)_{\text{max}} = 0.001$
1 restraint	$\Delta\rho_{\text{max}} = 0.64 \text{ e \AA}^{-3}$
	$\Delta\rho_{\text{min}} = -0.25 \text{ e \AA}^{-3}$

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (\AA^2)

x	y	z	$U_{\text{iso}}^*/U_{\text{eq}}$
-----	-----	-----	----------------------------------

O1	0.78175 (8)	0.89233 (13)	0.62825 (14)	0.0252 (3)
H1	0.7466 (13)	0.885 (3)	0.567 (2)	0.086 (12)*
O2	0.42198 (8)	0.93692 (14)	0.03223 (14)	0.0291 (4)
O3	0.43875 (7)	0.73467 (13)	-0.01081 (13)	0.0237 (3)
N1	0.67954 (9)	0.79668 (16)	0.46645 (16)	0.0222 (4)
C1	0.78970 (11)	0.77878 (18)	0.68363 (19)	0.0203 (4)
C2	0.84179 (10)	0.76051 (18)	0.78972 (18)	0.0186 (4)
C3	0.85050 (10)	0.63866 (18)	0.83515 (18)	0.0180 (4)
H3	0.8862	0.6243	0.9054	0.022*
C4	0.80999 (10)	0.53616 (18)	0.78384 (18)	0.0183 (4)
C5	0.75592 (10)	0.56031 (19)	0.68469 (18)	0.0199 (4)
H5	0.7253	0.4941	0.6509	0.024*
C6	0.74560 (10)	0.67938 (19)	0.63368 (18)	0.0201 (4)
C7	0.94204 (12)	0.9088 (2)	0.7566 (2)	0.0290 (5)
H7A	0.9137	0.9359	0.6780	0.044*
H7B	0.9735	0.8388	0.7402	0.044*
H7C	0.9722	0.9782	0.7930	0.044*
C8	0.88965 (11)	0.86740 (19)	0.84883 (19)	0.0219 (4)
C9	0.84263 (13)	0.9784 (2)	0.8811 (2)	0.0295 (5)
H9A	0.8124	1.0072	0.8051	0.044*
H9B	0.8744	1.0464	0.9158	0.044*
H9C	0.8112	0.9521	0.9428	0.044*
C10	0.93590 (13)	0.8274 (2)	0.9709 (2)	0.0304 (5)
H10A	0.9670	0.7569	0.9542	0.046*
H10B	0.9037	0.8024	1.0320	0.046*
H10C	0.9665	0.8974	1.0044	0.046*
C11	0.82432 (11)	0.40232 (18)	0.82969 (18)	0.0197 (4)
C12	0.75683 (11)	0.3511 (2)	0.8825 (2)	0.0254 (5)
H12A	0.7465	0.4030	0.9529	0.038*
H12B	0.7662	0.2652	0.9111	0.038*
H12C	0.7148	0.3526	0.8169	0.038*
C13	0.84080 (12)	0.3212 (2)	0.7192 (2)	0.0259 (5)
H13A	0.8838	0.3540	0.6858	0.039*
H13B	0.7988	0.3225	0.6535	0.039*
H13C	0.8502	0.2352	0.7479	0.039*
C14	0.88904 (11)	0.3930 (2)	0.9334 (2)	0.0247 (5)
H14A	0.9330	0.4251	0.9025	0.037*
H14B	0.8966	0.3056	0.9583	0.037*
H14C	0.8789	0.4423	1.0057	0.037*
C15	0.68922 (11)	0.69415 (19)	0.52586 (19)	0.0219 (4)
H15	0.6589	0.6250	0.4995	0.026*
C16	0.62487 (10)	0.8028 (2)	0.36008 (19)	0.0213 (4)
C17	0.59190 (11)	0.9167 (2)	0.3314 (2)	0.0235 (5)
H17	0.6062	0.9876	0.3815	0.028*
C18	0.53831 (11)	0.9279 (2)	0.23032 (19)	0.0228 (5)
H18	0.5157	1.0064	0.2116	0.027*
C19	0.51707 (10)	0.82475 (19)	0.15535 (19)	0.0202 (4)
C20	0.55214 (11)	0.7112 (2)	0.18075 (19)	0.0214 (4)
H20	0.5391	0.6411	0.1286	0.026*

supplementary materials

C21	0.60615 (11)	0.7002 (2)	0.28224 (19)	0.0234 (5)
H21	0.6305	0.6228	0.2990	0.028*
C22	0.45526 (11)	0.84073 (19)	0.05346 (19)	0.0209 (4)
C23	0.37799 (11)	0.7442 (2)	-0.1100 (2)	0.0265 (5)
H23A	0.3908	0.8002	-0.1768	0.032*
H23B	0.3347	0.7788	-0.0767	0.032*
C24	0.36159 (12)	0.6163 (2)	-0.1613 (2)	0.0281 (5)
H24A	0.3206	0.6204	-0.2286	0.042*
H24B	0.3488	0.5616	-0.0945	0.042*
H24C	0.4046	0.5830	-0.1944	0.042*

Atomic displacement parameters (\AA^2)

	U^{11}	U^{22}	U^{33}	U^{12}	U^{13}	U^{23}
O1	0.0265 (8)	0.0202 (8)	0.0271 (8)	-0.0029 (6)	-0.0030 (7)	0.0048 (6)
O2	0.0308 (8)	0.0244 (8)	0.0291 (8)	0.0063 (7)	-0.0075 (7)	0.0009 (7)
O3	0.0216 (7)	0.0233 (8)	0.0235 (8)	0.0036 (6)	-0.0069 (6)	-0.0009 (6)
N1	0.0198 (8)	0.0244 (9)	0.0225 (9)	0.0007 (7)	0.0031 (7)	0.0006 (8)
C1	0.0218 (10)	0.0195 (10)	0.0202 (10)	0.0052 (8)	0.0057 (8)	0.0059 (8)
C2	0.0173 (9)	0.0200 (10)	0.0187 (10)	0.0001 (8)	0.0031 (8)	0.0001 (8)
C3	0.0162 (9)	0.0229 (10)	0.0143 (9)	0.0016 (8)	-0.0008 (8)	0.0021 (8)
C4	0.0165 (9)	0.0206 (10)	0.0178 (10)	0.0008 (8)	0.0023 (8)	0.0009 (8)
C5	0.0173 (9)	0.0229 (11)	0.0191 (10)	-0.0008 (8)	0.0011 (8)	-0.0032 (8)
C6	0.0154 (9)	0.0273 (11)	0.0173 (10)	0.0025 (8)	0.0004 (8)	0.0000 (9)
C7	0.0294 (12)	0.0291 (12)	0.0286 (12)	-0.0085 (9)	0.0037 (10)	0.0017 (10)
C8	0.0252 (10)	0.0193 (10)	0.0204 (10)	-0.0026 (8)	0.0000 (8)	0.0017 (8)
C9	0.0393 (13)	0.0220 (11)	0.0273 (12)	-0.0012 (10)	0.0047 (10)	-0.0015 (9)
C10	0.0351 (12)	0.0253 (12)	0.0275 (12)	-0.0087 (10)	-0.0080 (10)	0.0024 (10)
C11	0.0201 (10)	0.0191 (10)	0.0192 (10)	0.0004 (8)	0.0002 (8)	-0.0015 (8)
C12	0.0262 (11)	0.0220 (11)	0.0280 (12)	-0.0025 (9)	0.0031 (9)	0.0009 (9)
C13	0.0309 (11)	0.0232 (11)	0.0232 (11)	0.0051 (9)	0.0023 (9)	-0.0025 (9)
C14	0.0269 (11)	0.0202 (11)	0.0254 (11)	0.0022 (9)	-0.0024 (9)	0.0009 (9)
C15	0.0212 (10)	0.0212 (11)	0.0235 (11)	-0.0011 (8)	0.0039 (8)	-0.0020 (9)
C16	0.0170 (9)	0.0283 (12)	0.0184 (10)	0.0001 (8)	0.0014 (8)	0.0009 (9)
C17	0.0216 (10)	0.0247 (11)	0.0238 (11)	-0.0014 (9)	0.0013 (9)	-0.0008 (9)
C18	0.0226 (10)	0.0227 (11)	0.0227 (11)	0.0021 (8)	0.0012 (9)	0.0015 (9)
C19	0.0187 (10)	0.0239 (11)	0.0174 (10)	0.0028 (8)	0.0004 (8)	0.0021 (8)
C20	0.0205 (10)	0.0228 (11)	0.0204 (10)	-0.0005 (8)	0.0016 (8)	0.0003 (9)
C21	0.0216 (10)	0.0241 (11)	0.0238 (11)	0.0060 (9)	0.0004 (8)	0.0039 (9)
C22	0.0194 (10)	0.0232 (11)	0.0198 (10)	0.0001 (8)	0.0012 (8)	0.0020 (9)
C23	0.0240 (11)	0.0295 (12)	0.0231 (11)	0.0018 (9)	-0.0082 (9)	0.0016 (9)
C24	0.0261 (11)	0.0306 (12)	0.0271 (12)	-0.0001 (10)	0.0012 (9)	-0.0006 (10)

Geometric parameters (\AA , $^\circ$)

O1—C1	1.355 (2)	C11—C14	1.531 (3)
O1—H1	0.866 (10)	C11—C13	1.537 (3)
O2—C22	1.207 (2)	C11—C12	1.537 (3)
O3—C22	1.346 (2)	C12—H12A	0.9800

O3—C23	1.450 (2)	C12—H12B	0.9800
N1—C15	1.273 (3)	C12—H12C	0.9800
N1—C16	1.428 (3)	C13—H13A	0.9800
C1—C6	1.406 (3)	C13—H13B	0.9800
C1—C2	1.410 (3)	C13—H13C	0.9800
C2—C3	1.397 (3)	C14—H14A	0.9800
C2—C8	1.535 (3)	C14—H14B	0.9800
C3—C4	1.402 (3)	C14—H14C	0.9800
C3—H3	0.9500	C15—H15	0.9500
C4—C5	1.389 (3)	C16—C17	1.381 (3)
C4—C11	1.529 (3)	C16—C21	1.399 (3)
C5—C6	1.393 (3)	C17—C18	1.379 (3)
C5—H5	0.9500	C17—H17	0.9500
C6—C15	1.465 (3)	C18—C19	1.395 (3)
C7—C8	1.540 (3)	C18—H18	0.9500
C7—H7A	0.9800	C19—C20	1.389 (3)
C7—H7B	0.9800	C19—C22	1.488 (3)
C7—H7C	0.9800	C20—C21	1.387 (3)
C8—C10	1.534 (3)	C20—H20	0.9500
C8—C9	1.539 (3)	C21—H21	0.9500
C9—H9A	0.9800	C23—C24	1.494 (3)
C9—H9B	0.9800	C23—H23A	0.9900
C9—H9C	0.9800	C23—H23B	0.9900
C10—H10A	0.9800	C24—H24A	0.9800
C10—H10B	0.9800	C24—H24B	0.9800
C10—H10C	0.9800	C24—H24C	0.9800
C1—O1—H1	106 (2)	H12A—C12—H12B	109.5
C22—O3—C23	114.99 (16)	C11—C12—H12C	109.5
C15—N1—C16	118.83 (18)	H12A—C12—H12C	109.5
O1—C1—C6	119.20 (18)	H12B—C12—H12C	109.5
O1—C1—C2	120.37 (18)	C11—C13—H13A	109.5
C6—C1—C2	120.43 (18)	C11—C13—H13B	109.5
C3—C2—C1	116.73 (18)	H13A—C13—H13B	109.5
C3—C2—C8	121.24 (17)	C11—C13—H13C	109.5
C1—C2—C8	121.96 (18)	H13A—C13—H13C	109.5
C2—C3—C4	124.26 (18)	H13B—C13—H13C	109.5
C2—C3—H3	117.9	C11—C14—H14A	109.5
C4—C3—H3	117.9	C11—C14—H14B	109.5
C5—C4—C3	116.83 (18)	H14A—C14—H14B	109.5
C5—C4—C11	120.03 (17)	C11—C14—H14C	109.5
C3—C4—C11	123.12 (17)	H14A—C14—H14C	109.5
C4—C5—C6	121.49 (19)	H14B—C14—H14C	109.5
C4—C5—H5	119.3	N1—C15—C6	122.09 (19)
C6—C5—H5	119.3	N1—C15—H15	119.0
C5—C6—C1	120.03 (18)	C6—C15—H15	119.0
C5—C6—C15	117.32 (18)	C17—C16—C21	119.51 (19)
C1—C6—C15	122.63 (19)	C17—C16—N1	117.79 (19)
C8—C7—H7A	109.5	C21—C16—N1	122.64 (19)
C8—C7—H7B	109.5	C18—C17—C16	120.3 (2)

supplementary materials

H7A—C7—H7B	109.5	C18—C17—H17	119.8
C8—C7—H7C	109.5	C16—C17—H17	119.8
H7A—C7—H7C	109.5	C17—C18—C19	120.5 (2)
H7B—C7—H7C	109.5	C17—C18—H18	119.7
C10—C8—C2	112.00 (17)	C19—C18—H18	119.7
C10—C8—C7	107.87 (18)	C20—C19—C18	119.40 (18)
C2—C8—C7	108.92 (17)	C20—C19—C22	122.69 (19)
C10—C8—C9	106.89 (17)	C18—C19—C22	117.87 (18)
C2—C8—C9	111.03 (17)	C21—C20—C19	120.0 (2)
C7—C8—C9	110.07 (18)	C21—C20—H20	120.0
C8—C9—H9A	109.5	C19—C20—H20	120.0
C8—C9—H9B	109.5	C20—C21—C16	120.18 (19)
H9A—C9—H9B	109.5	C20—C21—H21	119.9
C8—C9—H9C	109.5	C16—C21—H21	119.9
H9A—C9—H9C	109.5	O2—C22—O3	123.23 (18)
H9B—C9—H9C	109.5	O2—C22—C19	124.17 (19)
C8—C10—H10A	109.5	O3—C22—C19	112.57 (17)
C8—C10—H10B	109.5	O3—C23—C24	107.99 (17)
H10A—C10—H10B	109.5	O3—C23—H23A	110.1
C8—C10—H10C	109.5	C24—C23—H23A	110.1
H10A—C10—H10C	109.5	O3—C23—H23B	110.1
H10B—C10—H10C	109.5	C24—C23—H23B	110.1
C4—C11—C14	112.55 (16)	H23A—C23—H23B	108.4
C4—C11—C13	108.92 (16)	C23—C24—H24A	109.5
C14—C11—C13	108.30 (16)	C23—C24—H24B	109.5
C4—C11—C12	109.85 (16)	H24A—C24—H24B	109.5
C14—C11—C12	107.53 (17)	C23—C24—H24C	109.5
C13—C11—C12	109.64 (17)	H24A—C24—H24C	109.5
C11—C12—H12A	109.5	H24B—C24—H24C	109.5
C11—C12—H12B	109.5		
O1—C1—C2—C3	-175.11 (17)	C3—C4—C11—C13	-122.2 (2)
C6—C1—C2—C3	4.6 (3)	C5—C4—C11—C12	-64.1 (2)
O1—C1—C2—C8	1.8 (3)	C3—C4—C11—C12	117.6 (2)
C6—C1—C2—C8	-178.52 (18)	C16—N1—C15—C6	178.21 (17)
C1—C2—C3—C4	-1.5 (3)	C5—C6—C15—N1	-174.29 (19)
C8—C2—C3—C4	-178.36 (18)	C1—C6—C15—N1	4.0 (3)
C2—C3—C4—C5	-2.7 (3)	C15—N1—C16—C17	151.3 (2)
C2—C3—C4—C11	175.58 (19)	C15—N1—C16—C21	-31.4 (3)
C3—C4—C5—C6	3.9 (3)	C21—C16—C17—C18	3.2 (3)
C11—C4—C5—C6	-174.44 (18)	N1—C16—C17—C18	-179.46 (18)
C4—C5—C6—C1	-0.9 (3)	C16—C17—C18—C19	-0.4 (3)
C4—C5—C6—C15	177.43 (18)	C17—C18—C19—C20	-2.2 (3)
O1—C1—C6—C5	176.18 (18)	C17—C18—C19—C22	175.61 (19)
C2—C1—C6—C5	-3.5 (3)	C18—C19—C20—C21	2.0 (3)
O1—C1—C6—C15	-2.1 (3)	C22—C19—C20—C21	-175.68 (18)
C2—C1—C6—C15	178.18 (18)	C19—C20—C21—C16	0.8 (3)
C3—C2—C8—C10	-9.4 (3)	C17—C16—C21—C20	-3.4 (3)
C1—C2—C8—C10	173.86 (19)	N1—C16—C21—C20	179.43 (18)
C3—C2—C8—C7	109.8 (2)	C23—O3—C22—O2	0.8 (3)

C1—C2—C8—C7	-66.9 (2)	C23—O3—C22—C19	178.95 (17)
C3—C2—C8—C9	-128.8 (2)	C20—C19—C22—O2	177.1 (2)
C1—C2—C8—C9	54.5 (2)	C18—C19—C22—O2	-0.6 (3)
C5—C4—C11—C14	176.14 (18)	C20—C19—C22—O3	-1.0 (3)
C3—C4—C11—C14	-2.1 (3)	C18—C19—C22—O3	-178.74 (18)
C5—C4—C11—C13	56.0 (2)	C22—O3—C23—C24	-173.52 (17)

Hydrogen-bond geometry (Å, °)

<i>D</i> —H... <i>A</i>	<i>D</i> —H	H... <i>A</i>	<i>D</i> ... <i>A</i>	<i>D</i> —H... <i>A</i>
O1—H1...N1	0.87 (1)	1.80 (2)	2.609 (2)	154 (3)

Fig. 1




Fig. 2

