

6-Isopropyl-5-methoxy-3-phenyl-3*H*-1,2,3-triazolo[4,5-*d*]pyrimidin-7(6*H*)-one

Xiao-Hua Zeng,^a Hong-Mei Wang,^a Shou-Heng Deng^{b*}
and Li-Li Chen^a

^aInstitute of Medicinal Chemistry, Hubei Medical University, Shiyan 442000, People's Republic of China, and ^bCenter of Oncology, People's Hospital affiliated with Hubei Medical University, Shi Yan 442000, People's Republic of China
Correspondence e-mail: zkenn710@yahoo.com.cn


Received 9 October 2010; accepted 16 October 2010

Key indicators: single-crystal X-ray study; $T = 298\text{ K}$; mean $\sigma(\text{C}-\text{C}) = 0.005\text{ \AA}$; R factor = 0.065; wR factor = 0.185; data-to-parameter ratio = 11.3.

In the title compound, $\text{C}_{14}\text{H}_{15}\text{N}_5\text{O}_2$, the whole molecule apart from the terminal C atoms of the isopropyl group is located on a crystallographic mirror plane. An intramolecular C—H···N hydrogen-bonding interaction may stabilize the molecular conformation. The crystal packing features weak slipped π – π interactions between the pyrimidine and the phenyl rings of symmetry-related molecules [centroid–centroid distance = 3.746 (1) \AA , slippage of 1.574 \AA].

Related literature

For the biological activity of 8-azaguanine derivatives, see: Roblin *et al.* (1945); Ding *et al.* (2004); Mitchell *et al.* (1950); Levine *et al.* (1963); Montgomery *et al.* (1962); Yamamoto *et al.* (1967); Bariana (1971); Holland *et al.* (1975). For related structures, see: Chen & Shi (2006); Ferguson *et al.* (1998); Li *et al.* (2004); Maldonado *et al.* (2006); Wang *et al.* (2006); Xiao & Shi (2007); Zeng *et al.* (2006, 2009); Zhao, Hu *et al.* (2005); Zhao, Wang & Ding (2005); Zhao, Xie *et al.* (2005).



Experimental

Crystal data

$\text{C}_{14}\text{H}_{15}\text{N}_5\text{O}_2$
 $M_r = 285.31$
Orthorhombic, $Pnma$

$a = 14.921 (2)\text{ \AA}$
 $b = 6.7989 (11)\text{ \AA}$
 $c = 13.839 (2)\text{ \AA}$

$V = 1404.0 (4)\text{ \AA}^3$
 $Z = 4$
Mo $K\alpha$ radiation

$\mu = 0.10\text{ mm}^{-1}$
 $T = 298\text{ K}$
 $0.16 \times 0.12 \times 0.10\text{ mm}$

Data collection

Bruker SMART CCD area-detector diffractometer
Absorption correction: multi-scan (*SADABS*; Sheldrick, 2008)
 $T_{\min} = 0.985$, $T_{\max} = 0.991$

7422 measured reflections
1418 independent reflections
1045 reflections with $I > 2\sigma(I)$
 $R_{\text{int}} = 0.039$

Refinement

$R[F^2 > 2\sigma(F^2)] = 0.065$
 $wR(F^2) = 0.185$
 $S = 1.07$
1418 reflections

125 parameters
H-atom parameters constrained
 $\Delta\rho_{\max} = 0.23\text{ e \AA}^{-3}$
 $\Delta\rho_{\min} = -0.29\text{ e \AA}^{-3}$

Table 1
Hydrogen-bond geometry (\AA , $^\circ$).

$D-\text{H} \cdots A$	$D-\text{H}$	$\text{H} \cdots A$	$D \cdots A$	$D-\text{H} \cdots A$
C2—H2···N4	0.93	2.37	3.021 (4)	127

Data collection: *SMART* (Bruker, 2001); cell refinement: *SAINT* (Bruker, 2001); data reduction: *SAINT*; program(s) used to solve structure: *SHELXS97* (Sheldrick, 2008); program(s) used to refine structure: *SHELXL97* (Sheldrick, 2008); molecular graphics: *ORTEPIII* (Burnett & Johnson, 1996), *ORTEP-3 for Windows* (Farrugia, 1999) and *PLATON* (Spek, 2009); software used to prepare material for publication: *SHELXTL* (Sheldrick, 2008).

We gratefully acknowledge financial support of this work by the National Basic Research Program of China (2003CB114400), the National Natural Science Foundation of China (20372023, 20102001), the Educational Commission of Hubei Province (grant Nos. B200624004, B20092412), the Shiyan Municipal Science and Technology Bureau (grant No. 20061835) and Hubei Medical University (grant Nos. 2007QDJ15, 2007ZQB19, 2007ZQB20).

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: DN2610).

References

- Bariana, D. S. (1971). *J. Med. Chem.* **14**, 535–543.
- Bruker (2001). *SMART* and *SAINT*. Bruker AXS Inc., Madison, Wisconsin, USA.
- Burnett, M. N. & Johnson, C. K. (1996). *ORTEPIII*. Report ORNL-6895. Oak Ridge National Laboratory, Tennessee, USA.
- Chen, X.-B. & Shi, D.-Q. (2006). *Acta Cryst. E* **62**, o4780–o4782.
- Ding, M. W., Xu, S. Z. & Zhao, J. F. (2004). *J. Org. Chem.* **69**, 8366–8371.
- Farrugia, L. J. (1999). *J. Appl. Cryst.* **32**, 837–838.
- Ferguson, G., Low, J. N., Nogueras, M., Cobo, J., Lopez, M. D., Quijano, M. L. & Sanchez, A. (1998). *Acta Cryst. C* **54**, IUC9800031.
- Holland, A., Jackson, D., Chaplen, P., Lunt, E., Marshall, S., Pain, C. L. & Wooldridge, K. R. H. (1975). *Eur. J. Med. Chem.* **10**, 447–449.
- Levine, R. J., Hall, T. C. & Harris, C. A. (1963). *Cancer (N. Y.)*, **16**, 269–272.
- Li, M., Wen, L. R., Fu, W. J., Hu, F. Z. & Yang, H. Z. (2004). *Chin. J. Struct. Chem.* **23**, 11–14.
- Maldonado, C. R., Quirós, M. & Salas, J. M. (2006). *Acta Cryst. C* **62**, o489–o491.
- Mitchell, J. H., Skipper, H. E. & Bennett, L. L. (1950). *Cancer Res.* **10**, 647–649.
- Montgomery, J. A., Schabel, F. M. & Skipper, H. E. (1962). *Cancer Res.* **22**, 504–509.

organic compounds

- Roblin, R. O., Lampen, J. O., English, J. P., Cole, Q. P. & Vaughan, J. R. (1945). *J. Am. Chem. Soc.* **67**, 290–294.
- Sheldrick, G. M. (2008). *Acta Cryst. A* **64**, 112–122.
- Spek, A. L. (2009). *Acta Cryst. D* **65**, 148–155.
- Wang, H.-M., Zeng, X.-H., Hu, Z.-Q., Li, G.-H. & Tian, J.-H. (2006). *Acta Cryst. E* **62**, o5038–o5040.
- Xiao, L.-X. & Shi, D.-Q. (2007). *Acta Cryst. E* **63**, o2843.
- Yamamoto, I., Inoki, R., Tamari, Y. & Iwatsubo, K. (1967). *Jpn J. Pharmacol.* **17**, 140–142.
- Zeng, X.-H., Liu, X.-L., Deng, S.-H., Chen, P. & Wang, H.-M. (2009). *Acta Cryst. E* **65**, o2583–o2584.
- Zeng, X.-H., Wang, H.-M., Ding, M.-W. & He, H.-W. (2006). *Acta Cryst. E* **62**, o1888–o1890.
- Zhao, J.-F., Hu, Y.-G., Ding, M.-W. & He, H.-W. (2005). *Acta Cryst. E* **61**, o2791–o2792.
- Zhao, J. F., Wang, C. G. & Ding, M. W. (2005). *Chin. J. Struct. Chem.* **24**, 439–444.
- Zhao, J. F., Xie, C., Ding, M. W. & He, H. W. (2005). *Chem. Lett.* **34**, 1020–1022.

supplementary materials

Acta Cryst. (2010). E66, o2947-o2948 [doi:10.1107/S1600536810041978]

6-Isopropyl-5-methoxy-3-phenyl-3*H*-1,2,3-triazolo[4,5-*d*]pyrimidin-7(6*H*)-one

X.-H. Zeng, H.-M. Wang, S.-H. Deng and L.-L. Chen

Comment

The derivatives of heterocycles containing 8-azaguanine system, which are well known bioisosteres of guanine, are of great importance because of their remarkable biological properties, such as antimicrobial or antifungal activities (Roblin *et al.*, 1945; Ding *et al.*, 2004), encephaloma cell inhibitor (Mitchell *et al.*, 1950; Levine *et al.*, 1963), antileukemie (Montgomery *et al.*, 1962), hypersusceptibility inhibitor and acesodyne activities (Yamamoto *et al.*, 1967; Bariana, 1971; Holland *et al.*, 1975).

In recent years, Zhao's group succeeded in synthesizing the derivatives of 8-azaguanine *via* aza-Wittig reaction of beta-ethoxycarbonyl iminophosphorane with aromatic isocyanates (Zhao, Xie *et al.*, 2005). As a continuation of the quest for new biologically active derivatives of 8-azaguanine, the title compound, (I), was obtained from beta-ethoxycarbonyl iminophosphorane with aliphatic isocyanate, and structurally characterized.

In the title compound, C₁₄H₁₅N₅O₂, the whole molecule but the terminal C atoms of the isopropyl group is located in a mirror plane and is then perfectly planar (Fig. 1). The bond lengths and angles in the triazolopyrimidinone moiety are in good agreement with those observed for closely related structures (Zhao, Hu *et al.*, 2005; Zhao, Wang & Ding, 2005). the triazolopyrimidine ring system is perfectly coplanar (Chen & Shi, 2006; Ferguson *et al.*, 1998; Li *et al.*, 2004; Maldonado *et al.*, 2006; Wang *et al.*, 2006; Xiao & Shi, 2007; Zeng *et al.*, 2009).

The molecules are packed along the b axis with weak slippage π – π interaction between the pyrimidin and the phenyl rings of symmetry related molecules (Centroid to centroid distance = 3.746 (1) Å, interplanar distance = 3.399 Å with a slippage of 1.574 Å).

Experimental

To the solution of carbodiimide prepared according to Zeng *et al.* (2006) in a mixed solvent (CH₂Cl₂/MeOH, 1:4 v/v, 15 ml) was added a fresh prepared solution of Na/MeOH (0.1 g/2 ml). After stirring the reaction mixture for 6 h, the solvent was removed under reduced pressure and the residue was recrystallized from EtOH to give the title compound (I) in 89% yield (m.p. 471 K). Elemental analysis: calculated for C₁₄H₁₅N₅O₂: C, 58.94; H, 5.30; N, 24.55%. Found: C, 57.62; H, 5.72; N, 24.01%. Crystals suitable for X-ray diffraction study were obtained by recrystallization from hexane and dichloromethane (1:3 v/v) at room temperature.

Refinement

All H atoms attached to C atoms and N atom were fixed geometrically and treated as riding with C—H = 0.98 Å (methine), 0.96 Å (methyl) or 0.93 Å (aromatic) with U_{iso}(H) = 1.2U_{eq}(C) or U_{iso}(H) = 1.5U_{eq}(C_{methyl}).

supplementary materials

Figures




Fig. 1. View of the molecule of showing the atom-labeling scheme. Displacement ellipsoids are drawn at the 30% probability level. H-atoms are represented as small spheres of arbitrary radii. [Symmetry code: (i) $-x-1/2, y+1/2, z-1/2$].




Fig. 2. Packing view showing the stacking of the molecules along the b axis. H atoms have been omitted for clarity.

6-Isopropyl-5-methoxy-3-phenyl-3*H*-1,2,3-triazolo[4,5-*d*]pyrimidin-7(6*H*)-one

Crystal data

C ₁₄ H ₁₅ N ₅ O ₂	$F(000) = 600$
$M_r = 285.31$	$D_x = 1.350 \text{ Mg m}^{-3}$
Orthorhombic, <i>Pnma</i>	Mo $K\alpha$ radiation, $\lambda = 0.71073 \text{ \AA}$
Hall symbol: -P 2ac 2n	Cell parameters from 982 reflections
$a = 14.921 (2) \text{ \AA}$	$\theta = 2.9\text{--}20.5^\circ$
$b = 6.7989 (11) \text{ \AA}$	$\mu = 0.10 \text{ mm}^{-1}$
$c = 13.839 (2) \text{ \AA}$	$T = 298 \text{ K}$
$V = 1404.0 (4) \text{ \AA}^3$	Block, colourless
$Z = 4$	$0.16 \times 0.12 \times 0.10 \text{ mm}$

Data collection

Bruker SMART CCD area-detector diffractometer	1418 independent reflections
Radiation source: fine-focus sealed tube graphite	1045 reflections with $I > 2\sigma(I)$
φ and ω scans	$R_{\text{int}} = 0.039$
Absorption correction: multi-scan (<i>SADABS</i> ; Sheldrick, 2008)	$\theta_{\max} = 25.5^\circ, \theta_{\min} = 2.0^\circ$
$T_{\min} = 0.985, T_{\max} = 0.991$	$h = -17 \rightarrow 18$
7422 measured reflections	$k = -8 \rightarrow 8$
	$l = -13 \rightarrow 16$

Refinement

Refinement on F^2	Primary atom site location: structure-invariant direct methods
Least-squares matrix: full	Secondary atom site location: difference Fourier map
$R[F^2 > 2\sigma(F^2)] = 0.065$	Hydrogen site location: inferred from neighbouring sites
$wR(F^2) = 0.185$	H-atom parameters constrained

$S = 1.07$	$w = 1/[\sigma^2(F_o^2) + (0.0883P)^2 + 0.4734P]$
	where $P = (F_o^2 + 2F_c^2)/3$
1418 reflections	$(\Delta/\sigma)_{\max} < 0.001$
125 parameters	$\Delta\rho_{\max} = 0.23 \text{ e \AA}^{-3}$
0 restraints	$\Delta\rho_{\min} = -0.29 \text{ e \AA}^{-3}$

Special details

Geometry. All esds (except the esd in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell esds are taken into account individually in the estimation of esds in distances, angles and torsion angles; correlations between esds in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell esds is used for estimating esds involving l.s. planes.

Refinement. Refinement of F^2 against ALL reflections. The weighted R -factor wR and goodness of fit S are based on F^2 , conventional R -factors R are based on F , with F set to zero for negative F^2 . The threshold expression of $F^2 > \sigma(F^2)$ is used only for calculating R -factors(gt) etc. and is not relevant to the choice of reflections for refinement. R -factors based on F^2 are statistically about twice as large as those based on F , and R -factors based on ALL data will be even larger.

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (\AA^2)

	x	y	z	$U_{\text{iso}}^*/U_{\text{eq}}$	Occ. (<1)
O1	0.36483 (19)	0.2500	0.3850 (3)	0.1092 (13)	
O2	0.24610 (18)	0.2500	0.6874 (2)	0.0829 (9)	
N1	0.06296 (18)	0.2500	0.41475 (18)	0.0507 (7)	
N2	0.0868 (3)	0.2500	0.3193 (2)	0.0843 (12)	
N3	0.1730 (3)	0.2500	0.3126 (2)	0.0908 (12)	
N4	0.14519 (17)	0.2500	0.5659 (2)	0.0495 (7)	
N5	0.30395 (19)	0.2500	0.5368 (3)	0.0629 (9)	
C1	-0.0298 (2)	0.2500	0.4402 (2)	0.0462 (8)	
C2	-0.0554 (2)	0.2500	0.5362 (2)	0.0546 (9)	
H2	-0.0124	0.2500	0.5848	0.065*	
C3	-0.1452 (2)	0.2500	0.5591 (3)	0.0599 (10)	
H3	-0.1626	0.2500	0.6236	0.072*	
C4	-0.2092 (3)	0.2500	0.4886 (3)	0.0630 (10)	
H4	-0.2697	0.2500	0.5049	0.076*	
C5	-0.1835 (3)	0.2500	0.3938 (3)	0.0672 (11)	
H5	-0.2270	0.2500	0.3457	0.081*	
C6	-0.0949 (3)	0.2500	0.3683 (3)	0.0580 (10)	
H6	-0.0783	0.2500	0.3035	0.070*	
C7	0.1382 (2)	0.2500	0.4688 (2)	0.0464 (8)	
C8	0.2070 (2)	0.2500	0.4039 (3)	0.0608 (10)	
C9	0.2979 (3)	0.2500	0.4346 (3)	0.0721 (11)	
C10	0.2277 (2)	0.2500	0.5942 (3)	0.0589 (9)	
C11	0.3973 (3)	0.2500	0.5789 (4)	0.0879 (14)	
H11	0.4338	0.2500	0.5200	0.105*	
C12	0.4225 (2)	0.0624 (6)	0.6212 (3)	0.1210 (15)	
H12A	0.4861	0.0599	0.6319	0.181*	
H12B	0.4063	-0.0421	0.5780	0.181*	

supplementary materials

H12C	0.3919	0.0454	0.6816	0.181*	
C13	0.1714 (3)	0.2500	0.7534 (3)	0.1016 (17)	
H13A	0.1933	0.2500	0.8186	0.152*	
H13B	0.1356	0.1347	0.7427	0.152*	0.50
H13C	0.1356	0.3653	0.7427	0.152*	0.50

Atomic displacement parameters (\AA^2)

	U^{11}	U^{22}	U^{33}	U^{12}	U^{13}	U^{23}
O1	0.0561 (19)	0.156 (3)	0.115 (3)	0.000	0.0314 (18)	0.000
O2	0.0545 (17)	0.125 (3)	0.0687 (18)	0.000	-0.0150 (14)	0.000
N1	0.0539 (18)	0.0584 (17)	0.0399 (14)	0.000	0.0009 (13)	0.000
N2	0.075 (2)	0.131 (3)	0.0470 (19)	0.000	0.0067 (17)	0.000
N3	0.074 (3)	0.142 (4)	0.056 (2)	0.000	0.0142 (18)	0.000
N4	0.0419 (16)	0.0507 (16)	0.0559 (18)	0.000	-0.0071 (12)	0.000
N5	0.0365 (16)	0.0593 (19)	0.093 (2)	0.000	-0.0019 (15)	0.000
C1	0.050 (2)	0.0395 (17)	0.0485 (19)	0.000	-0.0052 (15)	0.000
C2	0.043 (2)	0.066 (2)	0.055 (2)	0.000	-0.0077 (16)	0.000
C3	0.052 (2)	0.065 (2)	0.063 (2)	0.000	0.0000 (17)	0.000
C4	0.047 (2)	0.060 (2)	0.082 (3)	0.000	-0.0074 (19)	0.000
C5	0.054 (2)	0.064 (2)	0.083 (3)	0.000	-0.030 (2)	0.000
C6	0.067 (3)	0.056 (2)	0.051 (2)	0.000	-0.0148 (18)	0.000
C7	0.048 (2)	0.0419 (18)	0.0497 (19)	0.000	-0.0003 (15)	0.000
C8	0.050 (2)	0.072 (2)	0.060 (2)	0.000	0.0094 (17)	0.000
C9	0.060 (3)	0.078 (3)	0.078 (3)	0.000	0.016 (2)	0.000
C10	0.055 (2)	0.055 (2)	0.067 (2)	0.000	-0.0120 (19)	0.000
C11	0.045 (2)	0.086 (3)	0.133 (4)	0.000	-0.013 (2)	0.000
C12	0.082 (2)	0.118 (3)	0.163 (4)	0.011 (2)	-0.034 (2)	0.043 (3)
C13	0.079 (3)	0.172 (5)	0.054 (2)	0.000	-0.016 (2)	0.000

Geometric parameters (\AA , $^\circ$)

O1—C9	1.212 (5)	C3—H3	0.9300
O2—C10	1.319 (4)	C4—C5	1.367 (6)
O2—C13	1.441 (5)	C4—H4	0.9300
N1—C7	1.350 (4)	C5—C6	1.369 (5)
N1—N2	1.368 (4)	C5—H5	0.9300
N1—C1	1.428 (4)	C6—H6	0.9300
N2—N3	1.290 (5)	C7—C8	1.364 (5)
N3—C8	1.360 (5)	C8—C9	1.421 (5)
N4—C10	1.293 (4)	C11—C12 ⁱ	1.453 (4)
N4—C7	1.348 (4)	C11—C12	1.453 (4)
N5—C10	1.387 (5)	C11—H11	0.9800
N5—C9	1.418 (5)	C12—H12A	0.9600
N5—C11	1.510 (5)	C12—H12B	0.9600
C1—C2	1.382 (5)	C12—H12C	0.9600
C1—C6	1.391 (4)	C13—H13A	0.9600
C2—C3	1.377 (5)	C13—H13B	0.9600

C2—H2	0.9300	C13—H13C	0.9600
C3—C4	1.365 (5)		
C10—O2—C13	117.3 (3)	N4—C7—C8	126.8 (3)
C7—N1—N2	108.6 (3)	N1—C7—C8	105.1 (3)
C7—N1—C1	132.0 (3)	N3—C8—C7	109.4 (3)
N2—N1—C1	119.4 (3)	N3—C8—C9	129.3 (4)
N3—N2—N1	109.2 (3)	C7—C8—C9	121.4 (4)
N2—N3—C8	107.8 (3)	O1—C9—N5	120.8 (4)
C10—N4—C7	112.0 (3)	O1—C9—C8	128.1 (4)
C10—N5—C9	121.2 (3)	N5—C9—C8	111.1 (3)
C10—N5—C11	122.4 (4)	N4—C10—O2	119.6 (3)
C9—N5—C11	116.4 (3)	N4—C10—N5	127.5 (4)
C2—C1—C6	119.6 (3)	O2—C10—N5	112.9 (3)
C2—C1—N1	120.4 (3)	C12 ⁱ —C11—C12	122.8 (5)
C6—C1—N1	120.0 (3)	C12 ⁱ —C11—N5	113.2 (2)
C3—C2—C1	119.4 (3)	C12—C11—N5	113.2 (2)
C3—C2—H2	120.3	C12 ⁱ —C11—H11	101.0
C1—C2—H2	120.3	C12—C11—H11	101.0
C4—C3—C2	121.1 (4)	N5—C11—H11	101.0
C4—C3—H3	119.5	C11—C12—H12A	109.5
C2—C3—H3	119.5	C11—C12—H12B	109.5
C3—C4—C5	119.3 (4)	H12A—C12—H12B	109.5
C3—C4—H4	120.3	C11—C12—H12C	109.5
C5—C4—H4	120.3	H12A—C12—H12C	109.5
C4—C5—C6	121.2 (3)	H12B—C12—H12C	109.5
C4—C5—H5	119.4	O2—C13—H13A	109.5
C6—C5—H5	119.4	O2—C13—H13B	109.5
C5—C6—C1	119.4 (3)	H13A—C13—H13B	109.5
C5—C6—H6	120.3	O2—C13—H13C	109.5
C1—C6—H6	120.3	H13A—C13—H13C	109.5
N4—C7—N1	128.1 (3)	H13B—C13—H13C	109.5

Symmetry codes: (i) $x, -y+1/2, z$.

Hydrogen-bond geometry (\AA , $^\circ$)

$D—H\cdots A$	$D—H$	$H\cdots A$	$D\cdots A$	$D—H\cdots A$
C2—H2 \cdots N4	0.93	2.37	3.021 (4)	127
C6—H6 \cdots N2	0.93	2.47	2.794 (5)	100
C11—H11 \cdots O1	0.98	2.13	2.727 (7)	117
C12—H12C \cdots O2	0.96	2.58	3.065 (5)	111

supplementary materials

Fig. 1




Fig. 2

