

***rac*-2-Hydroxy-2-(2-oxocyclopentyl)-1*H*-indene-1,3(2*H*)-dione**

J. Kalyana Sundar,^a S. Maharani,^b R. Ranjith Kumar,^b
S. Natarajan,^a J. Suresh^c and P. L. Nilantha Lakshman^{d*}

^aDepartment of Physics, Madurai Kamaraj University, Madurai 625 021, India,
^bDepartment of Organic Chemistry, School of Chemistry, Madurai Kamaraj University, Madurai 625 021, India, ^cDepartment of Physics, The Madura College, Madurai 625 011, India, and ^dDepartment of Food Science and Technology, University of Ruhuna, Mapalana, Kamburupitiya 81100, Sri Lanka
Correspondence e-mail: plakshmannilantha@ymail.com


Received 16 September 2010; accepted 21 October 2010

Key indicators: single-crystal X-ray study; $T = 293\text{ K}$; mean $\sigma(\text{C}-\text{C}) = 0.002\text{ \AA}$; R factor = 0.038; wR factor = 0.104; data-to-parameter ratio = 12.5.

In the title compound, $\text{C}_{14}\text{H}_{12}\text{O}_4$, the indene unit is essentially planar [r.m.s. deviation = 0.0309 (1) \AA] and the cyclopentanone ring adopts a twist form. In the crystal, molecules are joined via pairs of $\text{O}-\text{H} \cdots \text{O}$ hydrogen bonds into centrosymmetric dimers.

Related literature

For a similar structure, see: Pentala *et al.* (2009).



Experimental

Crystal data

$\text{C}_{14}\text{H}_{12}\text{O}_4$
 $M_r = 244.24$

Triclinic, $P\bar{1}$
 $a = 8.044 (3)\text{ \AA}$

$b = 8.404 (4)\text{ \AA}$	$Z = 2$
$c = 10.239 (3)\text{ \AA}$	$\text{Mo } K\alpha$ radiation
$\alpha = 66.95 (3)^\circ$	$\mu = 0.10\text{ mm}^{-1}$
$\beta = 74.36 (2)^\circ$	$T = 293\text{ K}$
$\gamma = 68.50 (3)^\circ$	$0.26 \times 0.22 \times 0.19\text{ mm}$
$V = 586.1 (4)\text{ \AA}^3$	

Data collection

Nonius MACH3 diffractometer
Absorption correction: ψ scan
(North *et al.*, 1968)
 $T_{\min} = 0.974$, $T_{\max} = 0.981$
2534 measured reflections

2054 independent reflections
1886 reflections with $I > 2\sigma(I)$
 $R_{\text{int}} = 0.016$
3 standard reflections every 60 min
intensity decay: none

Refinement

$R[F^2 > 2\sigma(F^2)] = 0.038$
 $wR(F^2) = 0.104$
 $S = 1.07$
2054 reflections

164 parameters
H-atom parameters constrained
 $\Delta\rho_{\max} = 0.25\text{ e \AA}^{-3}$
 $\Delta\rho_{\min} = -0.23\text{ e \AA}^{-3}$

Table 1
Hydrogen-bond geometry (\AA , $^\circ$).

$D-\text{H} \cdots A$	$D-\text{H}$	$\text{H} \cdots A$	$D \cdots A$	$D-\text{H} \cdots A$
O2—H2 \cdots O4 ⁱ	0.82	2.09	2.791 (2)	143

Symmetry code: (i) $-x, -y, -z + 1$.

Data collection: *CAD-4 EXPRESS* (Enraf–Nonius, 1994); cell refinement: *CAD-4 EXPRESS*; data reduction: *XCAD4* (Harms & Wocadlo, 1996); program(s) used to solve structure: *SHELXS97* (Sheldrick, 2008); program(s) used to refine structure: *SHELXL97* (Sheldrick, 2008); molecular graphics: *PLATON* (Spek, 2009); software used to prepare material for publication: *SHELXL97*.

JKS thanks the UGC for an RFSMS fellowship.

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: GK2301).

References

- Enraf–Nonius (1994). *CAD-4 EXPRESS*. Enraf–Nonius, Delft, The Netherlands.
- Harms, K. & Wocadlo, S. (1996). *XCAD4*. University of Marburg, Germany.
- North, A. C. T., Phillips, D. C. & Mathews, F. S. (1968). *Acta Cryst. A* **24**, 351–359.
- Pentala, N. R., Reddy, T. R. Y., Parkin, S. & Crooks, P. A. (2009). *Acta Cryst. E* **65**, o1877.
- Sheldrick, G. M. (2008). *Acta Cryst. A* **64**, 112–122.
- Spek, A. L. (2009). *Acta Cryst. D* **65**, 148–155.

supplementary materials

Acta Cryst. (2010). E66, o2967 [doi:10.1107/S1600536810042856]

***rac*-2-Hydroxy-2-(2-oxocyclopentyl)-1*H*-indene-1,3(2*H*)-dione**

J. K. Sundar, S. Maharani, R. R. Kumar, S. Natarajan, J. Suresh and P. L. N. Lakshman

Comment

Ninhydrin derivatives constitute a versatile class of compounds with profound biological activities such as antibacterial, anticonvulsant, anticancer and anti-inflammatory activities. The present work constitutes the synthesis of various ninhydrin derivatives which are being tested for anti-tubercular and other biological activities.

Experimental

A mixture of cyclopentanone (0.5 g, 0.006 mol) and ninhydrin (1.05 g, 0.006 mol) in methanol (10 ml) was heated under reflux for 4 h in the presence of solid sodium ethoxide (0.4 g, 0.006 mol). After completion of the reaction, as was evident from TLC, the reaction mixture was poured into crushed ice, extracted with dichloromethane and subjected to column chromatographic purification using petroleum ether:ethyl acetate mixture (90:10 v/v) to obtain the product in 50% yield. The compound was further recrystallized from ethyl acetate to obtain suitable crystals for X-ray studies (m.p. 543 K).

Refinement

All H atoms were placed at calculated positions and allowed to ride on their carrier atoms with C—H = 0.93–0.97 Å, $U_{\text{iso}} = 1.2U_{\text{eq}}(\text{C})$ and O—H = 0.82 Å, and $U_{\text{iso}} = 1.5U_{\text{eq}}(\text{O})$.

Figures




Fig. 1. The molecular structure of the title compound showing 30% probability displacement ellipsoids and the atom-numbering scheme.




Fig. 2. The packing diagram of the title compound.

***rac*-2-Hydroxy-2-(2-oxocyclopentyl)-1*H*-indene-1,3(2*H*)-dione**

Crystal data

$\text{C}_{14}\text{H}_{12}\text{O}_4$

$Z = 2$

$M_r = 244.24$

$F(000) = 256$

supplementary materials

Triclinic, $P\bar{1}$	$D_x = 1.384 \text{ Mg m}^{-3}$
Hall symbol: -P 1	Mo $K\alpha$ radiation, $\lambda = 0.71069 \text{ \AA}$
$a = 8.044 (3) \text{ \AA}$	Cell parameters from 25 reflections
$b = 8.404 (4) \text{ \AA}$	$\theta = 2\text{--}25^\circ$
$c = 10.239 (3) \text{ \AA}$	$\mu = 0.10 \text{ mm}^{-1}$
$\alpha = 66.95 (3)^\circ$	$T = 293 \text{ K}$
$\beta = 74.36 (2)^\circ$	Block, colourless
$\gamma = 68.50 (3)^\circ$	$0.26 \times 0.22 \times 0.19 \text{ mm}$
$V = 586.1 (4) \text{ \AA}^3$	

Data collection

Nonius MACH3 diffractometer	1886 reflections with $I > 2\sigma(I)$
Radiation source: fine-focus sealed tube	$R_{\text{int}} = 0.016$
graphite	$\theta_{\text{max}} = 25.0^\circ, \theta_{\text{min}} = 2.2^\circ$
$\omega\text{-}2\theta$ scans	$h = -1 \rightarrow 9$
Absorption correction: ψ scan (North <i>et al.</i> , 1968)	$k = -9 \rightarrow 9$
$T_{\text{min}} = 0.974, T_{\text{max}} = 0.981$	$l = -11 \rightarrow 12$
2534 measured reflections	3 standard reflections every 60 min
2054 independent reflections	intensity decay: none

Refinement

Refinement on F^2	Secondary atom site location: difference Fourier map
Least-squares matrix: full	Hydrogen site location: inferred from neighbouring sites
$R[F^2 > 2\sigma(F^2)] = 0.038$	H-atom parameters constrained
$wR(F^2) = 0.104$	$w = 1/[\sigma^2(F_o^2) + (0.0487P)^2 + 0.2034P]$ where $P = (F_o^2 + 2F_c^2)/3$
$S = 1.07$	$(\Delta/\sigma)_{\text{max}} < 0.001$
2054 reflections	$\Delta\rho_{\text{max}} = 0.25 \text{ e \AA}^{-3}$
164 parameters	$\Delta\rho_{\text{min}} = -0.23 \text{ e \AA}^{-3}$
0 restraints	Extinction correction: <i>SHELXL97</i> (Sheldrick, 2008), $F_c^* = kF_c[1 + 0.001xF_c^2\lambda^3/\sin(2\theta)]^{-1/4}$
Primary atom site location: structure-invariant direct methods	Extinction coefficient: 0.059 (7)

Special details

Geometry. All e.s.d.'s (except the e.s.d. in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell e.s.d.'s are taken into account individually in the estimation of e.s.d.'s in distances, angles and torsion angles; correlations between e.s.d.'s in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell e.s.d.'s is used for estimating e.s.d.'s involving l.s. planes.

Refinement. Refinement of F^2 against ALL reflections. The weighted R -factor wR and goodness of fit S are based on F^2 , conventional R -factors R are based on F , with F set to zero for negative F^2 . The threshold expression of $F^2 > \sigma(F^2)$ is used only for calculating R -

factors(gt) etc. and is not relevant to the choice of reflections for refinement. *R*-factors based on F^2 are statistically about twice as large as those based on F , and *R*-factors based on ALL data will be even larger.

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (\AA^2)

	<i>x</i>	<i>y</i>	<i>z</i>	$U_{\text{iso}}^*/U_{\text{eq}}$
C1	0.4901 (2)	0.0143 (2)	0.33717 (14)	0.0367 (4)
C2	0.6584 (2)	-0.1372 (2)	0.32374 (19)	0.0505 (4)
H2A	0.7411	-0.1548	0.3852	0.061*
H2B	0.7191	-0.1129	0.2254	0.061*
C3	0.5929 (3)	-0.3013 (2)	0.3708 (2)	0.0551 (5)
H3A	0.5821	-0.3580	0.4742	0.066*
H3B	0.6745	-0.3896	0.3258	0.066*
C4	0.4084 (2)	-0.2245 (2)	0.3207 (2)	0.0494 (4)
H4A	0.3353	-0.3064	0.3719	0.059*
H4B	0.4206	-0.2008	0.2184	0.059*
C5	0.3268 (2)	-0.04868 (19)	0.35711 (15)	0.0367 (4)
H5	0.2797	-0.0818	0.4598	0.044*
C6	0.17097 (19)	0.09543 (19)	0.27852 (15)	0.0353 (3)
C7	0.2311 (2)	0.1817 (2)	0.11679 (15)	0.0387 (4)
C8	0.2136 (2)	0.3728 (2)	0.09029 (16)	0.0379 (4)
C9	0.2631 (2)	0.5004 (2)	-0.03520 (18)	0.0502 (4)
H9	0.3169	0.4710	-0.1186	0.060*
C10	0.2304 (3)	0.6721 (2)	-0.0329 (2)	0.0573 (5)
H10	0.2618	0.7600	-0.1163	0.069*
C11	0.1513 (3)	0.7168 (2)	0.0916 (2)	0.0561 (5)
H11	0.1308	0.8338	0.0902	0.067*
C12	0.1028 (2)	0.5900 (2)	0.21723 (19)	0.0470 (4)
H12	0.0508	0.6193	0.3008	0.056*
C13	0.13396 (19)	0.4179 (2)	0.21488 (16)	0.0366 (3)
C14	0.09696 (19)	0.2590 (2)	0.33215 (15)	0.0362 (3)
O1	0.48152 (16)	0.16243 (15)	0.33474 (12)	0.0484 (3)
O2	0.03172 (15)	0.02227 (15)	0.29221 (13)	0.0487 (3)
H2	-0.0080	-0.0170	0.3773	0.073*
O3	0.2801 (2)	0.10548 (17)	0.02872 (13)	0.0602 (4)
O4	0.01604 (16)	0.25544 (16)	0.45179 (12)	0.0502 (3)

Atomic displacement parameters (\AA^2)

	U^{11}	U^{22}	U^{33}	U^{12}	U^{13}	U^{23}
C1	0.0390 (8)	0.0451 (9)	0.0268 (7)	-0.0147 (7)	-0.0051 (6)	-0.0103 (6)
C2	0.0385 (9)	0.0636 (11)	0.0529 (10)	-0.0090 (8)	-0.0067 (7)	-0.0282 (9)
C3	0.0563 (11)	0.0472 (10)	0.0590 (11)	0.0008 (8)	-0.0178 (8)	-0.0231 (8)
C4	0.0534 (10)	0.0382 (9)	0.0599 (10)	-0.0117 (7)	-0.0158 (8)	-0.0162 (7)
C5	0.0392 (8)	0.0369 (8)	0.0330 (7)	-0.0144 (6)	-0.0043 (6)	-0.0082 (6)
C6	0.0368 (8)	0.0355 (8)	0.0349 (7)	-0.0164 (6)	-0.0048 (6)	-0.0077 (6)
C7	0.0419 (8)	0.0409 (8)	0.0338 (8)	-0.0138 (6)	-0.0051 (6)	-0.0117 (6)
C8	0.0378 (8)	0.0392 (8)	0.0345 (7)	-0.0138 (6)	-0.0071 (6)	-0.0064 (6)

supplementary materials

C9	0.0581 (10)	0.0529 (10)	0.0353 (8)	-0.0246 (8)	-0.0039 (7)	-0.0043 (7)
C10	0.0668 (12)	0.0465 (10)	0.0520 (10)	-0.0299 (9)	-0.0110 (9)	0.0044 (8)
C11	0.0639 (11)	0.0356 (9)	0.0691 (12)	-0.0193 (8)	-0.0185 (9)	-0.0073 (8)
C12	0.0483 (9)	0.0397 (9)	0.0527 (10)	-0.0113 (7)	-0.0083 (7)	-0.0158 (7)
C13	0.0335 (7)	0.0367 (8)	0.0372 (8)	-0.0108 (6)	-0.0057 (6)	-0.0087 (6)
C14	0.0312 (7)	0.0400 (8)	0.0343 (8)	-0.0118 (6)	-0.0028 (6)	-0.0091 (6)
O1	0.0516 (7)	0.0463 (7)	0.0537 (7)	-0.0207 (5)	-0.0116 (5)	-0.0145 (5)
O2	0.0464 (7)	0.0514 (7)	0.0525 (7)	-0.0268 (5)	-0.0123 (5)	-0.0066 (5)
O3	0.0894 (10)	0.0528 (7)	0.0402 (6)	-0.0226 (7)	-0.0035 (6)	-0.0195 (6)
O4	0.0528 (7)	0.0511 (7)	0.0383 (6)	-0.0173 (5)	0.0087 (5)	-0.0143 (5)

Geometric parameters (\AA , $^\circ$)

C1—O1	1.2116 (19)	C6—C7	1.545 (2)
C1—C2	1.499 (2)	C7—O3	1.2059 (19)
C1—C5	1.526 (2)	C7—C8	1.479 (2)
C2—C3	1.512 (3)	C8—C9	1.384 (2)
C2—H2A	0.9700	C8—C13	1.393 (2)
C2—H2B	0.9700	C9—C10	1.378 (3)
C3—C4	1.528 (3)	C9—H9	0.9300
C3—H3A	0.9700	C10—C11	1.391 (3)
C3—H3B	0.9700	C10—H10	0.9300
C4—C5	1.531 (2)	C11—C12	1.379 (3)
C4—H4A	0.9700	C11—H11	0.9300
C4—H4B	0.9700	C12—C13	1.383 (2)
C5—C6	1.534 (2)	C12—H12	0.9300
C5—H5	0.9800	C13—C14	1.469 (2)
C6—O2	1.4160 (18)	C14—O4	1.2169 (18)
C6—C14	1.536 (2)	O2—H2	0.8200
O1—C1—C2	126.74 (15)	C5—C6—C14	110.99 (12)
O1—C1—C5	124.60 (14)	O2—C6—C7	107.82 (12)
C2—C1—C5	108.66 (13)	C5—C6—C7	113.02 (12)
C1—C2—C3	104.51 (14)	C14—C6—C7	102.49 (12)
C1—C2—H2A	110.9	O3—C7—C8	126.80 (14)
C3—C2—H2A	110.9	O3—C7—C6	125.30 (14)
C1—C2—H2B	110.9	C8—C7—C6	107.89 (13)
C3—C2—H2B	110.9	C9—C8—C13	120.56 (15)
H2A—C2—H2B	108.9	C9—C8—C7	129.09 (15)
C2—C3—C4	103.75 (14)	C13—C8—C7	110.35 (13)
C2—C3—H3A	111.0	C10—C9—C8	118.04 (17)
C4—C3—H3A	111.0	C10—C9—H9	121.0
C2—C3—H3B	111.0	C8—C9—H9	121.0
C4—C3—H3B	111.0	C9—C10—C11	121.33 (16)
H3A—C3—H3B	109.0	C9—C10—H10	119.3
C3—C4—C5	102.74 (13)	C11—C10—H10	119.3
C3—C4—H4A	111.2	C12—C11—C10	120.90 (17)
C5—C4—H4A	111.2	C12—C11—H11	119.6
C3—C4—H4B	111.2	C10—C11—H11	119.6
C5—C4—H4B	111.2	C11—C12—C13	117.90 (17)

H4A—C4—H4B	109.1	C11—C12—H12	121.1
C1—C5—C4	103.78 (13)	C13—C12—H12	121.1
C1—C5—C6	114.77 (12)	C12—C13—C8	121.28 (14)
C4—C5—C6	117.09 (13)	C12—C13—C14	129.02 (15)
C1—C5—H5	106.9	C8—C13—C14	109.69 (14)
C4—C5—H5	106.9	O4—C14—C13	126.49 (15)
C6—C5—H5	106.9	O4—C14—C6	124.45 (13)
O2—C6—C5	111.26 (12)	C13—C14—C6	109.03 (12)
O2—C6—C14	110.92 (12)	C6—O2—H2	109.5
O1—C1—C2—C3	166.12 (15)	O3—C7—C8—C13	173.24 (16)
C5—C1—C2—C3	−13.03 (17)	C6—C7—C8—C13	−5.46 (17)
C1—C2—C3—C4	33.50 (18)	C13—C8—C9—C10	−0.3 (2)
C2—C3—C4—C5	−41.27 (18)	C7—C8—C9—C10	179.19 (16)
O1—C1—C5—C4	168.45 (14)	C8—C9—C10—C11	0.5 (3)
C2—C1—C5—C4	−12.37 (16)	C9—C10—C11—C12	−0.1 (3)
O1—C1—C5—C6	39.4 (2)	C10—C11—C12—C13	−0.5 (3)
C2—C1—C5—C6	−141.39 (14)	C11—C12—C13—C8	0.7 (2)
C3—C4—C5—C1	32.54 (16)	C11—C12—C13—C14	179.53 (15)
C3—C4—C5—C6	160.13 (14)	C9—C8—C13—C12	−0.3 (2)
C1—C5—C6—O2	174.96 (11)	C7—C8—C13—C12	−179.90 (14)
C4—C5—C6—O2	52.92 (18)	C9—C8—C13—C14	−179.32 (14)
C1—C5—C6—C14	−61.01 (16)	C7—C8—C13—C14	1.09 (17)
C4—C5—C6—C14	176.94 (12)	C12—C13—C14—O4	7.0 (3)
C1—C5—C6—C7	53.49 (17)	C8—C13—C14—O4	−174.07 (15)
C4—C5—C6—C7	−68.56 (17)	C12—C13—C14—C6	−175.15 (15)
O2—C6—C7—O3	−54.4 (2)	C8—C13—C14—C6	3.77 (16)
C5—C6—C7—O3	69.0 (2)	O2—C6—C14—O4	56.38 (19)
C14—C6—C7—O3	−171.52 (16)	C5—C6—C14—O4	−67.85 (19)
O2—C6—C7—C8	124.29 (13)	C7—C6—C14—O4	171.22 (14)
C5—C6—C7—C8	−112.31 (14)	O2—C6—C14—C13	−121.51 (13)
C14—C6—C7—C8	7.21 (15)	C5—C6—C14—C13	114.26 (13)
O3—C7—C8—C9	−6.3 (3)	C7—C6—C14—C13	−6.67 (15)
C6—C7—C8—C9	174.99 (15)		

Hydrogen-bond geometry (Å, °)

<i>D</i> —H··· <i>A</i>	<i>D</i> —H	H··· <i>A</i>	<i>D</i> ··· <i>A</i>	<i>D</i> —H··· <i>A</i>
O2—H2···O4 ⁱ	0.82	2.09	2.791 (2)	143
C2—H2A···O4 ⁱⁱ	0.97	2.60	3.560 (2)	171
C2—H2B···O3 ⁱⁱⁱ	0.97	2.58	3.423 (2)	146

Symmetry codes: (i) $-x, -y, -z+1$; (ii) $-x+1, -y, -z+1$; (iii) $-x+1, -y, -z$.

supplementary materials

Fig. 1




Fig. 2

