

Acta Crystallographica Section E

Structure Reports

Online

ISSN 1600-5368

2-(4-Formyl-2,6-dimethoxyphenoxy)-acetic acid

Alain Collas, Christophe M. L. Vande Velde† and Frank Blockhuys*

Department of Chemistry, University of Antwerp, Universiteitsplein 1, B-2610 Wilrijk, Belgium

Correspondence e-mail: frank.blockhuys@ua.ac.be


Received 22 October 2010; accepted 27 October 2010

Key indicators: single-crystal X-ray study; $T = 293$ K; mean $\sigma(\text{C}-\text{C}) = 0.003$ Å; disorder in main residue; R factor = 0.036; wR factor = 0.093; data-to-parameter ratio = 11.8.

In the title compound, $\text{C}_{11}\text{H}_{12}\text{O}_6$, the aldehyde group is disordered over two sites in a 0.79:0.21 ratio. The carboxylic acid chain is found in the $[ap,ap]$ conformation due to two intramolecular $\text{O}-\text{H}\cdots\text{O}$ hydrogen bonds.

Related literature

For related acetic acids substituted in the α -position, see: Lundquist *et al.* (1987). For conformational and geometric considerations of carboxylic acids, see: Lide (1964); Leiserowitz (1976). For applications of PPV oligomers, see: Chemla (1987); Bandyopadhyay & Pal (2003). For hydrogen bonding and crystal engineering, see: Desiraju (1997); Steiner (2002).



Experimental

Crystal data

$\text{C}_{11}\text{H}_{12}\text{O}_6$
 $M_r = 240.21$
 Monoclinic, $P2_1/c$
 $a = 9.350$ (2) Å
 $b = 7.416$ (1) Å
 $c = 17.374$ (8) Å
 $\beta = 113.67$ (3)°

$V = 1103.4$ (6) Å³
 $Z = 4$
 Mo $K\alpha$ radiation
 $\mu = 0.12$ mm⁻¹
 $T = 293$ K
 $0.27 \times 0.24 \times 0.15$ mm

Data collection

Enraf–Nonius MACH3
 diffractometer
 4025 measured reflections
 2015 independent reflections

1292 reflections with $I > 2\sigma(I)$
 $R_{\text{int}} = 0.030$
 3 standard reflections every 60 min
 intensity decay: 1%

Refinement

$R[F^2 > 2\sigma(F^2)] = 0.036$
 $wR(F^2) = 0.093$
 $S = 1.01$
 2015 reflections
 171 parameters

8 restraints
 H-atom parameters constrained
 $\Delta\rho_{\text{max}} = 0.15$ e Å⁻³
 $\Delta\rho_{\text{min}} = -0.15$ e Å⁻³

Table 1

Hydrogen-bond geometry (Å, °).

$D-\text{H}\cdots A$	$D-\text{H}$	$\text{H}\cdots A$	$D\cdots A$	$D-\text{H}\cdots A$
O42—H42 ⁱ ···O4	0.82	2.11	2.617 (2)	120
O42—H42 ⁱ ···O5	0.82	2.18	2.932 (2)	153
C6—H6 ⁱ ···O41 ⁱ	0.93	2.54	3.468 (3)	176
C41—H41A ⁱ ···O11B ⁱⁱ	0.97	2.71	3.189 (9)	111
C51—H51B ⁱ ···O11A ⁱⁱⁱ	0.96	2.57	3.369 (3)	141
C31—H31B ⁱ ···O11A ^{iv}	0.96	2.70	3.454 (3)	136
C41—H41B ⁱ ···O4 ^v	0.97	2.56	3.527 (3)	173

Symmetry codes: (i) $x, -y + \frac{3}{2}, z - \frac{1}{2}$; (ii) $x, -y + \frac{3}{2}, z + \frac{1}{2}$; (iii) $x - 1, y, z$; (iv) $-x + 2, y + \frac{1}{2}, -z + \frac{3}{2}$; (v) $-x + 1, y - \frac{1}{2}, -z + \frac{3}{2}$.

Data collection: *CAD-4 EXPRESS* (Enraf–Nonius, 1994); cell refinement: *CAD-4 EXPRESS*; data reduction: *XCAD4* (Harms & Wocadlo, 1996); program(s) used to solve structure: *SHELXS97* (Sheldrick, 2008); program(s) used to refine structure: *SHELXL97* (Sheldrick, 2008); molecular graphics: *ORTEP-3 for Windows* (Farrugia, 1997); software used to prepare material for publication: *WinGX* (Farrugia, 1999).

CMLVV thanks the Fund for Scientific Research (FWO Vlaanderen) for a grant as a research assistant. AC thanks the Institute for the Promotion of Innovation by Science and Technology in Flanders (IWT) for a predoctoral grant.

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: ZL2319).

References

- Bandyopadhyay, A. & Pal, A. J. (2003). *Appl. Phys. Lett.* **82**, 1215–1217.
 Chemla, D. S. (1987). *Nonlinear Optical Properties of Organic Molecules and Crystals*. Boston: Academic Press.
 Desiraju, G. R. (1997). *J. Chem. Soc. Chem. Commun.* pp. 1475–1482.
 Enraf–Nonius (1994). *CAD-4 EXPRESS*. Enraf–Nonius, Delft, The Netherlands.
 Farrugia, L. J. (1997). *J. Appl. Cryst.* **30**, 565.
 Farrugia, L. J. (1999). *J. Appl. Cryst.* **32**, 837–838.
 Harms, K. & Wocadlo, S. (1996). *XCAD4*. University of Marburg, Germany.
 Leiserowitz, L. (1976). *Acta Cryst.* **B32**, 775–802.
 Lide, D. R. Jr (1964). *Annu. Rev. Phys. Chem.* **15**, 225–250.
 Lundquist, K., Stomberg, R. & von Unge, S. (1987). *Acta Chem. Scand. B*, **41**, 499–510.
 Sheldrick, G. M. (2008). *Acta Cryst.* **A64**, 112–122.
 Spek, A. L. (2009). *Acta Cryst.* **D65**, 148–155.
 Steiner, T. (2002). *Angew. Chem. Int. Ed.* **41**, 48–76.

† Current address: Karel de Grote University College, Department of Applied Engineering, Salesianenlaan 30, B-2660 Antwerp, Belgium.

supplementary materials

Acta Cryst. (2010). E66, o3003 [doi:10.1107/S1600536810043990]

2-(4-Formyl-2,6-dimethoxyphenoxy)acetic acid

A. Collas, C. M. L. Vande Velde and F. Blockhuys

Comment

The title compound was synthesized as a precursor for asymmetric PPV-type [poly(*p*-phenylene vinylene)] oligomers. These compounds are promising candidates as the active materials in organic memories (Bandyopadhyay & Pal, 2003) and as nonlinear optical (NLO) materials with a high first-order hyperpolarizability (Chemla, 1987). Besides the condition that these oligomers should bear acceptor and donor substituents connected by a π -system, it is of critical importance for their usefulness as an NLO material and as a bistable organic memory material that the crystal packing is non-centrosymmetric. In particular, the compound should crystallize in a polar space group. To engineer the crystal packing in order to meet this criterion, it is necessary to introduce certain statistically well chosen synthons (Desiraju, 1997). Therefore, we opted for carboxylic acid functional groups (powerful hydrogen bond donors) in the basic structure of the organic semiconductors. To combine the electronic (A- π -D) and the structural (non-centrosymmetric space group) requirements, we used the Williamson ether synthesis to prepare the title compound as a building block for a PPV-based semiconductor bearing a carboxylic acid moiety.

The carboxylic acid moiety is found in the [*ap,ap*] conformation (Fig. 1): the carboxyl H atom points in the opposite direction of the carbonyl group and the O4—C41—C42—O41 torsion angle is $-168.64(19)^\circ$. The reason for this unexpected conformation is the presence of an intramolecular bifurcated hydrogen bond involving H42, O4 and O5 (Table 1 and Fig. 2). Indeed, Lide showed, based on microwave experiments on gaseous formic acid, that, in general, the *sp* conformer is more stable than the *ap* conformer by about 16 kJ mol^{-1} (Lide, 1964). The acetic acid chain is twisted out-of-plane by about 113° and the torsion angle O4—C41—C42—O42 is about 11° .

A sawtooth motif is generated along the *c* axis by the CH \cdots O interaction between H6 and O41 (Fig. 2). Its effect is reinforced by H41A contacting O11B, the aldehyde O atom of the minor conformer. Perpendicular to these sawtooth ribbons (along the *a* axis) C51—H51B \cdots O11A hydrogen bonds continue the structure (Fig. 2). These corrugated sheets are then stacked along the *b* axis by two weak CH \cdots O hydrogen bonds and a CH \cdots π interaction (Fig. 3): H31B contacts O11A, H41B contacts O4 and H41A contacts the centroid of the ring [C41—H41A \cdots Cg^{vi}, 2.68 Å, 144° , symm. code vi = $1 - x, 1/2 + y, 3/2 - z$]. Finally, there is a modest π - π interaction [Cg \cdots Cg^{vii}, 4.133 (2) Å, 34.09° , symm. code vii = $1 - x, 2 - y, 1 - z$].

Experimental

Sodium hydroxide (5.0 g, 0.125 mol) in distilled water (20 ml) was added to a stirred solution of syringaldehyde (11.8 g, 0.065 mol) and iodoacetic acid (12.1 g, 0.065 mol) in distilled water (100 ml). The mixture was refluxed overnight and then poured into water (100 ml), which was then acidified using phosphoric acid. The mixture was left to cool in the refrigerator and the solvent was partially evaporated to precipitate the compound. The brown needle-like crystals were filtered off. The product changed colour to purple when it was exposed to air. The purple product was recrystallized from a 5:1 dichloromethane/acetone mixture to yield (54%) colourless crystals. $^1\text{H NMR}$ (CDCl_3 , 400 MHz, TMS): δ 3.98 (s, 6H,

supplementary materials

H31 and H51), 4.69 (s, 2H, H41), 7.16 (s, 2H, H2 and H6), 9.89 (s, 1H, H11). ^{13}C NMR (CDCl_3 , 100 MHz, TMS): δ 56.6 (C31 and C51), 70.3, (C41), 106.6 (C2 and C6), 132.9 (C1), 141.1 (C4), 152.6 (C3 and C5), 170.2 (C42), 190.5 (C11).

Refinement

The aldehyde moiety proved to be orientationally disordered over two sites in a 79:21 ratio (refined values 0.785 (6):0.215 (6)). The structure was initially refined with the aldehyde group in one position and then the second site was located in the residual electron density map. For the two aldehyde functions, consisting of three atoms, O11A and O11B were restrained to have the similar anisotropic displacement parameters, and the C1—C11A/C1—C11B and C11A—O11A/C11B—O11B distances were restrained to be equal. C11A and C11B were constrained to possess identical anisotropic displacement parameters. H atoms H11A and H11B were placed on the geometrically calculated positions and refined as 'riding'.

Figures




Fig. 1. Molecular structure of the title compound with the numbering scheme. Displacement ellipsoids are drawn at the 50% probability level and H atoms are represented as spheres with an arbitrary radius.




Fig. 2. Corrugated sheets in the *ac* plane generated by weak intermolecular hydrogen bonds and the bifurcated intramolecular hydrogen bond.




Fig. 3. The two weak hydrogen bonds and the $\text{CH}\cdots\pi$ interaction responsible for the stacking of the sheets along the *b* axis.

2-(4-Formyl-2,6-dimethoxyphenoxy)acetic acid

Crystal data

$\text{C}_{11}\text{H}_{12}\text{O}_6$

$M_r = 240.21$

Monoclinic, $P2_1/c$

Hall symbol: $-P\ 2ybc$

$a = 9.350$ (2) Å

$b = 7.416$ (1) Å

$c = 17.374$ (8) Å

$\beta = 113.67$ (3)°

$F(000) = 504$

$D_x = 1.446$ Mg m $^{-3}$

Melting point: 415 K

Mo $K\alpha$ radiation, $\lambda = 0.71073$ Å

Cell parameters from 25 reflections

$\theta = 9.0$ – 14.2 °

$\mu = 0.12$ mm $^{-1}$

$T = 293$ K

$V = 1103.4 (6) \text{ \AA}^3$
 $Z = 4$

Block, colourless
 $0.27 \times 0.24 \times 0.15 \text{ mm}$

Data collection

Enraf–Nonius MACH3
 diffractometer

$R_{\text{int}} = 0.030$

Radiation source: fine-focus sealed tube
 graphite

$\theta_{\text{max}} = 25.3^\circ$, $\theta_{\text{min}} = 2.4^\circ$

$\omega/2\theta$ scans

$h = -11 \rightarrow 11$

4025 measured reflections

$k = 0 \rightarrow 8$

2015 independent reflections

$l = -20 \rightarrow 20$

1292 reflections with $I > 2\sigma(I)$

3 standard reflections every 60 min

intensity decay: 1%

Refinement

Refinement on F^2

Secondary atom site location: difference Fourier map

Least-squares matrix: full

Hydrogen site location: inferred from neighbouring sites

$R[F^2 > 2\sigma(F^2)] = 0.036$

H-atom parameters constrained

$wR(F^2) = 0.093$

$w = 1/[\sigma^2(F_o^2) + (0.0427P)^2 + 0.1341P]$

$S = 1.00$

where $P = (F_o^2 + 2F_c^2)/3$

2015 reflections

$(\Delta/\sigma)_{\text{max}} < 0.001$

171 parameters

$\Delta\rho_{\text{max}} = 0.15 \text{ e \AA}^{-3}$

8 restraints

$\Delta\rho_{\text{min}} = -0.15 \text{ e \AA}^{-3}$

Primary atom site location: structure-invariant direct methods

Extinction correction: *SHELXL97* (Sheldrick, 2008),

$F_c^* = kFc[1 + 0.001xFc^2\lambda^3/\sin(2\theta)]^{-1/4}$

Extinction coefficient: 0.0086 (18)

Special details

Geometry. All e.s.d.'s (except the e.s.d. in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell e.s.d.'s are taken into account individually in the estimation of e.s.d.'s in distances, angles and torsion angles; correlations between e.s.d.'s in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell e.s.d.'s is used for estimating e.s.d.'s involving l.s. planes.

Refinement. Refinement of F^2 against ALL reflections. The weighted R -factor wR and goodness of fit S are based on F^2 , conventional R -factors R are based on F , with F set to zero for negative F^2 . The threshold expression of $F^2 > \sigma(F^2)$ is used only for calculating R -factors(gt) *etc.* and is not relevant to the choice of reflections for refinement. R -factors based on F^2 are statistically about twice as large as those based on F , and R -factors based on ALL data will be even larger.

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (\AA^2)

	x	y	z	$U_{\text{iso}}^*/U_{\text{eq}}$	Occ. (<1)
O11A	0.8177 (2)	0.6876 (3)	0.52772 (13)	0.0677 (9)	0.785 (6)
C11A	0.6811 (5)	0.6988 (12)	0.5115 (3)	0.0490 (13)	0.785 (6)
H11A	0.6121	0.6482	0.4613	0.059*	0.785 (6)

supplementary materials

C11B	0.720 (3)	0.712 (5)	0.5283 (16)	0.0490 (13)	0.215 (6)
H11B	0.8278	0.7181	0.5571	0.059*	0.215 (6)
O11B	0.6629 (10)	0.6437 (14)	0.4603 (6)	0.078 (3)	0.215 (6)
O4	0.39907 (13)	1.04247 (17)	0.70196 (7)	0.0376 (3)	
O5	0.22382 (14)	0.89227 (19)	0.55872 (8)	0.0449 (4)	
O3	0.71034 (14)	1.0208 (2)	0.76755 (8)	0.0478 (4)	
C6	0.4523 (2)	0.7946 (3)	0.53433 (11)	0.0388 (5)	
H6	0.3923	0.7439	0.4824	0.047*	
C2	0.7062 (2)	0.8612 (3)	0.64352 (11)	0.0414 (5)	
H2	0.8144	0.8545	0.6638	0.050*	
C5	0.38175 (19)	0.8774 (2)	0.58135 (10)	0.0345 (4)	
O42	0.13206 (15)	0.9328 (2)	0.70058 (9)	0.0604 (5)	
H42	0.1655	0.9578	0.6649	0.091*	
C4	0.47318 (19)	0.9529 (2)	0.65853 (10)	0.0334 (4)	
O41	0.22968 (17)	0.8754 (2)	0.83608 (9)	0.0623 (5)	
C1	0.6143 (2)	0.7884 (3)	0.56601 (11)	0.0392 (5)	
C41	0.4098 (2)	0.9607 (3)	0.77859 (11)	0.0380 (4)	
H41A	0.4634	1.0416	0.8252	0.046*	
H41B	0.4701	0.8502	0.7880	0.046*	
C3	0.6358 (2)	0.9442 (2)	0.69064 (11)	0.0369 (4)	
C42	0.2502 (2)	0.9191 (3)	0.77494 (12)	0.0429 (5)	
C51	0.1233 (2)	0.8060 (3)	0.48261 (12)	0.0551 (6)	
H51A	0.1374	0.8599	0.4359	0.083*	
H51B	0.0167	0.8202	0.4755	0.083*	
H51C	0.1484	0.6800	0.4853	0.083*	
C31	0.8750 (3)	0.9877 (4)	0.80939 (15)	0.0730 (7)	
H31A	0.8941	0.8602	0.8120	0.110*	
H31B	0.9128	1.0359	0.8653	0.110*	
H31C	0.9282	1.0449	0.7788	0.110*	

Atomic displacement parameters (\AA^2)

	U^{11}	U^{22}	U^{33}	U^{12}	U^{13}	U^{23}
O11A	0.0510 (16)	0.0877 (17)	0.0786 (16)	0.0113 (11)	0.0409 (12)	0.0007 (12)
C11A	0.050 (3)	0.052 (2)	0.049 (3)	0.002 (3)	0.025 (3)	0.002 (2)
C11B	0.050 (3)	0.052 (2)	0.049 (3)	0.002 (3)	0.025 (3)	0.002 (2)
O11B	0.084 (6)	0.101 (7)	0.053 (6)	0.007 (5)	0.031 (5)	-0.019 (5)
O4	0.0424 (7)	0.0359 (7)	0.0372 (7)	0.0054 (6)	0.0188 (6)	0.0017 (6)
O5	0.0315 (7)	0.0590 (9)	0.0409 (7)	0.0027 (6)	0.0109 (6)	-0.0073 (6)
O3	0.0327 (7)	0.0590 (9)	0.0445 (8)	-0.0028 (6)	0.0080 (6)	-0.0074 (7)
C6	0.0424 (11)	0.0409 (11)	0.0330 (9)	0.0001 (9)	0.0149 (8)	0.0011 (9)
C2	0.0340 (10)	0.0435 (11)	0.0503 (11)	0.0039 (9)	0.0206 (9)	0.0078 (9)
C5	0.0327 (9)	0.0366 (10)	0.0354 (9)	0.0020 (8)	0.0148 (8)	0.0043 (8)
O42	0.0361 (7)	0.0922 (13)	0.0540 (9)	0.0028 (8)	0.0193 (7)	0.0085 (9)
C4	0.0364 (9)	0.0309 (10)	0.0375 (9)	0.0034 (8)	0.0196 (8)	0.0026 (8)
O41	0.0623 (9)	0.0806 (12)	0.0556 (9)	0.0015 (9)	0.0359 (8)	0.0100 (8)
C1	0.0452 (10)	0.0375 (11)	0.0438 (11)	0.0053 (9)	0.0271 (9)	0.0068 (9)
C41	0.0399 (10)	0.0419 (11)	0.0321 (9)	0.0023 (9)	0.0142 (8)	-0.0017 (8)

C3	0.0349 (9)	0.0359 (10)	0.0394 (10)	-0.0019 (8)	0.0144 (8)	0.0036 (9)
C42	0.0434 (11)	0.0454 (12)	0.0433 (11)	0.0047 (9)	0.0210 (9)	0.0012 (9)
C51	0.0376 (11)	0.0738 (16)	0.0470 (12)	-0.0074 (11)	0.0095 (9)	-0.0118 (11)
C31	0.0452 (13)	0.0778 (18)	0.0681 (15)	0.0078 (12)	-0.0066 (11)	-0.0103 (14)

Geometric parameters (Å, °)

O11A—C11A	1.195 (5)	C2—C1	1.384 (3)
C11A—C1	1.485 (4)	C2—H2	0.9300
C11A—H11A	0.9300	C5—C4	1.386 (2)
C11B—O11B	1.196 (17)	O42—C42	1.324 (2)
C11B—C1	1.498 (16)	O42—H42	0.8200
C11B—H11B	0.9300	C4—C3	1.395 (2)
O4—C4	1.382 (2)	O41—C42	1.197 (2)
O4—C41	1.430 (2)	C41—C42	1.500 (3)
O5—C5	1.371 (2)	C41—H41A	0.9700
O5—C51	1.430 (2)	C41—H41B	0.9700
O3—C3	1.359 (2)	C51—H51A	0.9600
O3—C31	1.435 (3)	C51—H51B	0.9600
C6—C5	1.383 (2)	C51—H51C	0.9600
C6—C1	1.389 (3)	C31—H31A	0.9600
C6—H6	0.9300	C31—H31B	0.9600
C2—C3	1.383 (3)	C31—H31C	0.9600
O11A—C11A—C1	124.3 (3)	C6—C1—C11B	130.1 (9)
O11A—C11A—H11A	117.8	O4—C41—C42	110.56 (15)
C1—C11A—H11A	117.8	O4—C41—H41A	109.5
O11B—C11B—C1	118.8 (19)	C42—C41—H41A	109.5
O11B—C11B—H11B	120.6	O4—C41—H41B	109.5
C1—C11B—H11B	120.6	C42—C41—H41B	109.5
C4—O4—C41	116.19 (13)	H41A—C41—H41B	108.1
C5—O5—C51	117.50 (14)	O3—C3—C2	126.16 (16)
C3—O3—C31	116.68 (16)	O3—C3—C4	114.84 (16)
C5—C6—C1	118.91 (17)	C2—C3—C4	119.00 (16)
C5—C6—H6	120.5	O41—C42—O42	121.25 (19)
C1—C6—H6	120.5	O41—C42—C41	121.93 (18)
C3—C2—C1	119.53 (17)	O42—C42—C41	116.83 (16)
C3—C2—H2	120.2	O5—C51—H51A	109.5
C1—C2—H2	120.2	O5—C51—H51B	109.5
O5—C5—C6	125.46 (16)	H51A—C51—H51B	109.5
O5—C5—C4	114.83 (15)	O5—C51—H51C	109.5
C6—C5—C4	119.71 (15)	H51A—C51—H51C	109.5
C42—O42—H42	109.5	H51B—C51—H51C	109.5
O4—C4—C5	118.19 (15)	O3—C31—H31A	109.5
O4—C4—C3	120.58 (16)	O3—C31—H31B	109.5
C5—C4—C3	121.20 (16)	H31A—C31—H31B	109.5
C2—C1—C6	121.65 (17)	O3—C31—H31C	109.5
C2—C1—C11A	122.7 (2)	H31A—C31—H31C	109.5
C6—C1—C11A	115.7 (2)	H31B—C31—H31C	109.5
C2—C1—C11B	108.2 (9)		

supplementary materials

C51—O5—C5—C6	3.9 (3)	O11A—C11A—C1—C6	177.2 (6)
C51—O5—C5—C4	-175.66 (17)	O11A—C11A—C1—C11B	0(7)
C1—C6—C5—O5	-179.30 (17)	O11B—C11B—C1—C2	-179 (3)
C1—C6—C5—C4	0.2 (3)	O11B—C11B—C1—C6	0(4)
C41—O4—C4—C5	112.81 (18)	O11B—C11B—C1—C11A	3(4)
C41—O4—C4—C3	-69.2 (2)	C4—O4—C41—C42	-120.89 (17)
O5—C5—C4—O4	-3.6 (2)	C31—O3—C3—C2	-10.5 (3)
C6—C5—C4—O4	176.83 (16)	C31—O3—C3—C4	169.40 (19)
O5—C5—C4—C3	178.46 (16)	C1—C2—C3—O3	179.70 (17)
C6—C5—C4—C3	-1.1 (3)	C1—C2—C3—C4	-0.2 (3)
C3—C2—C1—C6	-0.7 (3)	O4—C4—C3—O3	3.3 (2)
C3—C2—C1—C11A	179.4 (5)	C5—C4—C3—O3	-178.83 (17)
C3—C2—C1—C11B	178.6 (17)	O4—C4—C3—C2	-176.79 (17)
C5—C6—C1—C2	0.6 (3)	C5—C4—C3—C2	1.1 (3)
C5—C6—C1—C11A	-179.4 (4)	O4—C41—C42—O41	-168.63 (19)
C5—C6—C1—C11B	-178 (2)	O4—C41—C42—O42	11.4 (2)
O11A—C11A—C1—C2	-2.9 (10)		

Hydrogen-bond geometry (\AA , $^\circ$)

<i>D</i> —H \cdots <i>A</i>	<i>D</i> —H	H \cdots <i>A</i>	<i>D</i> \cdots <i>A</i>	<i>D</i> —H \cdots <i>A</i>
O42—H42 \cdots O4	0.82	2.11	2.617 (2)	120
O42—H42 \cdots O5	0.82	2.18	2.932 (2)	153
C6—H6 \cdots O41 ⁱ	0.93	2.54	3.468 (3)	176
C41—H41A \cdots O11B ⁱⁱ	0.97	2.71	3.189 (9)	111
C51—H51B \cdots O11A ⁱⁱⁱ	0.96	2.57	3.369 (3)	141
C31—H31B \cdots O11A ^{iv}	0.96	2.70	3.454 (3)	136
C41—H41B \cdots O4 ^v	0.97	2.56	3.527 (3)	173

Symmetry codes: (i) $x, -y+3/2, z-1/2$; (ii) $x, -y+3/2, z+1/2$; (iii) $x-1, y, z$; (iv) $-x+2, y+1/2, -z+3/2$; (v) $-x+1, y-1/2, -z+3/2$.

Fig. 1




Fig. 2




Fig. 3

