

N-Benzyl-2,3,4,5,6-pentafluorobenzamide

Arto Valkonen, Tanja Lahtinen and Kari Rissanen*

Nanoscience Center, Department of Chemistry, University of Jyväskylä, PO Box 35, FIN-40014 University of Jyväskylä, Finland
Correspondence e-mail: kari.t.rissanen@jyu.fi


Received 15 October 2010; accepted 24 October 2010

Key indicators: single-crystal X-ray study; $T = 123$ K; mean $\sigma(\text{C}-\text{C}) = 0.002$ Å; R factor = 0.033; wR factor = 0.086; data-to-parameter ratio = 11.2.

In the title compound, $\text{C}_{14}\text{H}_8\text{F}_5\text{NO}$, the dihedral angle between the planes of the pentafluorophenyl and phenyl rings is $18.34(5)^\circ$. An intermolecular $\text{N}-\text{H}\cdots\text{O}$ hydrogen bond between the amide groups connects these molecules to form an infinite chain through the crystal structure. One weak intermolecular $\text{C}-\text{H}\cdots\text{O}$ contact and one $\pi-\pi$ interaction [centroid-centroid distance = $3.772(3)$ Å] are also involved in crystal structure stabilization between the phenyl rings.

Related literature

For related structures, see: An & Rhee (2003); Cockroft *et al.* (2007); Forbes *et al.* (2001); Liu *et al.* (2007); Qadeer *et al.* (2007); Zhang & Zhang (2008). For anion $\cdots\pi$ interactions, see: Albrecht *et al.* (2010); Lahtinen & Rissanen (2007); Müller *et al.* (2010).



Experimental

Crystal data

$\text{C}_{14}\text{H}_8\text{F}_5\text{NO}$
 $M_r = 301.21$
Monoclinic, $P2_1/n$
 $a = 7.1649(2)$ Å
 $b = 22.9090(5)$ Å
 $c = 7.5363(1)$ Å
 $\beta = 99.205(2)^\circ$

$V = 1221.08(5)$ Å³
 $Z = 4$
Mo $K\alpha$ radiation
 $\mu = 0.16$ mm⁻¹
 $T = 123$ K
 $0.40 \times 0.28 \times 0.26$ mm

Data collection

Bruker Nonius KappaCCD with APEXII detector diffractometer
4246 measured reflections
2152 independent reflections
1891 reflections with $I > 2\sigma(I)$
 $R_{\text{int}} = 0.016$

Refinement

$R[F^2 > 2\sigma(F^2)] = 0.033$
 $wR(F^2) = 0.086$
 $S = 1.06$
2152 reflections
193 parameters
1 restraint
H atoms treated by a mixture of independent and constrained refinement
 $\Delta\rho_{\text{max}} = 0.20$ e Å⁻³
 $\Delta\rho_{\text{min}} = -0.18$ e Å⁻³

Table 1

Hydrogen-bond geometry (Å, °).

$D-\text{H}\cdots A$	$D-\text{H}$	$\text{H}\cdots A$	$D\cdots A$	$D-\text{H}\cdots A$
$\text{N8}-\text{H8}\cdots\text{O1}^{\text{i}}$	0.88 (1)	2.01 (1)	2.875 (2)	171 (2)
$\text{C5}-\text{H5}\cdots\text{O1}^{\text{ii}}$	0.95	2.37	3.276 (2)	158

Symmetry codes: (i) $x + \frac{1}{2}, -y + \frac{1}{2}, z + \frac{1}{2}$; (ii) $x, y, z + 1$.

Data collection: COLLECT (Bruker, 2008); cell refinement: DENZO-SMN (Otwinowski & Minor, 1997); data reduction: DENZO-SMN; program(s) used to solve structure: SIR2004 (Burla *et al.*, 2005); program(s) used to refine structure: SHELXL97 (Sheldrick, 2008); molecular graphics: ORTEP-3 (Farrugia, 1997) and Mercury (Macrae *et al.*, 2008); software used to prepare material for publication: SHELXL97.

This work was supported by the Academy of Finland (KR, project No. 212588).

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: BT5380).

References

- Albrecht, M., Müller, M., Mergel, O., Rissanen, K. & Valkonen, A. (2010). *Chem. Eur. J.* **16**, 5062–5069.
- An, G. & Rhee, H. (2003). *Synlett*, pp. 876–878.
- Bruker (2008). COLLECT. Bruker AXS Inc., Delft, The Netherlands.
- Burla, M. C., Caliandro, R., Camalli, M., Carrozzini, B., Cascarano, G. L., De Caro, L., Giacovazzo, C., Polidori, G. & Spagna, R. (2005). *J. Appl. Cryst.* **38**, 381–388.
- Cockroft, S. L., Perkins, J., Zonta, C., Adams, H., Spey, S. E., Low, C. M. R., Vinter, J. G., Lawson, K. R., Urch, C. J. & Hunter, C. A. (2007). *Org. Biomol. Chem.* **5**, 1062–1080.
- Farrugia, L. J. (1997). *J. Appl. Cryst.* **30**, 565.
- Forbes, C. C., Beatty, A. M. & Smith, B. D. (2001). *Org. Lett.* **3**, 3595–3598.
- Lahtinen, T. & Rissanen, K. (2007). *Acta Cryst.* **E63**, o4114.
- Liu, S.-L., Liang, E.-X., Yu, L.-C. & Huang, L. (2007). *Z. Kristallogr. New Cryst. Struct.* **222**, 433–434.
- Macrae, C. F., Bruno, I. J., Chisholm, J. A., Edgington, P. R., McCabe, P., Pidcock, E., Rodriguez-Monge, L., Taylor, R., van de Streek, J. & Wood, P. A. (2008). *J. Appl. Cryst.* **41**, 466–470.
- Müller, M., Albrecht, M., Gossen, V., Peters, T., Hoffmann, A., Raabe, G., Valkonen, A. & Rissanen, K. (2010). *Chem. Eur. J.* In the press.
- Otwinowski, Z. & Minor, W. (1997). *Methods in Enzymology*, Vol. 276, *Macromolecular Crystallography*, Part A, edited by C. W. Carter & R. M. Sweet, pp. 307–326. New York: Academic Press.
- Qadeer, G., Rama, N. H. & Wong, W.-Y. (2007). *Acta Cryst.* **E63**, o335–o336.
- Sheldrick, G. M. (2008). *Acta Cryst.* **A64**, 112–122.
- Zhang, Q.-X. & Zhang, B.-S. (2008). *Acta Cryst.* **E64**, o884.

supplementary materials

Acta Cryst. (2010). E66, o3007 [doi:10.1107/S1600536810043345]

***N*-Benzyl-2,3,4,5,6-pentafluorobenzamide**

A. Valkonen, T. Lahtinen and K. Rissanen

Comment

The title compound is synthesized by classical amide formation reaction between amine (benzylamine) and carboxylic acid halide (2,3,4,5,6-pentafluorobenzoylchloride). The molecule of this secondary amide is not planar and contains two aromatic ring systems (Fig. 1), the other one being electron poor due to electron-withdrawing force of connected F atoms. This pentafluorophenyl moiety has recently been found to be an excellent halogen $\cdots\pi$ contact acceptor for halide and polyhalide anions in similar ammonium salt structures (Albrecht *et al.*, 2010; Müller *et al.*, 2010) and also acceptor for C=O \cdots C(aromatic) anion $\cdots\pi$ -type contacts (Lahtinen & Rissanen, 2007). The pentafluorophenyl ring is found to be inclined to phenyl ring by 18.34 (5)°. The C10/C9/N8/O1 amide group is more significantly inclined to pentafluorophenyl ring by 56.95 (4)° and to phenyl ring by 56.21 (4)°, as also observed, for example, with few substituted *N*-phenyl-2,3,4,5,6-pentafluorobenzamides (Cockroft *et al.*, 2007) and *N*-Benzyl-4,5-dimethoxy-2-nitrobenzamide (Qadeer *et al.*, 2007).

The intermolecular interactions of the title compound include one N—H \cdots O, one C—H \cdots O (Table 1) and one π – π contacts. The N—H \cdots O hydrogen bonds connect the molecules to form infinite chain in $(x + 1/2, -y + 1/2, z + 1/2)$ direction (Fig. 2), where every second molecule is in same orientation and every second is rotated 180° on the direction of *b* axis. Similar chain was obtained, for example, with *N*-Benzyl-4-phenylbenzamide (An & Rhee, 2003). These chains are connected to each other by one C—H \cdots O (Table 1) and one π – π contacts (Fig. 3), the latter having centroid-to-centroid distance of 3.772 (3) Å and closest C \cdots C distance of 3.327 (3) Å. These distances are slightly longer than in the structure of *N*-(2-pyridyl)-2,3,4,5,6-pentafluorobenzamide (Forbes *et al.*, 2001). The C=O \cdots C(aromatic) anion $\cdots\pi$ -type contact, found from the related structure of *N*-[1-(silatran-1-yl)propyl]pentafluorobenzamide (Lahtinen & Rissanen, 2007), seems to be in this case forced by nearby N—H \cdots O contact. Fluorines F3 and F4 (Fig. 1) show distance 2.920 (2) Å to F3 and F4 of the neighbouring molecule in $(-x, -y, -z + 1)$ direction, but this contact is most probably too weak to be significant in crystal stabilization.

Experimental

Benzylamine (184 mg, 1.72 mmol) and triethylamine (470 μ l, 3.44 mmol) were mixed in dry DCM under inert atmosphere (Ar). The reaction mixture was cooled (ice–salt bath) and 2,3,4,5,6-pentafluorobenzoylchloride (240 μ l, 1.72 mmol) in dry DCM was added dropwise to the reaction mixture. After addition the reaction mixture was stirred in ice–salt bath for 1 h and in room temperature for additional 20 h. The reaction mixture was washed twice with water and organic layer was dried and evaporated to yield white solid product. For the single-crystal X-ray analysis the crude product was recrystallized from CHCl₃ yielding colourless needles.

Refinement

All H atoms were visible in electron density maps, but those bonded to C were ideally positioned and allowed to ride on their parent atoms at C—H distances of 0.95 Å (aromatic) and 0.99 Å (methylene), with $U_{\text{iso}}(\text{H})$ of 1.2 times $U_{\text{eq}}(\text{C})$. The

supplementary materials

N—H proton were found in the electron density map and was refined with a distance restraint [$N—H = 0.88(2) \text{ \AA}$], and $U_{\text{iso}}(\text{H}) = 1.2$ times $U_{\text{eq}}(\text{N})$ was used.

Figures




Fig. 1. View of the molecule of the title compound showing the atom-labelling scheme. Displacement ellipsoids are drawn at the 50% probability level. H atoms are represented by circles of arbitrary size.




Fig. 2. Part of the infinite chain formed by $N—H\cdots O$ hydrogen bonds in the crystal of the title compound.




Fig. 3. Packing diagram showing the $N—H\cdots O$, $C—H\cdots O$ and $\pi-\pi$ contacts.

N-Benzyl-2,3,4,5,6-pentafluorobenzamide

Crystal data

$C_{14}H_8F_5NO$

$M_r = 301.21$

Monoclinic, $P2_1/n$

$a = 7.1649(2) \text{ \AA}$

$b = 22.9090(5) \text{ \AA}$

$c = 7.5363(1) \text{ \AA}$

$\beta = 99.205(2)^\circ$

$V = 1221.08(5) \text{ \AA}^3$

$Z = 4$

$F(000) = 608$

$D_x = 1.638 \text{ Mg m}^{-3}$

Mo $K\alpha$ radiation, $\lambda = 0.71073 \text{ \AA}$

Cell parameters from 3094 reflections

$\theta = 0.4\text{--}28.3^\circ$

$\mu = 0.16 \text{ mm}^{-1}$

$T = 123 \text{ K}$

Block, colourless

$0.40 \times 0.28 \times 0.26 \text{ mm}$

Data collection

Bruker Nonius KappaCCD with APEXII detector diffractometer

Radiation source: fine-focus sealed tube graphite

Detector resolution: 9 pixels mm^{-1}

φ and ω scans

4246 measured reflections

2152 independent reflections

1891 reflections with $I > 2\sigma(I)$

$R_{\text{int}} = 0.016$

$\theta_{\text{max}} = 25.0^\circ$, $\theta_{\text{min}} = 2.9^\circ$

$h = -8 \rightarrow 8$

$k = -27 \rightarrow 27$

$l = -8 \rightarrow 8$

Refinement

Refinement on F^2

Primary atom site location: structure-invariant direct methods

Least-squares matrix: full	Secondary atom site location: difference Fourier map
$R[F^2 > 2\sigma(F^2)] = 0.033$	Hydrogen site location: inferred from neighbouring sites
$wR(F^2) = 0.086$	H atoms treated by a mixture of independent and constrained refinement
$S = 1.06$	$w = 1/[\sigma^2(F_o^2) + (0.0393P)^2 + 0.498P]$
2152 reflections	where $P = (F_o^2 + 2F_c^2)/3$
193 parameters	$(\Delta/\sigma)_{\max} < 0.001$
1 restraint	$\Delta\rho_{\max} = 0.20 \text{ e } \text{\AA}^{-3}$
	$\Delta\rho_{\min} = -0.18 \text{ e } \text{\AA}^{-3}$

Special details

Geometry. All s.u.'s (except the s.u. in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell s.u.'s are taken into account individually in the estimation of s.u.'s in distances, angles and torsion angles; correlations between s.u.'s in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell s.u.'s is used for estimating s.u.'s involving l.s. planes.

Refinement. Refinement of F^2 against ALL reflections. The weighted R -factor wR and goodness of fit S are based on F^2 , conventional R -factors R are based on F , with F set to zero for negative F^2 . The threshold expression of $F^2 > 2\sigma(F^2)$ is used only for calculating R -factors(gt) *etc.* and is not relevant to the choice of reflections for refinement. R -factors based on F^2 are statistically about twice as large as those based on F , and R -factors based on ALL data will be even larger.

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (\AA^2)

	x	y	z	$U_{\text{iso}}^*/U_{\text{eq}}$
F1	0.11992 (14)	0.19154 (4)	-0.04946 (11)	0.0315 (2)
F2	0.17000 (14)	0.07495 (4)	-0.05119 (12)	0.0350 (3)
F3	0.14683 (15)	0.01169 (4)	0.24882 (14)	0.0379 (3)
F4	0.06766 (15)	0.06526 (4)	0.54949 (12)	0.0363 (3)
F5	0.00325 (13)	0.18090 (4)	0.54897 (11)	0.0277 (2)
O1	-0.10620 (15)	0.27306 (4)	0.12572 (13)	0.0260 (3)
N8	0.12659 (19)	0.28716 (5)	0.36339 (16)	0.0232 (3)
H8	0.218 (2)	0.2705 (7)	0.437 (2)	0.028*
C1	0.1073 (2)	0.37470 (6)	0.55552 (19)	0.0207 (3)
C2	0.1399 (2)	0.43426 (7)	0.5826 (2)	0.0240 (3)
H2	0.1652	0.4579	0.4856	0.029*
C3	0.1359 (2)	0.45945 (7)	0.7488 (2)	0.0278 (4)
H3	0.1570	0.5002	0.7649	0.033*
C4	0.1011 (2)	0.42535 (7)	0.8916 (2)	0.0277 (4)
H4	0.0983	0.4425	1.0059	0.033*
C5	0.0703 (2)	0.36600 (7)	0.8668 (2)	0.0262 (4)
H5	0.0474	0.3424	0.9648	0.031*
C6	0.0729 (2)	0.34078 (7)	0.6995 (2)	0.0226 (3)
H6	0.0509	0.3001	0.6835	0.027*
C7	0.1041 (2)	0.35043 (6)	0.3685 (2)	0.0262 (4)
H7A	-0.0174	0.3612	0.2935	0.031*
H7B	0.2068	0.3689	0.3149	0.031*

supplementary materials

C9	0.0197 (2)	0.25426 (6)	0.24216 (18)	0.0197 (3)
C10	0.0625 (2)	0.18984 (6)	0.25080 (18)	0.0197 (3)
C11	0.1035 (2)	0.16115 (7)	0.09949 (19)	0.0224 (3)
C12	0.1315 (2)	0.10168 (7)	0.0975 (2)	0.0249 (4)
C13	0.1202 (2)	0.06950 (7)	0.2499 (2)	0.0258 (4)
C14	0.0806 (2)	0.09667 (7)	0.4021 (2)	0.0253 (4)
C15	0.0511 (2)	0.15619 (7)	0.40118 (19)	0.0215 (3)

Atomic displacement parameters (\AA^2)

	U^{11}	U^{22}	U^{33}	U^{12}	U^{13}	U^{23}
F1	0.0414 (6)	0.0350 (5)	0.0190 (5)	-0.0003 (4)	0.0075 (4)	0.0023 (4)
F2	0.0382 (6)	0.0351 (5)	0.0326 (5)	0.0004 (4)	0.0089 (4)	-0.0129 (4)
F3	0.0420 (6)	0.0199 (5)	0.0515 (6)	0.0005 (4)	0.0070 (5)	-0.0003 (4)
F4	0.0463 (6)	0.0303 (5)	0.0315 (5)	-0.0016 (4)	0.0039 (4)	0.0131 (4)
F5	0.0343 (5)	0.0314 (5)	0.0176 (4)	-0.0017 (4)	0.0050 (4)	-0.0009 (4)
O1	0.0271 (6)	0.0248 (6)	0.0223 (5)	-0.0011 (5)	-0.0072 (5)	0.0037 (4)
N8	0.0258 (7)	0.0204 (7)	0.0203 (6)	0.0035 (5)	-0.0057 (5)	-0.0006 (5)
C1	0.0169 (7)	0.0222 (8)	0.0215 (7)	0.0018 (6)	-0.0015 (6)	0.0001 (6)
C2	0.0239 (8)	0.0241 (8)	0.0232 (8)	-0.0008 (6)	0.0016 (6)	0.0022 (6)
C3	0.0255 (8)	0.0233 (8)	0.0333 (9)	-0.0014 (7)	0.0006 (7)	-0.0045 (7)
C4	0.0249 (9)	0.0354 (9)	0.0227 (8)	0.0027 (7)	0.0034 (6)	-0.0067 (7)
C5	0.0217 (8)	0.0350 (9)	0.0222 (8)	0.0016 (7)	0.0046 (6)	0.0041 (6)
C6	0.0209 (8)	0.0201 (7)	0.0256 (8)	0.0005 (6)	0.0001 (6)	0.0019 (6)
C7	0.0345 (9)	0.0212 (8)	0.0212 (8)	0.0008 (7)	-0.0004 (6)	0.0009 (6)
C9	0.0191 (8)	0.0242 (8)	0.0158 (7)	-0.0019 (6)	0.0025 (6)	0.0024 (6)
C10	0.0148 (7)	0.0243 (8)	0.0186 (7)	-0.0017 (6)	-0.0020 (5)	-0.0001 (6)
C11	0.0193 (8)	0.0288 (8)	0.0184 (7)	-0.0028 (6)	0.0006 (6)	0.0028 (6)
C12	0.0188 (8)	0.0291 (8)	0.0262 (8)	-0.0014 (6)	0.0017 (6)	-0.0074 (7)
C13	0.0212 (8)	0.0192 (8)	0.0355 (9)	-0.0010 (6)	0.0000 (7)	-0.0002 (6)
C14	0.0229 (8)	0.0260 (8)	0.0256 (8)	-0.0031 (6)	-0.0002 (6)	0.0074 (6)
C15	0.0183 (8)	0.0269 (8)	0.0183 (7)	-0.0018 (6)	-0.0008 (6)	-0.0014 (6)

Geometric parameters (\AA , $^\circ$)

F1—C11	1.3419 (17)	C3—H3	0.9500
F2—C12	1.3439 (18)	C4—C5	1.385 (2)
F3—C13	1.3381 (18)	C4—H4	0.9500
F4—C14	1.3393 (17)	C5—C6	1.390 (2)
F5—C15	1.3418 (17)	C5—H5	0.9500
O1—C9	1.2313 (17)	C6—H6	0.9500
N8—C9	1.3287 (19)	C7—H7A	0.9900
N8—C7	1.4596 (19)	C7—H7B	0.9900
N8—H8	0.876 (14)	C9—C10	1.507 (2)
C1—C6	1.388 (2)	C10—C15	1.384 (2)
C1—C2	1.394 (2)	C10—C11	1.388 (2)
C1—C7	1.512 (2)	C11—C12	1.377 (2)
C2—C3	1.383 (2)	C12—C13	1.378 (2)
C2—H2	0.9500	C13—C14	1.374 (2)

C3—C4	1.385 (2)	C14—C15	1.380 (2)
C9—N8—C7	121.84 (13)	N8—C7—H7B	108.8
C9—N8—H8	118.6 (11)	C1—C7—H7B	108.8
C7—N8—H8	119.4 (11)	H7A—C7—H7B	107.7
C6—C1—C2	118.69 (14)	O1—C9—N8	124.59 (14)
C6—C1—C7	122.99 (13)	O1—C9—C10	119.63 (13)
C2—C1—C7	118.28 (13)	N8—C9—C10	115.78 (12)
C3—C2—C1	120.92 (14)	C15—C10—C11	117.21 (14)
C3—C2—H2	119.5	C15—C10—C9	122.86 (13)
C1—C2—H2	119.5	C11—C10—C9	119.79 (13)
C2—C3—C4	120.05 (14)	F1—C11—C12	118.15 (13)
C2—C3—H3	120.0	F1—C11—C10	120.00 (14)
C4—C3—H3	120.0	C12—C11—C10	121.84 (14)
C3—C4—C5	119.56 (14)	F2—C12—C11	120.61 (14)
C3—C4—H4	120.2	F2—C12—C13	119.90 (14)
C5—C4—H4	120.2	C11—C12—C13	119.49 (14)
C4—C5—C6	120.37 (14)	F3—C13—C14	120.15 (14)
C4—C5—H5	119.8	F3—C13—C12	119.81 (14)
C6—C5—H5	119.8	C14—C13—C12	120.04 (14)
C1—C6—C5	120.40 (14)	F4—C14—C13	119.99 (14)
C1—C6—H6	119.8	F4—C14—C15	120.27 (14)
C5—C6—H6	119.8	C13—C14—C15	119.73 (14)
N8—C7—C1	113.87 (12)	F5—C15—C14	118.18 (13)
N8—C7—H7A	108.8	F5—C15—C10	120.08 (13)
C1—C7—H7A	108.8	C14—C15—C10	121.69 (14)
C6—C1—C2—C3	-0.8 (2)	F1—C11—C12—F2	-1.4 (2)
C7—C1—C2—C3	177.02 (14)	C10—C11—C12—F2	179.65 (13)
C1—C2—C3—C4	0.7 (2)	F1—C11—C12—C13	178.39 (13)
C2—C3—C4—C5	0.0 (2)	C10—C11—C12—C13	-0.6 (2)
C3—C4—C5—C6	-0.5 (2)	F2—C12—C13—F3	-0.3 (2)
C2—C1—C6—C5	0.3 (2)	C11—C12—C13—F3	179.87 (14)
C7—C1—C6—C5	-177.44 (14)	F2—C12—C13—C14	-179.91 (14)
C4—C5—C6—C1	0.4 (2)	C11—C12—C13—C14	0.3 (2)
C9—N8—C7—C1	135.84 (15)	F3—C13—C14—F4	-0.3 (2)
C6—C1—C7—N8	-19.9 (2)	C12—C13—C14—F4	179.26 (13)
C2—C1—C7—N8	162.42 (14)	F3—C13—C14—C15	-179.20 (14)
C7—N8—C9—O1	-0.4 (2)	C12—C13—C14—C15	0.4 (2)
C7—N8—C9—C10	178.57 (13)	F4—C14—C15—F5	-2.2 (2)
O1—C9—C10—C15	-121.13 (16)	C13—C14—C15—F5	176.69 (13)
N8—C9—C10—C15	59.85 (19)	F4—C14—C15—C10	-179.71 (13)
O1—C9—C10—C11	54.5 (2)	C13—C14—C15—C10	-0.8 (2)
N8—C9—C10—C11	-124.55 (15)	C11—C10—C15—F5	-176.90 (12)
C15—C10—C11—F1	-178.80 (13)	C9—C10—C15—F5	-1.2 (2)
C9—C10—C11—F1	5.4 (2)	C11—C10—C15—C14	0.5 (2)
C15—C10—C11—C12	0.1 (2)	C9—C10—C15—C14	176.25 (14)
C9—C10—C11—C12	-175.70 (14)		

supplementary materials

Hydrogen-bond geometry (Å, °)

<i>D</i> —H··· <i>A</i>	<i>D</i> —H	H··· <i>A</i>	<i>D</i> ··· <i>A</i>	<i>D</i> —H··· <i>A</i>
N8—H8···O1 ⁱ	0.88 (1)	2.01 (1)	2.875 (2)	171 (2)
C5—H5···O1 ⁱⁱ	0.95	2.37	3.276 (2)	158

Symmetry codes: (i) $x+1/2, -y+1/2, z+1/2$; (ii) $x, y, z+1$.

Fig. 1




Fig. 2




Fig. 3

