

[2-Hydroxy-N'-(4-oxo-4-phenylbutan-2-ylidene)benzohydrazidato(2-)pyridine-copper(II)]

Shu-Ping Zhang, Ying Wei and Si-Chang Shao*

Department of Chemistry, Fuyang Normal College, Fuyang, Anhui 236041, People's Republic of China

Correspondence e-mail: shaosic@fync.edu.cn


Received 28 October 2010; accepted 17 November 2010

Key indicators: single-crystal X-ray study; $T = 298\text{ K}$; mean $\sigma(\text{C}-\text{C}) = 0.007\text{ \AA}$; R factor = 0.057; wR factor = 0.131; data-to-parameter ratio = 16.0.

The mononuclear title complex, $[\text{Cu}(\text{C}_{17}\text{H}_{14}\text{N}_2\text{O}_3)(\text{C}_5\text{H}_5\text{N})]$, was synthesized by the reaction of $\text{CuCl}_2 \cdot 2\text{H}_2\text{O}$ with *N*-(4-oxo-4-phenylbutan-2-ylidene)benzohydrazide (H_2L). The central Cu^{II} atom exhibits a distorted square-planar coordination geometry, defined by two O atoms, one N atom from the ligand and one pyridine N atom with $\text{Cu}-\text{N}$ distances of 1.874 (4) and 1.963 (4) \AA , while the $\text{Cu}-\text{O}$ distances are 1.857 (3) and 1.890 (3) \AA . An intramolecular $\text{O}-\text{H}\cdots\text{N}$ interaction occurs.

Related literature

For the biological properties of Schiff base–metal complexes, see: Cozzi (2004). For metallobiomolecules, see: Singh *et al.* (2007). For metal ions bonded to biologically active compounds, see: Canpolat & Kaya (2004); Yildiz *et al.* (2004). For a related structure, see: Shen *et al.* (1997).



Experimental

Crystal data

$[\text{Cu}(\text{C}_{17}\text{H}_{14}\text{N}_2\text{O}_3)(\text{C}_5\text{H}_5\text{N})]$	$V = 3705.6 (7)\text{ \AA}^3$
$M_r = 436.94$	$Z = 8$
Orthorhombic, $C222_1$	Mo $K\alpha$ radiation
$a = 7.7096 (8)\text{ \AA}$	$\mu = 1.21\text{ mm}^{-1}$
$b = 22.906 (2)\text{ \AA}$	$T = 298\text{ K}$
$c = 20.983 (2)\text{ \AA}$	$0.28 \times 0.20 \times 0.20\text{ mm}$

Data collection

Bruker SMART APEXII CCD diffractometer	13526 measured reflections
Absorption correction: multi-scan (<i>SADABS</i> ; Sheldrick, 1996)	4034 independent reflections
$R_{\text{int}} = 0.050$	3340 reflections with $I > 2\sigma(I)$
$T_{\min} = 0.728$, $T_{\max} = 0.794$	

Refinement

$R[F^2 > 2\sigma(F^2)] = 0.057$	$\Delta\rho_{\text{max}} = 0.45\text{ e \AA}^{-3}$
$wR(F^2) = 0.131$	$\Delta\rho_{\text{min}} = -0.56\text{ e \AA}^{-3}$
$S = 1.08$	Absolute structure: Flack (1983),
4034 reflections	1761 Friedel pairs
252 parameters	Flack parameter: 0.08 (3)
	H-atom parameters constrained

Table 1
Hydrogen-bond geometry (\AA , $^\circ$).

$D-\text{H}\cdots A$	$D-\text{H}$	$\text{H}\cdots A$	$D\cdots A$	$D-\text{H}\cdots A$
O1—H1 \cdots N1	0.82	1.78	2.500 (5)	146

Data collection: *APEX2* (Bruker, 2003); cell refinement: *SAINT* (Bruker, 2003); data reduction: *SAINT*; program(s) used to solve structure: *SHELXS97* (Sheldrick, 2008); program(s) used to refine structure: *SHELXL97* (Sheldrick, 2008); molecular graphics: *SHELXTL* (Sheldrick, 2008); software used to prepare material for publication: *SHELXTL*.

This work was supported by the Natural Science Foundation of Anhui Provincial Education Commission (No. KJ2009A047Z).

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: BX2325).

References

- Bruker (2003). *APEX2* and *SAINT*. Bruker AXS Inc., Madison, Wisconsin, USA.
- Canpolat, E. & Kaya, M. (2004). *J. Coord. Chem.* **57**, 1217–1223.
- Cozzi, P. G. (2004). *Chem. Soc. Rev.* **33**, 410–421.
- Flack, H. D. (1983). *Acta Cryst. A39*, 876–881.
- Sheldrick, G. M. (1996). *SADABS*. University of Gottingen, Germany.
- Sheldrick, G. M. (2008). *Acta Cryst. A64*, 112–122.
- Shen, X., Wu, D., Huang, X., Liu, Q., Huang, Z. & Kang, B. (1997). *Polyhedron*, **16**, 1477–1482.
- Singh, K., Barwa, M. S. & Tyagi, P. (2007). *Eur. J. Med. Chem.* **42**, 394–402.
- Yildiz, M., Dulger, B., Koyuncu, S. Y. & Yapici, B. M. (2004). *J. Indian Chem. Soc.* **81**, 7–12.

supplementary materials

Acta Cryst. (2010). E66, m1635 [doi:10.1107/S1600536810047719]

[2-Hydroxy-*N'*-(4-oxo-4-phenylbutan-2-ylidene)benzohydrazidato(2-)pyridinecopper(II)]

S.-P. Zhang, Y. Wei and S.-C. Shao

Comment

Schiff base metal complexes have been widely studied because they have industrial, antifungal, antibacterial, anticancer and herbicidal applications (Cozzi, 2004). It is well known that N atoms play a key role in the coordination of metals at the active sites of numerous metallobiomolecules (Singh, *et al.*, 2007). They serve as models for biological important species and find applications in biomimetic catalytic reactions. Chelating ligands containing N and O donor atoms show broad biological activity and are of special interest because of the variety of ways in which they are bonded to metal ions. It is known that the existence of metal ions bonded to biologically active compounds may enhance their activities (Canpolat, *et al.*, 2004; Yildiz, *et al.*, 2004). Therefore, it is an important study to design and synthesis of new multidentate ligands containing N and O atoms and apply to synthesize complexes.

The asymmetric unit is composed of one mononuclear complex, (Fig.1). The central Cu^{II} atom exhibits a distorted square-planar coordination geometry, defined by two O atoms, one N atom from the ligand molecule and one N atom of the pyridine molecule with Cu—N distances of 1.874 (4) and 1.963 (4) Å while Cu—O distances are 1.857 (3) and 1.890 (3) Å respectively. The Cu—N and Cu—O distances are comparable to those found in other crystallographically characterized Cu^{II} complex (Shen, *et al.* 1997). The crystal structure of the title compound is stabilized by one intramolecular O—H···N interactions with average H···N distances 1.78 Å and O—H···N angle 146.3°.

Experimental

All reagents and solvents were used as obtained commercially without further purification. CuCl₂.2H₂O (0.170 mg, 0.1 mmol) was dissolved in 6 ml deionized water, giving a transparent solution, and 1 mL pyridine solution dissolved with *L* (28.2 mg, 0.1 mmol) was dropwised for 0.5 h. After stirring for 8 h, the solution was filtered. Black single crystals of the title compound were obtained from the filtrate after 3 weeks. Analysis calculated (%): C, 60.47; H, 4.38; N, 9.62%; Found: C, 60.15; H, 4.59; N, 9.49%.

Refinement

H atoms bonded to C atoms were placed geometrically and treated as riding, with C—H distances 0.93–0.96 Å and *U*_{iso}(H) = 1.2*U*_{eq}(C) for the CH while *U*_{iso}(H) = 1.5*U*_{eq}(C) for the CH₃ groups. The hydroxyl H atoms were located from difference maps and refined with the O—H distances restrained to 0.82 Å and *U*_{iso}(H) = 1.5*U*_{eq}(O).

supplementary materials

Figures




Fig. 1. The molecular structure of the title compound, showing 30% probability displacement ellipsoids.

[2-Hydroxy-*N'*-(4-oxo-4-phenylbutan-2-ylidene)benzohydrazidato(2-)pyridinecopper(II)]

Crystal data

$[\text{Cu}(\text{C}_{17}\text{H}_{14}\text{N}_2\text{O}_3)(\text{C}_5\text{H}_5\text{N})]$	$F(000) = 1800$
$M_r = 436.94$	$D_x = 1.566 \text{ Mg m}^{-3}$
Orthorhombic, $C222_1$	Mo $K\alpha$ radiation, $\lambda = 0.71073 \text{ \AA}$
Hall symbol: C 2c 2	Cell parameters from 3542 reflections
$a = 7.7096 (8) \text{ \AA}$	$\theta = 2.1\text{--}23.1^\circ$
$b = 22.906 (2) \text{ \AA}$	$\mu = 1.21 \text{ mm}^{-1}$
$c = 20.983 (2) \text{ \AA}$	$T = 298 \text{ K}$
$V = 3705.6 (7) \text{ \AA}^3$	Block, dark green
$Z = 8$	$0.28 \times 0.20 \times 0.20 \text{ mm}$

Data collection

Bruker SMART APEXII CCD diffractometer	4034 independent reflections
Radiation source: fine-focus sealed tube graphite	3340 reflections with $I > 2\sigma(I)$
phi and ω scans	$R_{\text{int}} = 0.050$
Absorption correction: multi-scan (<i>SADABS</i> ; Sheldrick, 1996)	$\theta_{\text{max}} = 27.0^\circ, \theta_{\text{min}} = 1.8^\circ$
$T_{\text{min}} = 0.728, T_{\text{max}} = 0.794$	$h = -9 \rightarrow 9$
13526 measured reflections	$k = -29 \rightarrow 27$
	$l = -26 \rightarrow 26$

Refinement

Refinement on F^2	Secondary atom site location: difference Fourier map
Least-squares matrix: full	Hydrogen site location: inferred from neighbouring sites
$R[F^2 > 2\sigma(F^2)] = 0.057$	H-atom parameters constrained
$wR(F^2) = 0.131$	$w = 1/[\sigma^2(F_o^2) + (0.0614P)^2 + 1.8583P]$
$S = 1.08$	where $P = (F_o^2 + 2F_c^2)/3$
4034 reflections	$(\Delta/\sigma)_{\text{max}} = 0.003$
252 parameters	$\Delta\rho_{\text{max}} = 0.45 \text{ e \AA}^{-3}$
0 restraints	$\Delta\rho_{\text{min}} = -0.56 \text{ e \AA}^{-3}$
	Absolute structure: Flack (1983), 1761 Friedel pairs

Primary atom site location: structure-invariant direct Flack parameter: 0.08 (3)
methods

Special details

Geometry. All e.s.d.'s (except the e.s.d. in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell e.s.d.'s are taken into account individually in the estimation of e.s.d.'s in distances, angles and torsion angles; correlations between e.s.d.'s in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell e.s.d.'s is used for estimating e.s.d.'s involving l.s. planes.

Refinement. Refinement of F^2 against ALL reflections. The weighted R -factor wR and goodness of fit S are based on F^2 , conventional R -factors R are based on F , with F set to zero for negative F^2 . The threshold expression of $F^2 > \sigma(F^2)$ is used only for calculating R -factors(gt) etc. and is not relevant to the choice of reflections for refinement. R -factors based on F^2 are statistically about twice as large as those based on F , and R -factors based on ALL data will be even larger.

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (\AA^2)

	x	y	z	$U_{\text{iso}}^*/U_{\text{eq}}$
Cu1	0.26164 (7)	0.56424 (2)	0.75389 (2)	0.03955 (17)
C1	0.4395 (6)	0.4324 (2)	0.8613 (2)	0.0396 (11)
C2	0.5248 (7)	0.4551 (2)	0.9121 (2)	0.0496 (13)
H2	0.5322	0.4955	0.9165	0.059*
C3	0.5994 (7)	0.4205 (3)	0.9567 (2)	0.0558 (14)
H3	0.6569	0.4370	0.9913	0.067*
C4	0.5895 (8)	0.3611 (3)	0.9504 (3)	0.0653 (17)
H4	0.6401	0.3369	0.9807	0.078*
C5	0.5071 (9)	0.3384 (2)	0.9008 (2)	0.0534 (12)
H5	0.5021	0.2980	0.8964	0.064*
C6	0.4294 (7)	0.3728 (2)	0.8558 (2)	0.0440 (12)
C7	0.3627 (6)	0.4715 (2)	0.8143 (2)	0.0386 (11)
C8	0.0179 (5)	0.60424 (13)	0.58645 (12)	0.0411 (11)
C9	0.0827 (5)	0.66043 (14)	0.57885 (14)	0.0524 (13)
H9A	0.1631	0.6751	0.6078	0.063*
C10	0.0275 (5)	0.69465 (12)	0.52805 (16)	0.0589 (15)
H10A	0.0709	0.7322	0.5230	0.071*
C11	-0.0925 (5)	0.67268 (15)	0.48484 (14)	0.0617 (16)
H11A	-0.1295	0.6956	0.4509	0.074*
C12	-0.1574 (4)	0.61649 (16)	0.49245 (15)	0.0614 (15)
H12A	-0.2377	0.6018	0.4635	0.074*
C13	-0.1022 (5)	0.58227 (12)	0.54325 (16)	0.0589 (15)
H13A	-0.1456	0.5447	0.5483	0.071*
C14	0.0822 (6)	0.5682 (2)	0.6383 (2)	0.0423 (11)
C15	0.0672 (6)	0.5104 (2)	0.6359 (2)	0.0435 (12)
H15	0.0074	0.4954	0.6011	0.052*
C16	0.1308 (6)	0.4698 (2)	0.6795 (2)	0.0391 (11)
C17	0.1033 (7)	0.4076 (2)	0.6673 (2)	0.0476 (12)
H17A	0.0384	0.3908	0.7017	0.071*
H17B	0.2134	0.3883	0.6638	0.071*
H17C	0.0399	0.4029	0.6282	0.071*

supplementary materials

C18	0.3924 (7)	0.6464 (2)	0.8448 (2)	0.0532 (14)
H18	0.4366	0.6128	0.8636	0.064*
C19	0.4202 (9)	0.6974 (3)	0.8736 (3)	0.0635 (17)
H19	0.4847	0.6989	0.9110	0.076*
C20	0.3560 (11)	0.7459 (3)	0.8488 (3)	0.080 (2)
H20	0.3752	0.7818	0.8682	0.096*
C21	0.2630 (12)	0.7419 (2)	0.7953 (3)	0.085 (2)
H21	0.2139	0.7750	0.7771	0.102*
C22	0.2413 (9)	0.6887 (2)	0.7680 (2)	0.0658 (17)
H22	0.1769	0.6862	0.7307	0.079*
N1	0.2780 (5)	0.44657 (14)	0.76881 (16)	0.0376 (8)
N2	0.2134 (4)	0.48748 (16)	0.72866 (15)	0.0353 (8)
N3	0.3066 (5)	0.64115 (17)	0.79176 (17)	0.0400 (9)
O1	0.3475 (6)	0.34649 (15)	0.80904 (17)	0.0602 (10)
H1	0.3070	0.3710	0.7848	0.090*
O2	0.3808 (5)	0.52533 (13)	0.82017 (15)	0.0428 (8)
O3	0.1558 (5)	0.59686 (14)	0.68281 (15)	0.0499 (9)

Atomic displacement parameters (\AA^2)

	U^{11}	U^{22}	U^{33}	U^{12}	U^{13}	U^{23}
Cu1	0.0412 (3)	0.0412 (3)	0.0363 (3)	0.0042 (3)	-0.0104 (3)	-0.0039 (2)
C1	0.034 (2)	0.045 (3)	0.039 (2)	0.004 (2)	0.0083 (18)	0.008 (2)
C2	0.058 (4)	0.050 (3)	0.041 (3)	0.005 (3)	0.000 (2)	-0.002 (2)
C3	0.059 (4)	0.066 (4)	0.043 (3)	0.008 (3)	-0.006 (2)	0.007 (2)
C4	0.056 (4)	0.085 (5)	0.055 (3)	0.014 (4)	0.005 (3)	0.027 (3)
C5	0.058 (3)	0.050 (3)	0.052 (3)	0.007 (3)	0.010 (3)	0.011 (2)
C6	0.050 (3)	0.041 (3)	0.041 (3)	0.006 (2)	0.014 (2)	-0.003 (2)
C7	0.030 (2)	0.050 (3)	0.036 (2)	0.002 (2)	-0.004 (2)	-0.004 (2)
C8	0.035 (3)	0.058 (3)	0.031 (2)	0.005 (3)	-0.005 (2)	-0.008 (2)
C9	0.060 (3)	0.056 (3)	0.041 (3)	0.003 (3)	-0.004 (2)	-0.005 (2)
C10	0.064 (4)	0.061 (3)	0.052 (3)	0.005 (3)	-0.001 (3)	0.003 (3)
C11	0.064 (4)	0.082 (5)	0.039 (3)	0.018 (3)	-0.002 (3)	-0.002 (3)
C12	0.058 (3)	0.075 (4)	0.051 (3)	0.007 (3)	-0.018 (3)	0.007 (3)
C13	0.049 (3)	0.074 (4)	0.054 (3)	0.006 (3)	-0.021 (3)	0.003 (3)
C14	0.034 (2)	0.056 (3)	0.036 (2)	0.012 (3)	-0.0044 (19)	-0.003 (2)
C15	0.040 (3)	0.057 (3)	0.034 (2)	-0.002 (2)	-0.008 (2)	-0.013 (2)
C16	0.032 (2)	0.042 (3)	0.043 (2)	-0.002 (2)	0.007 (2)	-0.009 (2)
C17	0.050 (3)	0.047 (3)	0.046 (3)	-0.004 (3)	-0.011 (2)	-0.008 (2)
C18	0.061 (4)	0.051 (3)	0.048 (3)	0.003 (3)	-0.019 (3)	-0.007 (2)
C19	0.084 (4)	0.049 (3)	0.057 (3)	0.007 (3)	-0.026 (3)	-0.006 (3)
C20	0.124 (7)	0.044 (3)	0.072 (4)	0.005 (4)	-0.018 (4)	-0.013 (3)
C21	0.133 (7)	0.045 (3)	0.078 (4)	0.028 (4)	-0.036 (5)	-0.012 (3)
C22	0.097 (5)	0.053 (3)	0.048 (3)	0.014 (4)	-0.028 (4)	0.002 (2)
N1	0.038 (2)	0.0335 (19)	0.0408 (17)	0.0034 (16)	0.0008 (17)	0.0041 (14)
N2	0.0283 (19)	0.045 (2)	0.0327 (16)	0.0030 (16)	0.0005 (15)	-0.0042 (15)
N3	0.043 (2)	0.042 (2)	0.0355 (18)	0.0042 (18)	-0.0054 (16)	0.0018 (17)
O1	0.081 (3)	0.040 (2)	0.059 (2)	0.003 (2)	-0.015 (2)	-0.0024 (18)

O2	0.051 (2)	0.0356 (19)	0.0417 (17)	0.0018 (15)	-0.0138 (16)	-0.0037 (14)
O3	0.064 (2)	0.044 (2)	0.0417 (17)	0.0059 (18)	-0.0178 (16)	-0.0048 (16)

Geometric parameters (\AA , $^\circ$)

Cu1—O3	1.857 (3)	C11—H11A	0.9300
Cu1—N2	1.874 (4)	C12—C13	1.3900
Cu1—O2	1.890 (3)	C12—H12A	0.9300
Cu1—N3	1.963 (4)	C13—H13A	0.9300
C1—C2	1.357 (7)	C14—O3	1.276 (5)
C1—C6	1.373 (7)	C14—C15	1.328 (7)
C1—C7	1.457 (7)	C15—C16	1.394 (7)
C2—C3	1.355 (7)	C15—H15	0.9300
C2—H2	0.9300	C16—N2	1.278 (6)
C3—C4	1.369 (9)	C16—C17	1.462 (7)
C3—H3	0.9300	C17—H17A	0.9600
C4—C5	1.327 (8)	C17—H17B	0.9600
C4—H4	0.9300	C17—H17C	0.9600
C5—C6	1.367 (7)	C18—N3	1.300 (6)
C5—H5	0.9300	C18—C19	1.332 (8)
C6—O1	1.314 (6)	C18—H18	0.9300
C7—O2	1.247 (5)	C19—C20	1.323 (8)
C7—N1	1.291 (6)	C19—H19	0.9300
C8—C9	1.3900	C20—C21	1.335 (9)
C8—C13	1.3900	C20—H20	0.9300
C8—C14	1.453 (5)	C21—C22	1.356 (7)
C9—C10	1.3900	C21—H21	0.9300
C9—H9A	0.9300	C22—N3	1.300 (6)
C10—C11	1.3900	C22—H22	0.9300
C10—H10A	0.9300	N1—N2	1.355 (5)
C11—C12	1.3900	O1—H1	0.8200
O3—Cu1—N2	93.64 (15)	C12—C13—H13A	120.0
O3—Cu1—O2	173.85 (15)	C8—C13—H13A	120.0
N2—Cu1—O2	82.05 (14)	O3—C14—C15	125.4 (4)
O3—Cu1—N3	92.39 (15)	O3—C14—C8	114.0 (4)
N2—Cu1—N3	172.50 (15)	C15—C14—C8	120.6 (4)
O2—Cu1—N3	92.27 (14)	C14—C15—C16	127.5 (4)
C2—C1—C6	118.3 (5)	C14—C15—H15	116.2
C2—C1—C7	119.5 (5)	C16—C15—H15	116.2
C6—C1—C7	122.1 (5)	N2—C16—C15	119.6 (4)
C3—C2—C1	121.6 (5)	N2—C16—C17	121.5 (4)
C3—C2—H2	119.2	C15—C16—C17	118.9 (4)
C1—C2—H2	119.2	C16—C17—H17A	109.5
C2—C3—C4	119.4 (5)	C16—C17—H17B	109.5
C2—C3—H3	120.3	H17A—C17—H17B	109.5
C4—C3—H3	120.3	C16—C17—H17C	109.5
C5—C4—C3	119.5 (5)	H17A—C17—H17C	109.5
C5—C4—H4	120.2	H17B—C17—H17C	109.5
C3—C4—H4	120.2	N3—C18—C19	123.4 (5)

supplementary materials

C4—C5—C6	121.7 (5)	N3—C18—H18	118.3
C4—C5—H5	119.2	C19—C18—H18	118.3
C6—C5—H5	119.2	C20—C19—C18	119.9 (6)
O1—C6—C5	117.5 (5)	C20—C19—H19	120.0
O1—C6—C1	123.1 (5)	C18—C19—H19	120.0
C5—C6—C1	119.4 (5)	C19—C20—C21	118.3 (6)
O2—C7—N1	124.5 (4)	C19—C20—H20	120.9
O2—C7—C1	119.7 (4)	C21—C20—H20	120.9
N1—C7—C1	115.7 (4)	C20—C21—C22	118.9 (6)
C9—C8—C13	120.0	C20—C21—H21	120.6
C9—C8—C14	119.3 (3)	C22—C21—H21	120.6
C13—C8—C14	120.6 (3)	N3—C22—C21	122.9 (5)
C10—C9—C8	120.0	N3—C22—H22	118.5
C10—C9—H9A	120.0	C21—C22—H22	118.5
C8—C9—H9A	120.0	C7—N1—N2	109.9 (4)
C9—C10—C11	120.0	C16—N2—N1	117.8 (4)
C9—C10—H10A	120.0	C16—N2—Cu1	128.6 (3)
C11—C10—H10A	120.0	N1—N2—Cu1	113.6 (3)
C12—C11—C10	120.0	C22—N3—C18	116.6 (4)
C12—C11—H11A	120.0	C22—N3—Cu1	122.0 (3)
C10—C11—H11A	120.0	C18—N3—Cu1	121.2 (4)
C11—C12—C13	120.0	C6—O1—H1	109.5
C11—C12—H12A	120.0	C7—O2—Cu1	109.9 (3)
C13—C12—H12A	120.0	C14—O3—Cu1	125.2 (3)
C12—C13—C8	120.0		
C6—C1—C2—C3	0.2 (8)	N3—C18—C19—C20	1.4 (10)
C7—C1—C2—C3	−179.5 (5)	C18—C19—C20—C21	0.5 (12)
C1—C2—C3—C4	0.2 (9)	C19—C20—C21—C22	−1.4 (13)
C2—C3—C4—C5	0.1 (9)	C20—C21—C22—N3	0.3 (13)
C3—C4—C5—C6	−0.9 (10)	O2—C7—N1—N2	0.3 (6)
C4—C5—C6—O1	−178.5 (5)	C1—C7—N1—N2	180.0 (3)
C4—C5—C6—C1	1.3 (9)	C15—C16—N2—N1	−177.9 (4)
C2—C1—C6—O1	178.9 (5)	C17—C16—N2—N1	1.1 (6)
C7—C1—C6—O1	−1.5 (8)	C15—C16—N2—Cu1	1.6 (6)
C2—C1—C6—C5	−1.0 (7)	C17—C16—N2—Cu1	−179.4 (4)
C7—C1—C6—C5	178.7 (5)	C7—N1—N2—C16	177.6 (4)
C2—C1—C7—O2	2.3 (7)	C7—N1—N2—Cu1	−1.9 (4)
C6—C1—C7—O2	−177.4 (5)	O3—Cu1—N2—C16	−1.8 (4)
C2—C1—C7—N1	−177.4 (4)	O2—Cu1—N2—C16	−177.4 (4)
C6—C1—C7—N1	3.0 (7)	O3—Cu1—N2—N1	177.7 (3)
C13—C8—C9—C10	0.0	O2—Cu1—N2—N1	2.1 (3)
C14—C8—C9—C10	177.5 (4)	C21—C22—N3—C18	1.5 (10)
C8—C9—C10—C11	0.0	C21—C22—N3—Cu1	176.4 (6)
C9—C10—C11—C12	0.0	C19—C18—N3—C22	−2.4 (8)
C10—C11—C12—C13	0.0	C19—C18—N3—Cu1	−177.3 (5)
C11—C12—C13—C8	0.0	O2—Cu1—N3—C22	−175.0 (5)
C9—C8—C13—C12	0.0	O3—Cu1—N3—C18	−176.3 (4)
C14—C8—C13—C12	−177.4 (4)	O2—Cu1—N3—C18	−0.3 (4)
C9—C8—C14—O3	19.2 (5)	N1—C7—O2—Cu1	1.4 (6)

C13—C8—C14—O3	−163.4 (3)	C1—C7—O2—Cu1	−178.3 (3)
C9—C8—C14—C15	−158.8 (4)	N2—Cu1—O2—C7	−1.8 (3)
C13—C8—C14—C15	18.7 (6)	N3—Cu1—O2—C7	173.3 (4)
O3—C14—C15—C16	−1.8 (9)	C15—C14—O3—Cu1	1.0 (7)
C8—C14—C15—C16	175.9 (4)	C8—C14—O3—Cu1	−176.8 (3)
C14—C15—C16—N2	0.4 (8)	N2—Cu1—O3—C14	0.5 (4)
C14—C15—C16—C17	−178.7 (5)	N3—Cu1—O3—C14	−175.0 (4)

Hydrogen-bond geometry (Å, °)

<i>D</i> —H··· <i>A</i>	<i>D</i> —H	H··· <i>A</i>	<i>D</i> ··· <i>A</i>	<i>D</i> —H··· <i>A</i>
O1—H1···N1	0.82	1.78	2.500 (5)	146

supplementary materials

Fig. 1

