

1-Cyanomethyl-1,4-diazoniabicyclo-[2.2.2]octane tetrabromidocadmamate(II)

Bin Wei

Ordered Matter Science Research Center, Southeast University, Nanjing 211189, People's Republic of China
Correspondence e-mail: seuwei@126.com


Received 12 October 2010; accepted 16 November 2010

Key indicators: single-crystal X-ray study; $T = 293\text{ K}$; mean $\sigma(\text{C}-\text{C}) = 0.007\text{ \AA}$; R factor = 0.033; wR factor = 0.079; data-to-parameter ratio = 24.3.

In the title salt, $(\text{C}_8\text{H}_{15}\text{N}_3)[\text{CdBr}_4]$, four Br atoms coordinate the Cd^{II} atom in a distorted tetrahedral geometry. In the crystal, weak $\text{N}-\text{H}\cdots\text{Br}$ interactions connect the anion to three symmetry-related cations. The crystal structure also displays very weak $\text{C}-\text{H}\cdots\text{Br}$ interactions.

Related literature

For background to 1,4-diazabicyclo[2.2.2]octane derivatives and their properties, see: Basavaiah *et al.* (2003); Chen *et al.* (2010); Wang *et al.* (2005); Xiong *et al.* (2002); Ye *et al.* (2006).



Experimental

Crystal data

$(\text{C}_8\text{H}_{15}\text{N}_3)[\text{CdBr}_4]$
 $M_r = 585.27$
Monoclinic, $P2_1/c$
 $a = 8.610 (3)\text{ \AA}$
 $b = 14.071 (4)\text{ \AA}$
 $c = 12.702 (4)\text{ \AA}$
 $\beta = 94.136 (4)^\circ$

$V = 1534.9 (8)\text{ \AA}^3$
 $Z = 4$
Mo $K\alpha$ radiation
 $\mu = 11.82\text{ mm}^{-1}$
 $T = 293\text{ K}$
 $0.2 \times 0.2 \times 0.2\text{ mm}$

Data collection

Rigaku Mercury CCD diffractometer
Absorption correction: multi-scan (*CrystalClear*; Rigaku, 2005)
 $T_{\min} = 0.470$, $T_{\max} = 1.000$

16630 measured reflections
3518 independent reflections
2861 reflections with $I > 2\sigma(I)$
 $R_{\text{int}} = 0.068$

Refinement

$R[F^2 > 2\sigma(F^2)] = 0.033$
 $wR(F^2) = 0.079$
 $S = 0.76$
3518 reflections

145 parameters
H-atom parameters constrained
 $\Delta\rho_{\max} = 0.61\text{ e \AA}^{-3}$
 $\Delta\rho_{\min} = -0.88\text{ e \AA}^{-3}$

Table 1
Hydrogen-bond geometry (\AA , $^\circ$).

$D-\text{H}\cdots A$	$D-\text{H}$	$\text{H}\cdots A$	$D\cdots A$	$D-\text{H}\cdots A$
N1—H1C \cdots Br1 ⁱ	0.90	2.85	3.466 (4)	127
N1—H1C \cdots Br2 ⁱⁱ	0.90	2.69	3.325 (4)	128
N1—H1C \cdots Br4 ⁱ	0.90	3.11	3.711 (4)	126
C2—H2B \cdots Br3 ⁱⁱⁱ	0.97	2.83	3.765 (5)	162
C4—H4B \cdots Br1	0.97	2.85	3.643 (4)	140
C7—H7A \cdots Br3 ^{iv}	0.97	2.90	3.626 (4)	132
C7—H7B \cdots Br2 ⁱⁱⁱ	0.97	2.78	3.683 (4)	154

Symmetry codes: (i) $-x + 2, -y + 1, -z + 1$; (ii) $-x + 2, y + \frac{1}{2}, -z + \frac{3}{2}$; (iii) $-x + 1, y + \frac{1}{2}, -z + \frac{3}{2}$; (iv) $x, -y + \frac{1}{2}, z + \frac{1}{2}$.

Data collection: *CrystalClear* (Rigaku, 2005); cell refinement: *CrystalClear*; data reduction: *CrystalClear*; program(s) used to solve structure: *SHELXS97* (Sheldrick, 2008); program(s) used to refine structure: *SHELXL97* (Sheldrick, 2008); molecular graphics: *SHELXTL* (Sheldrick, 2008); software used to prepare material for publication: *SHELXL97*.

The author is grateful to the starter fund of Southeast University for the purchase of the diffractometer.

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: BH2319).

References

- Basavaiah, D., Rao, A. J. & Satyanarayana, T. (2003). *Chem. Rev.* **103**, 811–892.
- Chen, L.-Z., Huang, Y., Xiong, R.-G. & Hu, H.-W. (2010). *J. Mol. Struct.* **963**, 16–21.
- Rigaku (2005). *CrystalClear*. Rigaku Corporation, Tokyo, Japan.
- Sheldrick, G. M. (2008). *Acta Cryst. A* **64**, 112–122.
- Wang, X.-S., Song, Y.-M. & Xiong, R.-G. (2005). *Chin. J. Inorg. Chem.* **21**, 1030–1033.
- Xiong, R. G., Xue, X., Zhao, H., You, X.-Z., Abrahams, B. F. & Xue, Z. (2002). *Angew. Chem. Int. Ed.* **41**, 3800–3803.
- Ye, Q., Song, Y.-M., Wang, G.-X., Chen, K., Fu, D.-W., Chan, P. W. H., Zhu, J.-S., Huang, S. D. & Xiong, R.-G. (2006). *J. Am. Chem. Soc.* **128**, 6554–6555.

supplementary materials

Acta Cryst. (2010). E66, m1672 [doi:10.1107/S1600536810047495]

1-Cyanomethyl-1,4-diazoniabicyclo[2.2.2]octane tetrabromidocadmate(II)

B. Wei

Comment

We are studying the dielectric-ferroelectric materials. 1,4-Diazabicyclo[2.2.2]octane (DABCO) has attracted attention in recent years because of its nucleophilicity (Basavaiah *et al.*, 2003; Xiong *et al.*, 2002) and ferroelectric properties of its derivatives (Chen *et al.*, 2010). For a project on the electric properties of DABCO derivatives (Ye *et al.*, 2006), the title compound was prepared. With no dielectric anomaly observed, the title compound should not be a real ferroelectrics or there may be no distinct phase transition occurring within the measured temperature range (Wang *et al.*, 2005).

The asymmetric unit of the title compound is shown in Fig 1. The Cd atoms are coordinated by four Br atoms with very similar distances in the range 2.5764 (10) to 2.6195 (12) Å. The Br—Cd—Br bond angles are between 98.29 (3) and 116.85 (4)°, which show that the coordination polyhedron can be described as an irregular tetrahedron. Cations ($C_8H_{14}N_3$)²⁺ and anions $CdBr_4^{2-}$ are connected *via* weak hydrogen bonds. Weak C—H···Br intramolecular and intermolecular hydrogen bonds also contribute to the stability of the crystal structure, forming one-dimensional chains running along the *a* axis (Fig. 2).

Experimental

1,4-Diaza-bicyclo[2.2.2]octane (DABCO) (10 mmol, 1.14 g) and bromoacetonitrile (20 mmol, 2.4 g) were dissolved in CH₃CN (10 ml) under stirring for 1 h. at room temperature. 1-(Cyanomethyl)-4-aza-1-azonia-bicyclo[2.2.2]octane bromide was obtained by filtering the solid precipitate, then washed with acetonitrile and dried (yield: 90%).

CdBr₂ (10 mmol, 0.271 g) and 4 ml 60% HBr were dissolved in MeOH (20 ml) and 1-(cyanomethyl)-4-aza-1-azonia-bicyclo[2.2.2]octane bromide (20 mmol, 0.464 g) dissolved in 10 ml of methanol was added. The mixture was stirred until the solution was clear. After slow evaporation (5 days) of the solvent, colourless plate crystals of the title compound were obtained in about 56% yield.

Refinement

H atoms bonded to C and N atoms were placed in idealized positions [C—H = 0.97 Å and N—H = 0.90 Å] and allowed to ride on their parent atoms with U_{iso} fixed at 1.2 U_{eq} (Carrier atom).

supplementary materials

Figures




Fig. 1. The structure of the title compound with labeling scheme and displacement ellipsoids drawn at the 30% probability level.




Fig. 2. The crystal structure of the title compound viewed down the c axis. Intermolecular interactions are shown as dashed lines.

1-Cyanomethyl-1,4-diazeniabicyclo[2.2.2]octane tetrabromidocadmate(II)

Crystal data

(C ₈ H ₁₅ N ₃)[CdBr ₄]	$F(000) = 1088$
$M_r = 585.27$	$D_x = 2.533 \text{ Mg m}^{-3}$
Monoclinic, $P2_1/c$	Mo $K\alpha$ radiation, $\lambda = 0.71073 \text{ \AA}$
Hall symbol: -P 2ybc	Cell parameters from 4263 reflections
$a = 8.610 (3) \text{ \AA}$	$\theta = 2.4\text{--}27.5^\circ$
$b = 14.071 (4) \text{ \AA}$	$\mu = 11.82 \text{ mm}^{-1}$
$c = 12.702 (4) \text{ \AA}$	$T = 293 \text{ K}$
$\beta = 94.136 (4)^\circ$	Prism, colourless
$V = 1534.9 (8) \text{ \AA}^3$	$0.2 \times 0.2 \times 0.2 \text{ mm}$
$Z = 4$	

Data collection

Rigaku Mercury CCD diffractometer	3518 independent reflections
Radiation source: fine-focus sealed tube graphite	2861 reflections with $I > 2\sigma(I)$
Detector resolution: 28.5714 pixels mm^{-1}	$R_{\text{int}} = 0.068$
ω scans	$\theta_{\text{max}} = 27.5^\circ, \theta_{\text{min}} = 3.1^\circ$
Absorption correction: multi-scan (<i>CrystalClear</i> ; Rigaku, 2005)	$h = -11 \rightarrow 11$
$T_{\text{min}} = 0.470, T_{\text{max}} = 1.000$	$k = -18 \rightarrow 18$
16630 measured reflections	$l = -16 \rightarrow 16$

Refinement

Refinement on F^2	Primary atom site location: structure-invariant direct methods
Least-squares matrix: full	Secondary atom site location: difference Fourier map
$R[F^2 > 2\sigma(F^2)] = 0.033$	Hydrogen site location: inferred from neighbouring sites
$wR(F^2) = 0.079$	H-atom parameters constrained
$S = 0.76$	$w = 1/[\sigma^2(F_o^2) + (0.0502P)^2 + 0.2386P]$ where $P = (F_o^2 + 2F_c^2)/3$
3518 reflections	$(\Delta/\sigma)_{\max} = 0.018$
145 parameters	$\Delta\rho_{\max} = 0.61 \text{ e \AA}^{-3}$
0 restraints	$\Delta\rho_{\min} = -0.88 \text{ e \AA}^{-3}$
0 constraints	

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (\AA^2)

	<i>x</i>	<i>y</i>	<i>z</i>	$U_{\text{iso}}^*/U_{\text{eq}}$
Cd1	0.78773 (4)	0.22670 (2)	0.50861 (3)	0.02924 (11)
Br1	0.82000 (6)	0.40556 (3)	0.46315 (4)	0.03047 (13)
Br2	0.79028 (6)	0.24284 (3)	0.71414 (4)	0.03420 (13)
Br3	0.52068 (6)	0.15357 (3)	0.45098 (4)	0.03464 (13)
Br4	1.03019 (6)	0.13565 (3)	0.45657 (4)	0.03694 (14)
C8	0.4254 (6)	0.4480 (3)	0.6975 (4)	0.0314 (11)
N2	0.6179 (4)	0.5753 (2)	0.7314 (3)	0.0205 (7)
C5	0.9032 (5)	0.5513 (3)	0.7405 (4)	0.0328 (11)
H5A	0.9784	0.5564	0.8008	0.039*
H5B	0.9444	0.5081	0.6898	0.039*
N1	0.8754 (4)	0.6469 (2)	0.6913 (3)	0.0242 (8)
H1C	0.9610	0.6750	0.6691	0.029*
C6	0.7506 (6)	0.5138 (3)	0.7750 (4)	0.0378 (12)
H6A	0.7350	0.4490	0.7501	0.045*
H6B	0.7532	0.5132	0.8515	0.045*
C4	0.6267 (5)	0.5849 (3)	0.6140 (3)	0.0294 (10)
H4A	0.5365	0.6195	0.5842	0.035*
H4B	0.6263	0.5224	0.5819	0.035*
C3	0.7744 (5)	0.6376 (3)	0.5904 (3)	0.0283 (10)
H3A	0.8293	0.6027	0.5387	0.034*
H3B	0.7486	0.7000	0.5620	0.034*
C2	0.6364 (6)	0.6733 (3)	0.7808 (4)	0.0316 (11)
H2A	0.5586	0.7160	0.7484	0.038*
H2B	0.6215	0.6696	0.8557	0.038*
C7	0.4643 (5)	0.5354 (3)	0.7579 (3)	0.0279 (10)
H7A	0.4677	0.5215	0.8328	0.034*
H7B	0.3835	0.5825	0.7424	0.034*
N3	0.3933 (6)	0.3830 (3)	0.6497 (4)	0.0459 (12)
C1	0.7981 (5)	0.7107 (3)	0.7646 (4)	0.0300 (10)

supplementary materials

H1A	0.7909	0.7744	0.7354	0.036*
H1B	0.8590	0.7135	0.8318	0.036*

Atomic displacement parameters (\AA^2)

	U^{11}	U^{22}	U^{33}	U^{12}	U^{13}	U^{23}
Cd1	0.02631 (19)	0.03071 (19)	0.0309 (2)	-0.00124 (14)	0.00314 (15)	0.00137 (14)
Br1	0.0324 (3)	0.0264 (2)	0.0335 (3)	0.00131 (19)	0.0088 (2)	-0.00169 (18)
Br2	0.0285 (3)	0.0477 (3)	0.0266 (3)	0.0031 (2)	0.0034 (2)	0.0044 (2)
Br3	0.0287 (3)	0.0421 (3)	0.0334 (3)	-0.0064 (2)	0.0042 (2)	-0.0070 (2)
Br4	0.0334 (3)	0.0351 (3)	0.0433 (3)	0.0054 (2)	0.0086 (2)	0.0016 (2)
C8	0.029 (3)	0.026 (2)	0.039 (3)	-0.008 (2)	0.003 (2)	0.010 (2)
N2	0.0180 (18)	0.0228 (18)	0.0213 (18)	0.0003 (14)	0.0053 (15)	0.0002 (14)
C5	0.023 (2)	0.032 (3)	0.043 (3)	0.007 (2)	0.003 (2)	0.002 (2)
N1	0.0148 (18)	0.031 (2)	0.029 (2)	-0.0028 (15)	0.0094 (15)	-0.0017 (15)
C6	0.027 (3)	0.034 (3)	0.052 (3)	0.008 (2)	-0.003 (2)	0.015 (2)
C4	0.027 (3)	0.045 (3)	0.016 (2)	-0.004 (2)	0.0037 (19)	-0.0024 (18)
C3	0.025 (2)	0.035 (3)	0.026 (2)	-0.002 (2)	0.007 (2)	0.0022 (18)
C2	0.033 (3)	0.031 (2)	0.033 (3)	-0.003 (2)	0.015 (2)	-0.013 (2)
C7	0.020 (2)	0.033 (2)	0.032 (3)	-0.0025 (19)	0.0067 (19)	0.0061 (19)
N3	0.051 (3)	0.032 (2)	0.053 (3)	-0.012 (2)	-0.004 (2)	0.011 (2)
C1	0.028 (2)	0.028 (2)	0.035 (3)	-0.003 (2)	0.008 (2)	-0.0084 (19)

Geometric parameters (\AA , $^\circ$)

Cd1—Br3	2.5766 (8)	N1—H1C	0.8997
Cd1—Br4	2.5760 (8)	C6—H6A	0.9700
Cd1—Br1	2.6015 (9)	C6—H6B	0.9700
Cd1—Br2	2.6191 (10)	C4—C3	1.521 (6)
C8—N3	1.122 (6)	C4—H4A	0.9700
C8—C7	1.475 (6)	C4—H4B	0.9700
N2—C7	1.497 (5)	C3—H3A	0.9700
N2—C6	1.506 (5)	C3—H3B	0.9700
N2—C4	1.505 (5)	C2—C1	1.516 (6)
N2—C2	1.519 (5)	C2—H2A	0.9700
C5—N1	1.495 (5)	C2—H2B	0.9700
C5—C6	1.510 (7)	C7—H7A	0.9700
C5—H5A	0.9700	C7—H7B	0.9700
C5—H5B	0.9700	C1—H1A	0.9700
N1—C1	1.485 (5)	C1—H1B	0.9700
N1—C3	1.501 (5)		
Br3—Cd1—Br4	116.84 (3)	N2—C4—C3	110.0 (3)
Br3—Cd1—Br1	115.51 (2)	N2—C4—H4A	109.7
Br4—Cd1—Br1	108.83 (2)	C3—C4—H4A	109.7
Br3—Cd1—Br2	105.13 (2)	N2—C4—H4B	109.7
Br4—Cd1—Br2	110.50 (2)	C3—C4—H4B	109.7
Br1—Cd1—Br2	98.28 (2)	H4A—C4—H4B	108.2
N3—C8—C7	178.1 (5)	N1—C3—C4	108.4 (3)

C7—N2—C6	111.2 (3)	N1—C3—H3A	110.0
C7—N2—C4	111.4 (3)	C4—C3—H3A	110.0
C6—N2—C4	109.0 (3)	N1—C3—H3B	110.0
C7—N2—C2	108.3 (3)	C4—C3—H3B	110.0
C6—N2—C2	108.4 (4)	H3A—C3—H3B	108.4
C4—N2—C2	108.5 (3)	C1—C2—N2	109.2 (3)
N1—C5—C6	108.6 (4)	C1—C2—H2A	109.8
N1—C5—H5A	110.0	N2—C2—H2A	109.8
C6—C5—H5A	110.0	C1—C2—H2B	109.8
N1—C5—H5B	110.0	N2—C2—H2B	109.8
C6—C5—H5B	110.0	H2A—C2—H2B	108.3
H5A—C5—H5B	108.3	C8—C7—N2	111.5 (4)
C5—N1—C1	110.3 (3)	C8—C7—H7A	109.3
C5—N1—C3	110.1 (3)	N2—C7—H7A	109.3
C1—N1—C3	109.2 (3)	C8—C7—H7B	109.3
C5—N1—H1C	114.6	N2—C7—H7B	109.3
C1—N1—H1C	110.2	H7A—C7—H7B	108.0
C3—N1—H1C	102.0	N1—C1—C2	109.4 (3)
N2—C6—C5	110.1 (4)	N1—C1—H1A	109.8
N2—C6—H6A	109.6	C2—C1—H1A	109.8
C5—C6—H6A	109.6	N1—C1—H1B	109.8
N2—C6—H6B	109.6	C2—C1—H1B	109.8
C5—C6—H6B	109.6	H1A—C1—H1B	108.2
H6A—C6—H6B	108.1		

Hydrogen-bond geometry (Å, °)

<i>D</i> —H··· <i>A</i>	<i>D</i> —H	H··· <i>A</i>	<i>D</i> ··· <i>A</i>	<i>D</i> —H··· <i>A</i>
N1—H1C···Br1 ⁱ	0.90	2.85	3.466 (4)	127
N1—H1C···Br2 ⁱⁱ	0.90	2.69	3.325 (4)	128
N1—H1C···Br4 ⁱ	0.90	3.11	3.711 (4)	126
C2—H2B···Br3 ⁱⁱⁱ	0.97	2.83	3.765 (5)	162
C4—H4B···Br1	0.97	2.85	3.643 (4)	140
C7—H7A···Br3 ^{iv}	0.97	2.90	3.626 (4)	132
C7—H7B···Br2 ⁱⁱⁱ	0.97	2.78	3.683 (4)	154

Symmetry codes: (i) $-x+2, -y+1, -z+1$; (ii) $-x+2, y+1/2, -z+3/2$; (iii) $-x+1, y+1/2, -z+3/2$; (iv) $x, -y+1/2, z+1/2$.

supplementary materials

Fig. 1




Fig. 2

