

Bis(μ -bis{[4-(2-pyridyl)pyrimidin-2-yl]-sulfanyl}methane)disilver(I) bis(perchlorate)

Hai-Bin Zhu

School of Chemistry and Chemical Engineering, Southeast University, Nanjing 211189, People's Republic of China
Correspondence e-mail: zhuhaibin@seu.edu.cn


Received 20 November 2010; accepted 25 November 2010

Key indicators: single-crystal X-ray study; $T = 298\text{ K}$; mean $\sigma(\text{C}-\text{C}) = 0.009\text{ \AA}$; disorder in solvent or counterion; R factor = 0.046; wR factor = 0.172; data-to-parameter ratio = 16.0.

In the macrocyclic centrosymmetric dinuclear complex, $[\text{Ag}_2(\text{C}_{19}\text{H}_{14}\text{N}_6\text{S}_2)_2](\text{ClO}_4)_2$, the Ag^{I} atom, bis{[4-(2-pyridyl)pyrimidin-2-yl]sulfanyl}methane (2-bppt) ligand and perchlorate anion each lie on a twofold rotation axis. The 2-bppt ligand chelates two four-coordinated Ag^{I} atoms through its two bipyridine-like arms. The O atoms of the perchlorate anion are disordered each over two positions of equal occupancy. Adjacent complex molecules are linked by $\pi-\pi$ interactions between the pyridine and pyrimidine rings [centroid-centroid distance = 3.663 (8) \AA].

Related literature

For $\text{Ag}(\text{I})$ coordination polymers, see: Chen *et al.* (2006). For the coordination chemistry of 4-(pyridin-*n*-yl)pyrimidin-2-thiol (*n* = 2, 3, 4) and their derivatives, see: Dong *et al.* (2009); Huang *et al.* (2007); Zhu *et al.* (2010).



Experimental

Crystal data

$[\text{Ag}_2(\text{C}_{19}\text{H}_{14}\text{N}_6\text{S}_2)_2](\text{ClO}_4)_2$	$V = 8761 (2)\text{ \AA}^3$
$M_r = 1195.64$	$Z = 8$
Orthorhombic, $Fdd2$	Mo $K\alpha$ radiation
$a = 10.4382 (16)\text{ \AA}$	$\mu = 1.27\text{ mm}^{-1}$
$b = 27.896 (4)\text{ \AA}$	$T = 298\text{ K}$
$c = 30.089 (5)\text{ \AA}$	$0.15 \times 0.12 \times 0.10\text{ mm}$

Data collection

Bruker APEXII CCD diffractometer	14503 measured reflections
Absorption correction: multi-scan (<i>SADABS</i> ; Bruker, 2001)	2705 independent reflections
$(SADABS$; Bruker, 2001)	1640 reflections with $I > 2\sigma(I)$
$T_{\min} = 0.832$, $T_{\max} = 0.880$	$R_{\text{int}} = 0.035$

Refinement

$R[F^2 > 2\sigma(F^2)] = 0.046$	24 restraints
$wR(F^2) = 0.172$	H-atom parameters constrained
$S = 1.05$	$\Delta\rho_{\text{max}} = 0.54\text{ e \AA}^{-3}$
2705 reflections	$\Delta\rho_{\text{min}} = -0.64\text{ e \AA}^{-3}$
169 parameters	

Table 1
Selected bond lengths (\AA).

Ag1–N1	2.277 (4)	Ag1–N2	2.398 (3)

Data collection: *APEX2* (Bruker, 2007); cell refinement: *SAINT-Plus* (Bruker, 2007); data reduction: *SAINT-Plus*; program(s) used to solve structure: *SHELXS97* (Sheldrick, 2008); program(s) used to refine structure: *SHELXL97* (Sheldrick, 2008); molecular graphics: *SHELXTL* (Sheldrick, 2008); software used to prepare material for publication: *SHELXTL*.

The author acknowledges financial support from the China Postdoctoral Research Fund (20070411010).

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: HY2383).

References

- Bruker (2001). *SADABS*. Bruker AXS Inc., Madison, Wisconsin, USA.
- Bruker (2007). *APEX2* and *SAINT-Plus*. Bruker AXS Inc., Madison, Wisconsin, USA.
- Chen, C.-L., Kang, B.-S. & Su, C.-Y. (2006). *Aust. J. Chem.* **59**, 3–18.
- Dong, H.-Z., Zhao, J., Gou, S.-H. & Zhu, H.-B. (2009). *Polyhedron*, **28**, 1040–1048.
- Huang, C.-H., Gou, S.-H., Zhu, H.-B. & Huang, W. (2007). *Inorg. Chem.* **46**, 5537–5543.
- Sheldrick, G. M. (2008). *Acta Cryst. A* **64**, 112–122.
- Zhu, H.-B., Li, L., Xu, G. & Gou, S.-H. (2010). *Eur. J. Inorg. Chem.* pp. 1143–1148.

supplementary materials

Acta Cryst. (2010). E66, m1691 [doi:10.1107/S160053681004924X]

Bis(μ -bis{[4-(2-pyridyl)pyrimidin-2-yl]sulfanyl}methane)disilver(I) bis(perchlorate)

H.-B. Zhu

Comment

The supramolecular chemistry of Ag(I) coordination polymers is being a dynamic and thriving research field, which has attracted considerable interest (Chen *et al.*, 2006). For a long time, we have focused on the coordination chemistry of 4-(pyridin-*n*-yl)pyrimidin-2-thiol (*n* = 2, 3, 4) and their derivatives (Dong *et al.*, 2009; Huang *et al.*, 2007; Zhu *et al.*, 2010). Herein, we report a macrocyclic Ag(I) complex with bis[4-(2-pyridyl)pyrimidin-2-ylthio]methane (2-bppt) ligand.

The title compound shows a discrete macrocyclic dinuclear structure, with perchlorate anions uncoordinated (Fig. 1). Each Ag^I ion is chelated by two sets of N,N-chelating donors from two 2-bppt ligands. The Ag—N bond distances are 2.277 (4) and 2.398 (3) Å (Table 1), while the N—Ag—N angles are in the range of 70.96 (13) to 158.7 (2)^o. The Ag—Ag separation across the macrocycle is 8.167 (1) Å.

Experimental

A CH₃CN solution of AgClO₄ (0.1 mmol) was layered above a CH₂Cl₂ solution of 2-bppt (0.1 mmol). Colorless crystals were obtained after one week. The crystals were collected and dried under vacuum (yield: 46%).

Refinement

All H atoms were positioned geometrically and allowed to ride on their parent atoms, with C—H = 0.93 and 0.97 Å and with *U*_{iso}(H) = 1.2*U*_{eq}(C).

Figures




Fig. 1. Molecular structure of the title compound, with the 30% probability displacement ellipsoids. [Symmetry codes: (A) 7/4 - *x*, 3/4 - *y*, *z*; (B) *x*, 3/4 - *y*, -1/4 - *z*; (C) 7/4 - *x*, *y*, -1/4 - *z*; (D) 9/4 - *x*, 1/4 - *y*, *z*.]

Bis(μ -bis{[4-(2-pyridyl)pyrimidin-2-yl]sulfanyl}methane)disilver(I) bis(perchlorate)

Crystal data

[Ag₂(C₁₉H₁₄N₆S₂)₂](ClO₄)₂

F(000) = 4768

*M*_r = 1195.64

*D*_x = 1.813 Mg m⁻³

supplementary materials

Orthorhombic, $Fddd$	Mo $K\alpha$ radiation, $\lambda = 0.71073 \text{ \AA}$
Hall symbol: -F 2uv 2vw	Cell parameters from 2705 reflections
$a = 10.4382 (16) \text{ \AA}$	$\theta = 2.3\text{--}25.5^\circ$
$b = 27.896 (4) \text{ \AA}$	$\mu = 1.27 \text{ mm}^{-1}$
$c = 30.089 (5) \text{ \AA}$	$T = 298 \text{ K}$
$V = 8761 (2) \text{ \AA}^3$	Block, colorless
$Z = 8$	$0.15 \times 0.12 \times 0.10 \text{ mm}$

Data collection

Bruker APEXII CCD diffractometer	2705 independent reflections
Radiation source: fine-focus sealed tube graphite	1640 reflections with $I > 2\sigma(I)$
φ and ω scans	$R_{\text{int}} = 0.035$
Absorption correction: multi-scan (<i>SADABS</i> ; Bruker, 2001)	$\theta_{\text{max}} = 28.3^\circ, \theta_{\text{min}} = 2.2^\circ$
$T_{\text{min}} = 0.832, T_{\text{max}} = 0.880$	$h = -13\text{--}12$
14503 measured reflections	$k = -36\text{--}35$
	$l = -35\text{--}40$

Refinement

Refinement on F^2	Primary atom site location: structure-invariant direct methods
Least-squares matrix: full	Secondary atom site location: difference Fourier map
$R[F^2 > 2\sigma(F^2)] = 0.046$	Hydrogen site location: inferred from neighbouring sites
$wR(F^2) = 0.172$	H-atom parameters constrained
$S = 1.05$	$w = 1/[\sigma^2(F_o^2) + (0.1P)^2]$ where $P = (F_o^2 + 2F_c^2)/3$
2705 reflections	$(\Delta/\sigma)_{\text{max}} < 0.001$
169 parameters	$\Delta\rho_{\text{max}} = 0.54 \text{ e \AA}^{-3}$
24 restraints	$\Delta\rho_{\text{min}} = -0.64 \text{ e \AA}^{-3}$

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (\AA^2)

	x	y	z	$U_{\text{iso}}^*/U_{\text{eq}}$	Occ. (<1)
Ag1	0.8750	0.3750	0.010711 (18)	0.0724 (3)	
S1	0.64901 (13)	0.37857 (5)	-0.07449 (5)	0.0703 (4)	
Cl1	1.1250	0.1250	-0.08929 (11)	0.1119 (9)	
C6	0.8563 (4)	0.27205 (15)	-0.03645 (15)	0.0602 (11)	
N2	0.7995 (3)	0.31493 (12)	-0.03991 (12)	0.0577 (9)	
N3	0.6715 (4)	0.28754 (15)	-0.10059 (13)	0.0708 (10)	
C9	0.7124 (4)	0.32014 (15)	-0.07181 (14)	0.0598 (10)	
C5	0.9565 (5)	0.26651 (16)	-0.00209 (16)	0.0652 (12)	
C10	0.5575 (6)	0.3750	-0.1250	0.0715 (18)	
H10A	0.5025	0.4030	-0.1267	0.086*	0.50
H10B	0.5025	0.3470	-0.1233	0.086*	0.50

N1	0.9792 (4)	0.30490 (14)	0.02472 (13)	0.0693 (10)	
C8	0.7272 (6)	0.24468 (18)	-0.09552 (17)	0.0804 (15)	
H8	0.7013	0.2199	-0.1141	0.096*	
C7	0.8198 (6)	0.23519 (16)	-0.06458 (18)	0.0761 (14)	
H7	0.8571	0.2050	-0.0625	0.091*	
C3	1.1171 (7)	0.2208 (3)	0.0336 (3)	0.107 (2)	
H3	1.1630	0.1924	0.0365	0.128*	
C1	1.0704 (5)	0.3010 (2)	0.05571 (19)	0.0873 (15)	
H1	1.0864	0.3268	0.0745	0.105*	
C2	1.1432 (5)	0.2585 (3)	0.0605 (3)	0.105 (2)	
H2	1.2078	0.2565	0.0817	0.125*	
C4	1.0244 (6)	0.2245 (2)	0.0026 (2)	0.0892 (16)	
H4	1.0062	0.1985	-0.0158	0.107*	
O1	1.1499 (10)	0.1740 (3)	-0.0861 (3)	0.120 (3)	0.50
O2	1.2437 (12)	0.1043 (4)	-0.0898 (4)	0.142 (4)	0.50
O3	1.0656 (13)	0.1194 (5)	-0.0423 (5)	0.166 (5)	0.50
O4	1.0504 (13)	0.1200 (6)	-0.1265 (5)	0.176 (5)	0.50

Atomic displacement parameters (\AA^2)

	U^{11}	U^{22}	U^{33}	U^{12}	U^{13}	U^{23}
Ag1	0.0848 (4)	0.0529 (3)	0.0795 (4)	-0.0060 (2)	0.000	0.000
S1	0.0718 (8)	0.0712 (7)	0.0679 (8)	0.0074 (5)	-0.0057 (6)	-0.0091 (6)
Cl1	0.1036 (17)	0.0619 (12)	0.170 (3)	0.0054 (11)	0.000	0.000
C6	0.068 (3)	0.051 (2)	0.062 (3)	-0.0062 (19)	0.026 (2)	0.0064 (19)
N2	0.059 (2)	0.0522 (18)	0.062 (2)	-0.0070 (15)	0.0100 (18)	-0.0039 (15)
N3	0.074 (2)	0.071 (3)	0.067 (2)	-0.022 (2)	0.007 (2)	-0.0131 (19)
C9	0.058 (2)	0.063 (2)	0.059 (2)	-0.0073 (19)	0.012 (2)	-0.006 (2)
C5	0.065 (3)	0.062 (3)	0.069 (3)	0.002 (2)	0.023 (2)	0.013 (2)
C10	0.050 (3)	0.089 (5)	0.076 (4)	0.000	0.000	0.000 (3)
N1	0.066 (2)	0.068 (2)	0.074 (2)	-0.0072 (19)	0.004 (2)	0.0169 (19)
C8	0.095 (4)	0.075 (3)	0.071 (3)	-0.026 (3)	0.020 (3)	-0.020 (3)
C7	0.091 (4)	0.050 (2)	0.087 (4)	-0.008 (2)	0.034 (3)	-0.007 (2)
C3	0.092 (5)	0.102 (5)	0.126 (6)	0.029 (4)	0.023 (4)	0.036 (5)
C1	0.074 (3)	0.101 (4)	0.087 (4)	-0.006 (3)	-0.007 (3)	0.023 (3)
C2	0.065 (4)	0.129 (6)	0.120 (5)	0.007 (3)	-0.003 (3)	0.050 (5)
C4	0.087 (4)	0.081 (3)	0.099 (4)	0.022 (3)	0.027 (3)	0.019 (3)
O1	0.152 (7)	0.076 (5)	0.132 (7)	0.008 (5)	-0.029 (5)	-0.009 (4)
O2	0.133 (7)	0.105 (6)	0.188 (8)	0.044 (6)	-0.017 (6)	-0.012 (6)
O3	0.176 (9)	0.166 (8)	0.155 (8)	-0.016 (7)	0.028 (7)	0.007 (7)
O4	0.167 (8)	0.198 (9)	0.162 (8)	-0.025 (8)	-0.076 (7)	-0.041 (7)

Geometric parameters (\AA , $^\circ$)

Ag1—N1	2.277 (4)	C5—C4	1.377 (6)
Ag1—N2	2.398 (3)	C10—S1 ⁱ	1.798 (4)
S1—C9	1.761 (4)	C10—H10A	0.9700
S1—C10	1.798 (4)	C10—H10B	0.9700

supplementary materials

Cl1—O4	1.370 (11)	N1—C1	1.337 (6)
Cl1—O2	1.367 (11)	C8—C7	1.368 (8)
Cl1—O1	1.395 (9)	C8—H8	0.9300
Cl1—O3	1.552 (13)	C7—H7	0.9300
C6—N2	1.339 (5)	C3—C2	1.355 (11)
C6—C7	1.385 (7)	C3—C4	1.348 (9)
C6—C5	1.479 (7)	C3—H3	0.9300
N2—C9	1.330 (5)	C1—C2	1.416 (9)
N3—C9	1.326 (5)	C1—H1	0.9300
N3—C8	1.338 (6)	C2—H2	0.9300
C5—N1	1.361 (6)	C4—H4	0.9300
N1—Ag1—N1 ⁱⁱ	158.7 (2)	S1 ⁱ —C10—H10A	108.3
N1—Ag1—N2	71.00 (14)	S1—C10—H10B	108.3
N1 ⁱⁱ —Ag1—N2	124.11 (13)	S1 ⁱ —C10—H10B	108.3
N1—Ag1—N2 ⁱⁱ	124.11 (13)	H10A—C10—H10B	107.4
N1 ⁱⁱ —Ag1—N2 ⁱⁱ	71.00 (14)	C5—N1—C1	118.3 (5)
N2—Ag1—N2 ⁱⁱ	101.13 (16)	C5—N1—Ag1	118.9 (3)
C9—S1—C10	100.78 (17)	C1—N1—Ag1	122.6 (4)
O4—Cl1—O4 ⁱⁱⁱ	70.5 (13)	N3—C8—C7	123.9 (4)
O4—Cl1—O2	117.6 (10)	N3—C8—H8	118.1
O4 ⁱⁱⁱ —Cl1—O2	61.2 (7)	C7—C8—H8	118.1
O4—Cl1—O2 ⁱⁱⁱ	61.2 (7)	C8—C7—C6	117.8 (5)
O4 ⁱⁱⁱ —Cl1—O2 ⁱⁱⁱ	117.6 (10)	C8—C7—H7	121.1
O2—Cl1—O2 ⁱⁱⁱ	178.7 (11)	C6—C7—H7	121.1
O4—Cl1—O1 ⁱⁱⁱ	81.5 (8)	C2—C3—C4	119.8 (6)
O4 ⁱⁱⁱ —Cl1—O1 ⁱⁱⁱ	105.2 (8)	C2—C3—H3	120.1
O2—Cl1—O1 ⁱⁱⁱ	75.8 (6)	C4—C3—H3	120.1
O2 ⁱⁱⁱ —Cl1—O1 ⁱⁱⁱ	104.3 (6)	N1—C1—C2	121.3 (6)
O4—Cl1—O1	105.2 (8)	N1—C1—H1	119.3
O4 ⁱⁱⁱ —Cl1—O1	81.5 (8)	C2—C1—H1	119.3
O2—Cl1—O1	104.3 (6)	C3—C2—C1	118.9 (6)
O2 ⁱⁱⁱ —Cl1—O1	75.8 (6)	C3—C2—H2	120.6
O1 ⁱⁱⁱ —Cl1—O1	172.0 (9)	C1—C2—H2	120.6
O4—Cl1—O3 ⁱⁱⁱ	168.9 (8)	C3—C4—C5	120.3 (6)
O4 ⁱⁱⁱ —Cl1—O3 ⁱⁱⁱ	120.5 (8)	C3—C4—H4	119.8
O2—Cl1—O3 ⁱⁱⁱ	72.0 (7)	C5—C4—H4	119.8
O2 ⁱⁱⁱ —Cl1—O3 ⁱⁱⁱ	109.3 (9)	Cl1—O1—O2 ⁱⁱⁱ	51.3 (4)
O1 ⁱⁱⁱ —Cl1—O3 ⁱⁱⁱ	96.3 (7)	Cl1—O1—O4 ⁱⁱⁱ	48.6 (5)
O1—Cl1—O3 ⁱⁱⁱ	76.3 (7)	O2 ⁱⁱⁱ —O1—O4 ⁱⁱⁱ	83.8 (7)
O4—Cl1—O3	120.5 (8)	Cl1—O1—O3 ⁱⁱⁱ	55.7 (6)
O4 ⁱⁱⁱ —Cl1—O3	168.9 (8)	O2 ⁱⁱⁱ —O1—O3 ⁱⁱⁱ	85.0 (7)
O2—Cl1—O3	109.3 (9)	O4 ⁱⁱⁱ —O1—O3 ⁱⁱⁱ	88.7 (8)
O2 ⁱⁱⁱ —Cl1—O3	72.0 (7)	Cl1—O2—O4 ⁱⁱⁱ	59.5 (7)

O1 ⁱⁱⁱ —Cl1—O3	76.3 (7)	Cl1—O2—O1 ⁱⁱⁱ	52.8 (5)
O1—Cl1—O3	96.3 (7)	O4 ⁱⁱⁱ —O2—O1 ⁱⁱⁱ	90.0 (9)
O3 ⁱⁱⁱ —Cl1—O3	48.6 (10)	Cl1—O2—O3 ⁱⁱⁱ	59.0 (6)
N2—C6—C7	119.6 (5)	O4 ⁱⁱⁱ —O2—O3 ⁱⁱⁱ	108.6 (10)
N2—C6—C5	117.4 (4)	O1 ⁱⁱⁱ —O2—O3 ⁱⁱⁱ	80.0 (8)
C7—C6—C5	123.0 (4)	O3 ⁱⁱⁱ —O3—Cl1	65.7 (5)
C9—N2—C6	117.2 (4)	O3 ⁱⁱⁱ —O3—O2 ⁱⁱⁱ	104.2 (9)
C9—N2—Ag1	127.3 (3)	Cl1—O3—O2 ⁱⁱⁱ	49.0 (5)
C6—N2—Ag1	115.4 (3)	O3 ⁱⁱⁱ —O3—O1 ⁱⁱⁱ	88.3 (12)
C9—N3—C8	113.5 (4)	Cl1—O3—O1 ⁱⁱⁱ	48.0 (5)
N3—C9—N2	128.0 (4)	O2 ⁱⁱⁱ —O3—O1 ⁱⁱⁱ	75.8 (7)
N3—C9—S1	119.0 (4)	Cl1—O4—O2 ⁱⁱⁱ	59.3 (7)
N2—C9—S1	113.0 (3)	Cl1—O4—O4 ⁱⁱⁱ	54.8 (6)
N1—C5—C4	121.3 (5)	O2 ⁱⁱⁱ —O4—O4 ⁱⁱⁱ	103.6 (10)
N1—C5—C6	117.1 (4)	Cl1—O4—O1 ⁱⁱⁱ	49.9 (6)
C4—C5—C6	121.6 (5)	O2 ⁱⁱⁱ —O4—O1 ⁱⁱⁱ	84.9 (9)
S1—C10—S1 ⁱ	115.8 (4)	O4 ⁱⁱⁱ —O4—O1 ⁱⁱⁱ	80.6 (11)
S1—C10—H10A	108.3		

Symmetry codes: (i) $x, -y+3/4, -z-1/4$; (ii) $-x+7/4, -y+3/4, z$; (iii) $-x+9/4, -y+1/4, z$.

supplementary materials

Fig. 1

