

## 1-(2-Hydroxybenzoyl)thiosemicarbazide hemihydrate

Ming-zhi Song\* and Chuangang Fan

College of Chemistry and Chemical Technology, Binzhou University, Binzhou 256600, Shandong, People's Republic of China  
Correspondence e-mail: fanchuangang2009@163.com


Received 14 October 2010; accepted 29 October 2010

Key indicators: single-crystal X-ray study;  $T = 298$  K; mean  $\sigma(\text{C}-\text{C}) = 0.005$  Å;  $R$  factor = 0.047;  $wR$  factor = 0.122; data-to-parameter ratio = 13.3.

The asymmetric unit of the title compound,  $\text{C}_8\text{H}_9\text{N}_3\text{O}_2\text{S} \cdot 0.5\text{H}_2\text{O}$ , contains two thiosemicarbazide molecules with the short distance of 3.521 (3) Å between the centroids of the benzene rings, and one water molecule. In the two independent molecules, the benzene rings and the thiosemicarbazone fragments are twisted at 9.2 (3) and 18.5 (3)°. An extensive three-dimensional hydrogen-bonding network, formed by intermolecular  $\text{N}-\text{H} \cdots \text{O}$ ,  $\text{N}-\text{H} \cdots \text{S}$  and  $\text{O}-\text{H} \cdots \text{O}$  hydrogen bonds, consolidates the crystal packing.

### Related literature

For the biological activities of thiosemicarbazide derivatives, see: Desai *et al.* (1984); Shukla *et al.* (1984). For related structures, see: Gors *et al.* (1979); Jin (2007).



### Experimental

#### Crystal data

$\text{C}_8\text{H}_9\text{N}_3\text{O}_2\text{S} \cdot 0.5\text{H}_2\text{O}$   
 $M_r = 440.50$   
Monoclinic,  $P2_1/n$   
 $a = 9.0718$  (11) Å  
 $b = 21.608$  (2) Å  
 $c = 10.1035$  (13) Å  
 $\beta = 90.173$  (1)°

$V = 1980.5$  (4) Å<sup>3</sup>  
 $Z = 4$   
Mo  $K\alpha$  radiation  
 $\mu = 0.31$  mm<sup>-1</sup>  
 $T = 298$  K  
 $0.45 \times 0.20 \times 0.17$  mm

#### Data collection

Bruker SMART APEX CCD  
area-detector diffractometer  
Absorption correction: multi-scan  
(*SADABS*; Sheldrick, 1996)  
 $T_{\min} = 0.873$ ,  $T_{\max} = 0.949$

9862 measured reflections  
3487 independent reflections  
2202 reflections with  $I > 2\sigma(I)$   
 $R_{\text{int}} = 0.041$

#### Refinement

$R[F^2 > 2\sigma(F^2)] = 0.047$   
 $wR(F^2) = 0.122$   
 $S = 1.01$   
3487 reflections

262 parameters  
H-atom parameters constrained  
 $\Delta\rho_{\text{max}} = 0.37$  e Å<sup>-3</sup>  
 $\Delta\rho_{\text{min}} = -0.33$  e Å<sup>-3</sup>

**Table 1**

Hydrogen-bond geometry (Å, °).

$D-\text{H} \cdots A$	$D-\text{H}$	$\text{H} \cdots A$	$D \cdots A$	$D-\text{H} \cdots A$
$\text{N2}-\text{H2} \cdots \text{S1}^{\text{i}}$	0.86	2.58	3.361 (3)	151
$\text{N5}-\text{H5} \cdots \text{S1}^{\text{ii}}$	0.86	2.83	3.404 (3)	125
$\text{N6}-\text{H6A} \cdots \text{O2}^{\text{iii}}$	0.86	2.16	2.984 (3)	160
$\text{N6}-\text{H6B} \cdots \text{S2}^{\text{iv}}$	0.86	2.85	3.647 (3)	156
$\text{O4}-\text{H4} \cdots \text{O1}^{\text{ii}}$	0.82	1.93	2.724 (3)	162
$\text{O5}-\text{H5B} \cdots \text{O5}^{\text{v}}$	0.85	1.95	2.728 (10)	152
$\text{N3}-\text{H3A} \cdots \text{O3}$	0.86	2.12	2.918 (3)	154
$\text{N3}-\text{H3B} \cdots \text{O5}$	0.86	2.30	3.110 (5)	156
$\text{N5}-\text{H5} \cdots \text{O4}$	0.86	1.99	2.629 (3)	130
$\text{O2}-\text{H2A} \cdots \text{O1}$	0.82	1.83	2.552 (3)	146
$\text{O5}-\text{H5D} \cdots \text{S1}$	0.85	2.66	3.398 (4)	147

Symmetry codes: (i)  $-x + 1, -y, -z + 1$ ; (ii)  $x + 1, y, z$ ; (iii)  $x + \frac{1}{2}, -y + \frac{1}{2}, z - \frac{1}{2}$ ; (iv)  $-x + 2, -y, -z$ ; (v)  $-x + 1, -y, -z$ .

Data collection: *SMART* (Bruker, 2007); cell refinement: *SAINT* (Bruker, 2007); data reduction: *SAINT*; program(s) used to solve structure: *SHELXS97* (Sheldrick, 2008); program(s) used to refine structure: *SHELXL97* (Sheldrick, 2008); molecular graphics: *SHELXTL* (Sheldrick, 2008); software used to prepare material for publication: *SHELXTL*.

The authors acknowledge the financial support of the Foundation of Binzhou University (grant No. BZXYQNLG200501).

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: CV2777).

### References

- Bruker (2007). *SMART* and *SAINT*. Bruker AXS Inc., Madison, Wisconsin, USA.
- Desai, N. C., Shukla, H. K., Paresh, B. P. & Thaker, K. A. (1984). *J. Indian Chem. Soc.* pp. 455–457.
- Gors, C., Baert, F., Henichart, J. P. & Houssin, R. (1979). *J. Mol. Struct.* **55**, 223–233.
- Jin, L.-F. (2007). *Acta Cryst.* **E63**, o3465.
- Sheldrick, G. M. (1996). *SADABS*. University of Göttingen, Germany.
- Sheldrick, G. M. (2008). *Acta Cryst.* **A64**, 112–122.
- Shukla, H. K., Desai, N. C., Astik, R. R. & Thaker, K. A. (1984). *J. Indian Chem. Soc.* pp. 168–196.

**supplementary materials**

*Acta Cryst.* (2010). E66, o3048 [ doi:10.1107/S1600536810044259 ]

## 1-(2-Hydroxybenzoyl)thiosemicarbazide hemihydrate

M. Song and C. Fan

### Comment

Thiosemicarbazide compounds exhibit various biological activities such as anti-bacterial, anti-fungal and especially anti-tuberculosis (Shukla *et al.*, 1984; Desai *et al.*, 1984). Herewith we present the title compound (I), a new thiosemicarbazide compound.

The asymmetric unit of (I) contains two independent molecules and one crystalline water molecule (Fig. 1). The bond lengths and angles are normal and comparable to the values observed in similar compounds (Gors *et al.*, 1979; Jin, 2007). In two independent molecules, the benzene rings and the thiosemicarbazone fragments (C/N/N) are twisted at 9.2 (3)° and 18.5 (3)°, respectively.

In the crystal structure, the three-dimensional network structure was formed by the intermolecular N—H···O, N—H···S and O—H···O hydrogen bonds (Table 1).

### Experimental

Salicylyl hydrazine (10 mmol), potassium thiocyanate (12 mmol) and 10 ml methanol-water (1:1) were mixed in 50 ml flask. After refluxing 12 h at 373 K, the resulting mixture was recrystallized from solution, affording the title compound as a colorless crystalline solid.

### Refinement

All H atoms were placed in geometrically idealized positions (N—H = 0.86, O—H = 0.82–0.85 and C—H = 0.93 Å) and treated as riding on their parent atoms, with  $U_{\text{iso}}(\text{H}) = 1.2U_{\text{eq}} + 1.5U_{\text{eq}}$  of the parent atom.

### Figures




Fig. 1. The content of asymmetric unit of the title compound showing the atomic numbering scheme and 30% probability displacement ellipsoids.

## 1-(2-Hydroxybenzoyl)thiosemicarbazide hemihydrate

### Crystal data

C<sub>16</sub>H<sub>20</sub>N<sub>6</sub>O<sub>5</sub>S<sub>2</sub>

$F(000) = 916$

# supplementary materials

---

$$M_r = 440.50$$

Monoclinic,  $P2_1/n$

$$a = 9.0718 (11) \text{ \AA}$$

$$b = 21.608 (2) \text{ \AA}$$

$$c = 10.1035 (13) \text{ \AA}$$

$$\beta = 90.173 (1)^\circ$$

$$V = 1980.5 (4) \text{ \AA}^3$$

$$Z = 4$$

$$D_x = 1.474 \text{ Mg m}^{-3}$$

Mo  $K\alpha$  radiation,  $\lambda = 0.71073 \text{ \AA}$

Cell parameters from 2367 reflections

$$\theta = 2.2\text{--}25.7^\circ$$

$$\mu = 0.31 \text{ mm}^{-1}$$

$$T = 298 \text{ K}$$

Block, yellow

$$0.45 \times 0.20 \times 0.17 \text{ mm}$$

## Data collection

Bruker SMART APEX CCD area-detector diffractometer

Radiation source: fine-focus sealed tube graphite

$\varphi$  and  $\omega$  scans

Absorption correction: multi-scan (*SADABS*; Sheldrick, 1996)

$$T_{\min} = 0.873, T_{\max} = 0.949$$

9862 measured reflections

3487 independent reflections

2202 reflections with  $I > 2\sigma(I)$

$$R_{\text{int}} = 0.041$$

$$\theta_{\max} = 25.0^\circ, \theta_{\min} = 1.9^\circ$$

$$h = -10 \rightarrow 10$$

$$k = -21 \rightarrow 25$$

$$l = -11 \rightarrow 12$$

## Refinement

Refinement on  $F^2$

Least-squares matrix: full

$$R[F^2 > 2\sigma(F^2)] = 0.047$$

$$wR(F^2) = 0.122$$

$$S = 1.01$$

3487 reflections

262 parameters

0 restraints

Primary atom site location: structure-invariant direct methods

Secondary atom site location: difference Fourier map

Hydrogen site location: inferred from neighbouring sites

H-atom parameters constrained

$$w = 1/[\sigma^2(F_o^2) + (0.0468P)^2 + 1.2463P]$$

$$\text{where } P = (F_o^2 + 2F_c^2)/3$$

$$(\Delta/\sigma)_{\max} < 0.001$$

$$\Delta\rho_{\max} = 0.37 \text{ e \AA}^{-3}$$

$$\Delta\rho_{\min} = -0.32 \text{ e \AA}^{-3}$$

## Special details

**Geometry.** All e.s.d.'s (except the e.s.d. in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell e.s.d.'s are taken into account individually in the estimation of e.s.d.'s in distances, angles and torsion angles; correlations between e.s.d.'s in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell e.s.d.'s is used for estimating e.s.d.'s involving l.s. planes.

**Refinement.** Refinement of  $F^2$  against ALL reflections. The weighted  $R$ -factor  $wR$  and goodness of fit  $S$  are based on  $F^2$ , conventional  $R$ -factors  $R$  are based on  $F$ , with  $F$  set to zero for negative  $F^2$ . The threshold expression of  $F^2 > \sigma(F^2)$  is used only for calculating  $R$ -factors(gt) *etc.* and is not relevant to the choice of reflections for refinement.  $R$ -factors based on  $F^2$  are statistically about twice as large as those based on  $F$ , and  $R$ -factors based on ALL data will be even larger.

*Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters ( $\text{\AA}^2$ )*

	<i>x</i>	<i>y</i>	<i>z</i>	$U_{\text{iso}}^*/U_{\text{eq}}$	Occ. (<1)
N1	0.4021 (3)	0.07227 (12)	0.5682 (3)	0.0401 (7)	
H1	0.3330	0.0634	0.6231	0.048*	
N2	0.5273 (3)	0.10117 (11)	0.6152 (3)	0.0392 (7)	
H2	0.6033	0.0794	0.6354	0.047*	
N3	0.4849 (3)	0.07466 (12)	0.3544 (3)	0.0443 (7)	
H3A	0.5612	0.0950	0.3806	0.053*	
H3B	0.4746	0.0655	0.2721	0.053*	
N4	0.8800 (3)	0.06010 (11)	0.3372 (3)	0.0399 (7)	
H4'	0.8275	0.0385	0.3908	0.048*	
N5	0.9398 (3)	0.11556 (11)	0.3803 (3)	0.0362 (7)	
H5	1.0322	0.1177	0.3992	0.043*	
N6	0.9674 (3)	0.07614 (13)	0.1292 (3)	0.0527 (8)	
H6A	0.9949	0.1125	0.1534	0.063*	
H6B	0.9827	0.0639	0.0495	0.063*	
O1	0.4251 (2)	0.19444 (10)	0.5943 (2)	0.0386 (6)	
O2	0.5712 (3)	0.29296 (10)	0.6415 (2)	0.0441 (6)	
H2A	0.5042	0.2722	0.6100	0.066*	
O3	0.7237 (2)	0.16510 (10)	0.3629 (2)	0.0439 (6)	
O4	1.1396 (2)	0.17295 (9)	0.5240 (2)	0.0408 (6)	
H4	1.2180	0.1806	0.5612	0.061*	
O5	0.3935 (7)	0.0087 (2)	0.0932 (4)	0.172 (2)	
H5D	0.3575	-0.0063	0.1638	0.206*	0.50
H5B	0.4780	0.0030	0.0591	0.206*	0.50
S1	0.23175 (9)	0.01686 (4)	0.39559 (9)	0.0415 (3)	
S2	0.84286 (14)	-0.03260 (5)	0.17448 (11)	0.0703 (4)	
C1	0.3839 (4)	0.05757 (13)	0.4406 (3)	0.0336 (8)	
C2	0.5320 (3)	0.16295 (14)	0.6297 (3)	0.0307 (7)	
C3	0.6647 (3)	0.19040 (13)	0.6870 (3)	0.0287 (7)	
C4	0.6763 (3)	0.25507 (14)	0.6910 (3)	0.0315 (7)	
C5	0.7998 (4)	0.28273 (16)	0.7474 (3)	0.0404 (8)	
H5A	0.8066	0.3256	0.7513	0.049*	
C6	0.9113 (4)	0.24687 (18)	0.7971 (3)	0.0448 (9)	
H6	0.9938	0.2657	0.8342	0.054*	
C7	0.9031 (4)	0.18322 (17)	0.7929 (3)	0.0445 (9)	
H7	0.9797	0.1593	0.8267	0.053*	
C8	0.7813 (3)	0.15549 (15)	0.7387 (3)	0.0377 (8)	
H8	0.7761	0.1125	0.7361	0.045*	
C9	0.9021 (4)	0.03945 (15)	0.2138 (4)	0.0412 (8)	
C10	0.8546 (3)	0.16613 (14)	0.3929 (3)	0.0307 (7)	
C11	0.9288 (3)	0.22405 (13)	0.4381 (3)	0.0288 (7)	
C12	1.0660 (3)	0.22665 (13)	0.5004 (3)	0.0303 (7)	
C13	1.1241 (4)	0.28340 (14)	0.5386 (3)	0.0370 (8)	
H13	1.2142	0.2849	0.5826	0.044*	
C14	1.0501 (4)	0.33711 (15)	0.5122 (3)	0.0425 (9)	
H14	1.0910	0.3749	0.5365	0.051*	

## supplementary materials

---

C15	0.9147 (4)	0.33536 (15)	0.4495 (3)	0.0431 (9)
H15	0.8642	0.3719	0.4314	0.052*
C16	0.8552 (4)	0.27936 (14)	0.4143 (3)	0.0373 (8)
H16	0.7633	0.2784	0.3733	0.045*

### Atomic displacement parameters ( $\text{\AA}^2$ )

	$U^{11}$	$U^{22}$	$U^{33}$	$U^{12}$	$U^{13}$	$U^{23}$
N1	0.0370 (17)	0.0357 (16)	0.0475 (18)	-0.0102 (13)	-0.0024 (14)	-0.0063 (14)
N2	0.0284 (16)	0.0278 (15)	0.0613 (19)	-0.0024 (12)	-0.0136 (13)	-0.0030 (13)
N3	0.0396 (18)	0.0449 (17)	0.0485 (18)	-0.0078 (14)	-0.0004 (15)	-0.0042 (14)
N4	0.0388 (17)	0.0296 (15)	0.0514 (18)	-0.0092 (12)	0.0036 (14)	-0.0057 (13)
N5	0.0256 (15)	0.0272 (14)	0.0558 (18)	-0.0054 (12)	-0.0035 (13)	-0.0093 (13)
N6	0.078 (2)	0.0351 (17)	0.0453 (18)	-0.0171 (16)	0.0066 (16)	-0.0064 (15)
O1	0.0281 (13)	0.0323 (12)	0.0555 (14)	0.0039 (10)	-0.0091 (11)	-0.0004 (11)
O2	0.0432 (15)	0.0294 (12)	0.0596 (15)	0.0017 (10)	-0.0031 (12)	-0.0032 (11)
O3	0.0241 (13)	0.0436 (14)	0.0639 (16)	-0.0005 (10)	-0.0073 (11)	-0.0033 (12)
O4	0.0315 (13)	0.0331 (13)	0.0577 (15)	0.0027 (10)	-0.0120 (11)	-0.0027 (11)
O5	0.299 (7)	0.114 (3)	0.102 (3)	-0.017 (4)	0.030 (4)	-0.006 (3)
S1	0.0350 (5)	0.0327 (5)	0.0567 (6)	-0.0036 (4)	-0.0089 (4)	-0.0056 (4)
S2	0.1012 (9)	0.0420 (6)	0.0677 (7)	-0.0309 (6)	0.0138 (6)	-0.0172 (5)
C1	0.0309 (19)	0.0198 (16)	0.050 (2)	0.0055 (13)	-0.0033 (16)	-0.0016 (15)
C2	0.0303 (18)	0.0316 (18)	0.0302 (17)	-0.0006 (15)	0.0035 (14)	0.0010 (14)
C3	0.0264 (17)	0.0303 (17)	0.0293 (17)	-0.0022 (13)	0.0012 (13)	-0.0030 (14)
C4	0.0308 (19)	0.0325 (18)	0.0313 (18)	0.0019 (14)	0.0051 (14)	-0.0033 (15)
C5	0.041 (2)	0.0376 (19)	0.043 (2)	-0.0130 (16)	0.0075 (17)	-0.0115 (16)
C6	0.030 (2)	0.064 (3)	0.040 (2)	-0.0169 (18)	0.0041 (16)	-0.0147 (19)
C7	0.030 (2)	0.058 (2)	0.045 (2)	-0.0004 (17)	-0.0064 (16)	-0.0021 (18)
C8	0.036 (2)	0.0357 (19)	0.0410 (19)	0.0022 (15)	-0.0036 (16)	-0.0021 (16)
C9	0.040 (2)	0.0313 (19)	0.052 (2)	-0.0035 (15)	-0.0020 (17)	-0.0036 (17)
C10	0.0270 (19)	0.0315 (18)	0.0337 (18)	-0.0003 (14)	0.0046 (14)	0.0012 (14)
C11	0.0253 (17)	0.0284 (17)	0.0328 (17)	-0.0003 (13)	0.0068 (14)	-0.0011 (14)
C12	0.0288 (18)	0.0285 (17)	0.0338 (18)	0.0000 (14)	0.0034 (14)	-0.0009 (14)
C13	0.0326 (19)	0.0354 (19)	0.043 (2)	-0.0050 (15)	0.0012 (16)	-0.0043 (16)
C14	0.048 (2)	0.0285 (19)	0.051 (2)	-0.0053 (17)	0.0093 (18)	-0.0096 (16)
C15	0.046 (2)	0.0274 (18)	0.056 (2)	0.0080 (16)	0.0064 (18)	0.0025 (16)
C16	0.034 (2)	0.0360 (19)	0.042 (2)	0.0057 (15)	0.0076 (15)	0.0017 (16)

### Geometric parameters ( $\text{\AA}$ , $^\circ$ )

N1—C1	1.338 (4)	S1—C1	1.697 (3)
N1—N2	1.379 (3)	S2—C9	1.694 (3)
N1—H1	0.8600	C2—C3	1.460 (4)
N2—C2	1.344 (4)	C3—C8	1.399 (4)
N2—H2	0.8600	C3—C4	1.402 (4)
N3—C1	1.319 (4)	C4—C5	1.391 (4)
N3—H3A	0.8600	C5—C6	1.369 (5)
N3—H3B	0.8600	C5—H5A	0.9300
N4—C9	1.340 (4)	C6—C7	1.378 (5)

N4—N5	1.385 (3)	C6—H6	0.9300
N4—H4'	0.8600	C7—C8	1.370 (4)
N5—C10	1.345 (4)	C7—H7	0.9300
N5—H5	0.8600	C8—H8	0.9300
N6—C9	1.308 (4)	C10—C11	1.492 (4)
N6—H6A	0.8600	C11—C16	1.390 (4)
N6—H6B	0.8600	C11—C12	1.394 (4)
O1—C2	1.237 (3)	C12—C13	1.389 (4)
O2—C4	1.351 (4)	C13—C14	1.367 (4)
O2—H2A	0.8200	C13—H13	0.9300
O3—C10	1.225 (3)	C14—C15	1.381 (5)
O4—C12	1.359 (3)	C14—H14	0.9300
O4—H4	0.8200	C15—C16	1.371 (4)
O5—H5D	0.8496	C15—H15	0.9300
O5—H5B	0.8500	C16—H16	0.9300
C1—N1—N2	122.6 (3)	C4—C5—H5A	120.0
C1—N1—H1	118.7	C5—C6—C7	120.9 (3)
N2—N1—H1	118.7	C5—C6—H6	119.5
C2—N2—N1	120.9 (3)	C7—C6—H6	119.5
C2—N2—H2	119.5	C8—C7—C6	119.5 (3)
N1—N2—H2	119.5	C8—C7—H7	120.3
C1—N3—H3A	120.0	C6—C7—H7	120.3
C1—N3—H3B	120.0	C7—C8—C3	121.4 (3)
H3A—N3—H3B	120.0	C7—C8—H8	119.3
C9—N4—N5	121.4 (3)	C3—C8—H8	119.3
C9—N4—H4'	119.3	N6—C9—N4	118.4 (3)
N5—N4—H4'	119.3	N6—C9—S2	123.2 (3)
C10—N5—N4	120.5 (3)	N4—C9—S2	118.4 (3)
C10—N5—H5	119.7	O3—C10—N5	121.3 (3)
N4—N5—H5	119.7	O3—C10—C11	121.8 (3)
C9—N6—H6A	120.0	N5—C10—C11	116.8 (3)
C9—N6—H6B	120.0	C16—C11—C12	118.1 (3)
H6A—N6—H6B	120.0	C16—C11—C10	116.9 (3)
C4—O2—H2A	109.5	C12—C11—C10	125.0 (3)
C12—O4—H4	109.5	O4—C12—C13	121.3 (3)
H5D—O5—H5B	129.4	O4—C12—C11	118.8 (3)
N3—C1—N1	119.1 (3)	C13—C12—C11	119.9 (3)
N3—C1—S1	122.3 (3)	C14—C13—C12	120.6 (3)
N1—C1—S1	118.6 (3)	C14—C13—H13	119.7
O1—C2—N2	119.4 (3)	C12—C13—H13	119.7
O1—C2—C3	122.4 (3)	C13—C14—C15	120.2 (3)
N2—C2—C3	118.2 (3)	C13—C14—H14	119.9
C8—C3—C4	118.1 (3)	C15—C14—H14	119.9
C8—C3—C2	123.4 (3)	C16—C15—C14	119.5 (3)
C4—C3—C2	118.5 (3)	C16—C15—H15	120.3
O2—C4—C5	117.2 (3)	C14—C15—H15	120.3
O2—C4—C3	122.7 (3)	C15—C16—C11	121.7 (3)
C5—C4—C3	120.0 (3)	C15—C16—H16	119.2
C6—C5—C4	120.1 (3)	C11—C16—H16	119.2

## supplementary materials

---

C6—C5—H5A

120.0

### Hydrogen-bond geometry (Å, °)

<i>D</i> —H $\cdots$ <i>A</i>	<i>D</i> —H	H $\cdots$ <i>A</i>	<i>D</i> $\cdots$ <i>A</i>	<i>D</i> —H $\cdots$ <i>A</i>
N2—H2 $\cdots$ S1 <sup>i</sup>	0.86	2.58	3.361 (3)	151
N5—H5 $\cdots$ S1 <sup>ii</sup>	0.86	2.83	3.404 (3)	125
N6—H6A $\cdots$ O2 <sup>iii</sup>	0.86	2.16	2.984 (3)	160
N6—H6B $\cdots$ S2 <sup>iv</sup>	0.86	2.85	3.647 (3)	156
O4—H4 $\cdots$ O1 <sup>ii</sup>	0.82	1.93	2.724 (3)	162
O5—H5B $\cdots$ O5 <sup>v</sup>	0.85	1.95	2.728 (10)	152
N3—H3A $\cdots$ O3	0.86	2.12	2.918 (3)	154
N3—H3B $\cdots$ O5	0.86	2.30	3.110 (5)	156
N5—H5 $\cdots$ O4	0.86	1.99	2.629 (3)	130
O2—H2A $\cdots$ O1	0.82	1.83	2.552 (3)	146
O5—H5D $\cdots$ S1	0.85	2.66	3.398 (4)	147

Symmetry codes: (i)  $-x+1, -y, -z+1$ ; (ii)  $x+1, y, z$ ; (iii)  $x+1/2, -y+1/2, z-1/2$ ; (iv)  $-x+2, -y, -z$ ; (v)  $-x+1, -y, -z$ .



Fig. 1

