

N-(2,3-Dichlorophenyl)-4-methylbenzenesulfonamide

K. Shakuntala,^a Sabine Foro,^b B. Thimme Gowda,^{a*}
P. G. Nirmala^a and Hartmut Fuess^b

^aDepartment of Chemistry, Mangalore University, Mangalagangothri 574 199, Mangalore, India, and ^bInstitute of Materials Science, Darmstadt University of Technology, Petersenstrasse 23, D-64287 Darmstadt, Germany
Correspondence e-mail: gowdabt@yahoo.com


Received 29 October 2010; accepted 29 October 2010

Key indicators: single-crystal X-ray study; $T = 299$ K; mean $\sigma(\text{C}-\text{C}) = 0.004$ Å; R factor = 0.065; wR factor = 0.176; data-to-parameter ratio = 13.9.

The title compound, $\text{C}_{13}\text{H}_{11}\text{Cl}_2\text{NO}_2\text{S}$, contains two molecules in the asymmetric unit in which the dihedral angles between the benzene rings are 76.0 (1) and 79.9 (1)°. The conformations of the N—H bonds with respect to their adjacent ortho-chlorine atoms are syn. In the crystal, N—H...O hydrogen bonds link the molecules into dimers.

Related literature

For our study of the effect of substituents on the structures of *N*-(aryl)arylsulfonamides, see: Gowda *et al.* (2009, 2010*a,b*). For related structures, see: Gelbrich *et al.* (2007); Perlovich *et al.* (2006).



Experimental

Crystal data

$\text{C}_{13}\text{H}_{11}\text{Cl}_2\text{NO}_2\text{S}$
 $M_r = 316.19$
Monoclinic, $P2_1/n$
 $a = 14.035$ (3) Å
 $b = 12.386$ (2) Å

$c = 15.993$ (3) Å
 $\beta = 98.30$ (1)°
 $V = 2751.1$ (9) Å³
 $Z = 8$
Cu $K\alpha$ radiation

$\mu = 5.64$ mm⁻¹
 $T = 299$ K

$0.50 \times 0.45 \times 0.40$ mm

Data collection

Enraf–Nonius CAD-4
diffractometer
9784 measured reflections
4903 independent reflections

4359 reflections with $I > 2\sigma(I)$
 $R_{\text{int}} = 0.131$
3 standard reflections every 120 min
intensity decay: 1.0%

Refinement

$R[F^2 > 2\sigma(F^2)] = 0.065$
 $wR(F^2) = 0.176$
 $S = 1.11$
4903 reflections
352 parameters
2 restraints

H atoms treated by a mixture of independent and constrained refinement
 $\Delta\rho_{\text{max}} = 0.81$ e Å⁻³
 $\Delta\rho_{\text{min}} = -0.56$ e Å⁻³

Table 1

Hydrogen-bond geometry (Å, °).

$D-H\cdots A$	$D-H$	$H\cdots A$	$D\cdots A$	$D-H\cdots A$
$\text{N1}-\text{H1N}\cdots\text{O3}$	0.84 (2)	2.29 (2)	3.086 (3)	160 (3)
$\text{N2}-\text{H2N}\cdots\text{O2}$	0.85 (2)	2.32 (2)	3.104 (3)	154 (3)

Data collection: *CAD-4-PC* (Enraf–Nonius, 1996); cell refinement: *CAD-4-PC*; data reduction: *REDU4* (Stoe & Cie, 1987); program(s) used to solve structure: *SHELXS97* (Sheldrick, 2008); program(s) used to refine structure: *SHELXL97* (Sheldrick, 2008); molecular graphics: *PLATON* (Spek, 2009); software used to prepare material for publication: *SHELXL97*.

KS thanks the University Grants Commission, Government of India, New Delhi for the award of a research fellowship under its faculty improvement program.

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: BQ2248).

References

- Enraf–Nonius (1996). *CAD-4-PC*. Enraf–Nonius, Delft, The Netherlands.
Gelbrich, T., Hursthouse, M. B. & Threlfall, T. L. (2007). *Acta Cryst.* **B63**, 621–632.
Gowda, B. T., Foro, S., Nirmala, P. G. & Fuess, H. (2009). *Acta Cryst.* **E65**, o2334.
Gowda, B. T., Foro, S., Nirmala, P. G. & Fuess, H. (2010*a*). *Acta Cryst.* **E66**, o188.
Gowda, B. T., Foro, S., Nirmala, P. G. & Fuess, H. (2010*b*). *Acta Cryst.* **E66**, o278.
Perlovich, G. L., Tkachev, V. V., Schaper, K.-J. & Raevsky, O. A. (2006). *Acta Cryst.* **E62**, o780–o782.
Sheldrick, G. M. (2008). *Acta Cryst.* **A64**, 112–122.
Spek, A. L. (2009). *Acta Cryst.* **D65**, 148–155.
Stoe & Cie (1987). *REDU4*. Stoe & Cie GmbH, Darmstadt, Germany.

supplementary materials

Acta Cryst. (2010). E66, o3062 [doi:10.1107/S1600536810044375]

***N*-(2,3-Dichlorophenyl)-4-methylbenzenesulfonamide**

K. Shakuntala, S. Foro, B. T. Gowda, P. G. Nirmala and H. Fues

Comment

As part of a study of the substituent effects on the crystal structures of *N*-(aryl)-arylsulfonamides (Gowda *et al.*, 2009; 2010*a*, *b*), in the present work, the structure of 4-methyl-*N*-(2,3-dichlorophenyl)-benzenesulfonamide (I) has been determined. The asymmetric unit of (I) contains two independent molecules. In one of the molecules, conformation of the N—C bond in the C—SO₂—NH—C segment of the structure has *gauche* torsions with respect to the S=O bonds (Fig. 1). The molecules are bent at the S atoms with the C—SO₂—NH—C torsion angles of 65.4 (2) and -61.7 (2). The conformations of the N—H bonds and the *ortho*-chloro groups in the anilino benzene rings are *syn* to each other. The benzene rings in the title compound are tilted relative to each other by 76.0 (1)° (molecule 1) and 79.9 (1)° (molecule 2). The other bond parameters in (I) are similar to those observed in 4-methyl-*N*-(2-chlorophenyl)-benzenesulfonamide (Gowda *et al.*, 2010*a*), 4-methyl-*N*-(3-chlorophenyl)benzenesulfonamide (Gowda *et al.*, 2010*b*), 4-methyl-*N*-(3,5-dichlorophenyl)benzenesulfonamide (Gowda *et al.*, 2009) and other aryl sulfonamides (Perlovich *et al.*, 2006; Gelbrich *et al.*, 2007). The N—H···O hydrogen bonds (Table 1) pack the molecules into infinite chains in the direction of *a*-axis (Fig. 2).

Experimental

The solution of toluene (10 ml) in chloroform (40 ml) was treated dropwise with chlorosulfonic acid (25 ml) at 0 °C. After the initial evolution of hydrogen chloride subsided, the reaction mixture was brought to room temperature and poured into crushed ice in a beaker. The chloroform layer was separated, washed with cold water and allowed to evaporate slowly. The residual 4-methylbenzenesulfonylchloride was treated with 2,3-dichloroaniline in the stoichiometric ratio and boiled for ten minutes. The reaction mixture was then cooled to room temperature and added to ice cold water (100 ml). The resultant 4-methyl-*N*-(2,3-dichlorophenyl)benzenesulfonamide was filtered under suction and washed thoroughly with cold water. It was then recrystallized to constant melting point from dilute ethanol. The purity of the compound was checked and characterized by recording its infrared and NMR spectra. The single crystals used in X-ray diffraction studies were grown in ethanolic solution by slow evaporation at room temperature.

Refinement

The H atoms of the NH groups were located in a difference map and later restrained to N—H = 0.86 (2) %A. The other H atoms were positioned with idealized geometry using a riding model with C—H = 0.93–0.96 Å. All H atoms were refined with isotropic displacement parameters (set to 1.2 times of the U_{eq} of the parent atom).

Figures




Fig. 1. Molecular structure of (I), showing the atom labeling scheme and displacement ellipsoids are drawn at the 50% probability level.




Fig. 2. Molecular packing of (I) with hydrogen bonding shown as dashed lines.

N-(2,3-Dichlorophenyl)-4-methylbenzenesulfonamide

Crystal data

$C_{13}H_{11}Cl_2NO_2S$

$M_r = 316.19$

Monoclinic, $P2_1/n$

Hall symbol: $-P\ 2_1/n$

$a = 14.035\ (3)\ \text{\AA}$

$b = 12.386\ (2)\ \text{\AA}$

$c = 15.993\ (3)\ \text{\AA}$

$\beta = 98.30\ (1)^\circ$

$V = 2751.1\ (9)\ \text{\AA}^3$

$Z = 8$

$F(000) = 1296$

$D_x = 1.527\ \text{Mg m}^{-3}$

Cu $K\alpha$ radiation, $\lambda = 1.54180\ \text{\AA}$

Cell parameters from 25 reflections

$\theta = 5.6\text{--}19.2^\circ$

$\mu = 5.64\ \text{mm}^{-1}$

$T = 299\ \text{K}$

Prism, colourless

$0.50 \times 0.45 \times 0.40\ \text{mm}$

Data collection

Enraf–Nonius CAD-4
diffractometer

Radiation source: fine-focus sealed tube
graphite

$\omega/2\theta$ scans

9784 measured reflections

4903 independent reflections

4359 reflections with $I > 2\sigma(I)$

$R_{\text{int}} = 0.131$

$\theta_{\text{max}} = 67.0^\circ$, $\theta_{\text{min}} = 3.9^\circ$

$h = -16 \rightarrow 16$

$k = -14 \rightarrow 0$

$l = -19 \rightarrow 19$

3 standard reflections every 120 min

intensity decay: 1.0%

Refinement

Refinement on F^2

Least-squares matrix: full

$R[F^2 > 2\sigma(F^2)] = 0.065$

Secondary atom site location: difference Fourier map

Hydrogen site location: inferred from neighbouring sites

H atoms treated by a mixture of independent and constrained refinement

$wR(F^2) = 0.176$	$w = 1/[\sigma^2(F_o^2) + (0.0914P)^2 + 0.5192P]$
$S = 1.11$	where $P = (F_o^2 + 2F_c^2)/3$
4903 reflections	$(\Delta/\sigma)_{\max} < 0.001$
352 parameters	$\Delta\rho_{\max} = 0.81 \text{ e } \text{\AA}^{-3}$
2 restraints	$\Delta\rho_{\min} = -0.56 \text{ e } \text{\AA}^{-3}$
Primary atom site location: structure-invariant direct methods	Extinction correction: <i>SHELXL97</i> (Sheldrick, 2008), $F_c^* = kFc[1+0.001xFc^2\lambda^3/\sin(2\theta)]^{-1/4}$ Extinction coefficient: 0.0038 (4)

Special details

Geometry. All e.s.d.'s (except the e.s.d. in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell e.s.d.'s are taken into account individually in the estimation of e.s.d.'s in distances, angles and torsion angles; correlations between e.s.d.'s in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell e.s.d.'s is used for estimating e.s.d.'s involving l.s. planes.

Refinement. Refinement of F^2 against ALL reflections. The weighted R -factor wR and goodness of fit S are based on F^2 , conventional R -factors R are based on F , with F set to zero for negative F^2 . The threshold expression of $F^2 > \sigma(F^2)$ is used only for calculating R -factors(gt) *etc.* and is not relevant to the choice of reflections for refinement. R -factors based on F^2 are statistically about twice as large as those based on F , and R -factors based on ALL data will be even larger.

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (\AA^2)

	x	y	z	$U_{\text{iso}}^*/U_{\text{eq}}$
C1	0.88559 (18)	0.0982 (2)	0.42938 (14)	0.0312 (6)
C2	0.9254 (2)	0.0053 (2)	0.40037 (17)	0.0404 (6)
H2	0.8955	-0.0612	0.4038	0.048*
C3	1.0094 (2)	0.0132 (3)	0.36653 (18)	0.0473 (7)
H3	1.0361	-0.0489	0.3468	0.057*
C4	1.05552 (19)	0.1107 (3)	0.36089 (16)	0.0412 (7)
C5	1.0132 (2)	0.2032 (3)	0.38929 (19)	0.0453 (7)
H5	1.0423	0.2700	0.3849	0.054*
C6	0.9292 (2)	0.1972 (2)	0.42354 (18)	0.0413 (6)
H6	0.9020	0.2592	0.4427	0.050*
C7	0.66060 (17)	0.0946 (2)	0.32818 (17)	0.0353 (6)
C8	0.64433 (18)	0.1597 (2)	0.25643 (17)	0.0375 (6)
C9	0.6067 (2)	0.1151 (3)	0.17875 (18)	0.0428 (7)
C10	0.5881 (2)	0.0062 (3)	0.1720 (2)	0.0578 (9)
H10	0.5641	-0.0237	0.1199	0.069*
C11	0.6048 (3)	-0.0574 (3)	0.2417 (3)	0.0637 (10)
H11	0.5925	-0.1311	0.2365	0.076*
C12	0.6398 (2)	-0.0151 (3)	0.3207 (2)	0.0530 (8)
H12	0.6493	-0.0597	0.3680	0.064*
C13	1.1486 (2)	0.1179 (4)	0.3250 (2)	0.0625 (10)
H13A	1.1354	0.1367	0.2662	0.075*
H13B	1.1809	0.0495	0.3311	0.075*
H13C	1.1888	0.1723	0.3548	0.075*

supplementary materials

N1	0.69333 (16)	0.1410 (2)	0.40784 (15)	0.0379 (5)
H1N	0.700 (3)	0.2081 (15)	0.410 (2)	0.045*
O1	0.75897 (16)	-0.02137 (18)	0.48773 (13)	0.0491 (5)
O2	0.78830 (16)	0.16207 (18)	0.54657 (12)	0.0470 (5)
Cl1	0.67147 (7)	0.29473 (6)	0.26491 (5)	0.0542 (3)
Cl2	0.58337 (8)	0.19562 (9)	0.09061 (5)	0.0690 (3)
S1	0.77998 (4)	0.08962 (6)	0.47581 (4)	0.0344 (2)
C14	0.62488 (18)	0.4300 (2)	0.57233 (14)	0.0302 (5)
C15	0.5861 (2)	0.3271 (2)	0.57427 (16)	0.0372 (6)
H15	0.6173	0.2680	0.5548	0.045*
C16	0.5007 (2)	0.3142 (2)	0.60547 (18)	0.0420 (6)
H16	0.4740	0.2455	0.6066	0.050*
C17	0.45287 (19)	0.4019 (3)	0.63566 (16)	0.0397 (6)
C18	0.4933 (2)	0.5035 (3)	0.63213 (18)	0.0437 (7)
H18	0.4619	0.5629	0.6510	0.052*
C19	0.5792 (2)	0.5186 (2)	0.60121 (17)	0.0383 (6)
H19	0.6059	0.5872	0.5998	0.046*
C20	0.83949 (17)	0.4415 (2)	0.68958 (16)	0.0322 (5)
C21	0.87507 (17)	0.3732 (2)	0.75535 (16)	0.0318 (5)
C22	0.89966 (18)	0.4133 (2)	0.83721 (17)	0.0375 (6)
C23	0.8875 (2)	0.5210 (3)	0.8543 (2)	0.0484 (7)
H23	0.9037	0.5475	0.9089	0.058*
C24	0.8515 (2)	0.5880 (3)	0.7900 (2)	0.0509 (7)
H24	0.8421	0.6605	0.8014	0.061*
C25	0.8284 (2)	0.5505 (2)	0.70714 (19)	0.0440 (7)
H25	0.8056	0.5982	0.6639	0.053*
C26	0.3602 (2)	0.3853 (3)	0.6705 (2)	0.0560 (9)
H26A	0.3739	0.3697	0.7298	0.067*
H26B	0.3257	0.3259	0.6418	0.067*
H26C	0.3218	0.4495	0.6620	0.067*
N2	0.81882 (17)	0.39956 (19)	0.60644 (14)	0.0370 (5)
H2N	0.826 (2)	0.3316 (16)	0.603 (2)	0.044*
O3	0.73700 (16)	0.37678 (18)	0.46284 (11)	0.0475 (5)
O4	0.75082 (16)	0.55963 (18)	0.52425 (13)	0.0480 (5)
Cl3	0.88652 (6)	0.23746 (6)	0.73614 (5)	0.0526 (3)
Cl4	0.94255 (7)	0.32756 (8)	0.91866 (5)	0.0600 (3)
S2	0.73465 (5)	0.44706 (5)	0.53342 (4)	0.0342 (2)

Atomic displacement parameters (\AA^2)

	U^{11}	U^{22}	U^{33}	U^{12}	U^{13}	U^{23}
C1	0.0288 (12)	0.0371 (14)	0.0268 (11)	0.0076 (10)	0.0013 (10)	-0.0017 (10)
C2	0.0439 (15)	0.0369 (14)	0.0404 (13)	0.0106 (12)	0.0061 (12)	-0.0029 (11)
C3	0.0477 (16)	0.0519 (17)	0.0430 (14)	0.0209 (15)	0.0094 (13)	-0.0032 (13)
C4	0.0297 (13)	0.0630 (19)	0.0302 (12)	0.0149 (13)	0.0021 (10)	0.0059 (12)
C5	0.0375 (15)	0.0479 (16)	0.0507 (16)	0.0007 (13)	0.0070 (13)	0.0035 (13)
C6	0.0379 (14)	0.0394 (15)	0.0475 (14)	0.0070 (12)	0.0089 (12)	-0.0044 (12)
C7	0.0203 (11)	0.0388 (14)	0.0458 (14)	0.0026 (10)	0.0010 (10)	-0.0021 (11)

C8	0.0250 (11)	0.0403 (14)	0.0472 (14)	0.0050 (11)	0.0055 (11)	-0.0079 (12)
C9	0.0314 (13)	0.0523 (17)	0.0436 (14)	0.0073 (13)	0.0021 (11)	-0.0059 (13)
C10	0.0478 (17)	0.056 (2)	0.0638 (19)	0.0036 (16)	-0.0109 (16)	-0.0218 (17)
C11	0.060 (2)	0.0422 (17)	0.080 (2)	-0.0048 (16)	-0.0183 (19)	-0.0116 (17)
C12	0.0467 (17)	0.0407 (16)	0.067 (2)	-0.0051 (14)	-0.0069 (16)	0.0032 (15)
C13	0.0407 (17)	0.099 (3)	0.0501 (17)	0.0165 (18)	0.0151 (14)	0.0137 (19)
N1	0.0304 (11)	0.0368 (12)	0.0465 (12)	0.0050 (10)	0.0062 (10)	-0.0011 (10)
O1	0.0523 (12)	0.0440 (12)	0.0517 (11)	-0.0005 (10)	0.0096 (10)	0.0120 (9)
O2	0.0531 (12)	0.0547 (12)	0.0342 (9)	0.0095 (10)	0.0102 (9)	-0.0046 (9)
Cl1	0.0739 (6)	0.0381 (4)	0.0496 (4)	0.0011 (3)	0.0049 (4)	0.0001 (3)
Cl2	0.0888 (7)	0.0725 (6)	0.0424 (4)	0.0130 (5)	-0.0019 (4)	0.0002 (4)
S1	0.0341 (4)	0.0386 (4)	0.0314 (3)	0.0055 (3)	0.0076 (3)	0.0024 (2)
C14	0.0315 (12)	0.0331 (12)	0.0247 (10)	0.0062 (10)	-0.0002 (10)	0.0018 (9)
C15	0.0435 (14)	0.0300 (12)	0.0386 (12)	0.0062 (12)	0.0076 (12)	-0.0001 (11)
C16	0.0411 (15)	0.0397 (15)	0.0445 (14)	-0.0033 (13)	0.0034 (12)	0.0034 (12)
C17	0.0295 (13)	0.0564 (17)	0.0308 (12)	0.0052 (12)	-0.0035 (10)	0.0041 (12)
C18	0.0374 (14)	0.0464 (16)	0.0472 (15)	0.0149 (13)	0.0062 (12)	-0.0023 (13)
C19	0.0418 (14)	0.0304 (13)	0.0410 (13)	0.0066 (12)	0.0005 (12)	-0.0027 (11)
C20	0.0214 (11)	0.0346 (13)	0.0403 (13)	-0.0015 (10)	0.0036 (10)	-0.0017 (11)
C21	0.0220 (11)	0.0311 (12)	0.0424 (13)	0.0003 (10)	0.0047 (10)	-0.0023 (11)
C22	0.0259 (11)	0.0449 (15)	0.0406 (13)	-0.0018 (11)	0.0007 (10)	0.0028 (12)
C23	0.0416 (15)	0.0538 (18)	0.0476 (15)	-0.0032 (14)	-0.0011 (13)	-0.0150 (14)
C24	0.0511 (17)	0.0367 (15)	0.0613 (18)	0.0000 (14)	-0.0044 (15)	-0.0127 (14)
C25	0.0441 (15)	0.0334 (14)	0.0509 (15)	0.0025 (12)	-0.0055 (13)	0.0003 (12)
C26	0.0375 (15)	0.083 (3)	0.0479 (16)	0.0029 (16)	0.0082 (13)	0.0082 (17)
N2	0.0370 (12)	0.0364 (12)	0.0379 (11)	0.0087 (10)	0.0061 (10)	-0.0006 (9)
O3	0.0581 (13)	0.0548 (12)	0.0315 (9)	0.0080 (11)	0.0128 (9)	-0.0008 (9)
O4	0.0569 (12)	0.0404 (11)	0.0480 (11)	-0.0015 (10)	0.0121 (10)	0.0119 (9)
Cl3	0.0709 (5)	0.0328 (4)	0.0515 (4)	0.0090 (3)	0.0001 (4)	0.0004 (3)
Cl4	0.0666 (5)	0.0630 (5)	0.0455 (4)	0.0009 (4)	-0.0081 (4)	0.0085 (4)
S2	0.0375 (4)	0.0354 (4)	0.0306 (3)	0.0034 (3)	0.0077 (3)	0.0043 (2)

Geometric parameters (Å, °)

C1—C6	1.380 (4)	C14—C19	1.384 (4)
C1—C2	1.387 (4)	C14—C15	1.388 (4)
C1—S1	1.754 (2)	C14—S2	1.756 (2)
C2—C3	1.370 (4)	C15—C16	1.373 (4)
C2—H2	0.9300	C15—H15	0.9300
C3—C4	1.380 (5)	C16—C17	1.399 (4)
C3—H3	0.9300	C16—H16	0.9300
C4—C5	1.396 (4)	C17—C18	1.385 (4)
C4—C13	1.503 (4)	C17—C26	1.500 (4)
C5—C6	1.371 (4)	C18—C19	1.379 (4)
C5—H5	0.9300	C18—H18	0.9300
C6—H6	0.9300	C19—H19	0.9300
C7—C12	1.392 (4)	C20—C21	1.386 (4)
C7—C8	1.394 (4)	C20—C25	1.392 (4)
C7—N1	1.413 (4)	C20—N2	1.418 (3)

supplementary materials

C8—C9	1.392 (4)	C21—C22	1.396 (4)
C8—C11	1.716 (3)	C21—C13	1.721 (3)
C9—C10	1.375 (5)	C22—C23	1.378 (4)
C9—C12	1.719 (3)	C22—C14	1.720 (3)
C10—C11	1.358 (6)	C23—C24	1.361 (5)
C10—H10	0.9300	C23—H23	0.9300
C11—C12	1.390 (5)	C24—C25	1.397 (4)
C11—H11	0.9300	C24—H24	0.9300
C12—H12	0.9300	C25—H25	0.9300
C13—H13A	0.9600	C26—H26A	0.9600
C13—H13B	0.9600	C26—H26B	0.9600
C13—H13C	0.9600	C26—H26C	0.9600
N1—S1	1.638 (2)	N2—S2	1.645 (2)
N1—H1N	0.837 (18)	N2—H2N	0.851 (18)
O1—S1	1.425 (2)	O3—S2	1.430 (2)
O2—S1	1.436 (2)	O4—S2	1.424 (2)
C6—C1—C2	120.7 (2)	C19—C14—C15	121.2 (2)
C6—C1—S1	119.5 (2)	C19—C14—S2	119.7 (2)
C2—C1—S1	119.8 (2)	C15—C14—S2	119.1 (2)
C3—C2—C1	118.9 (3)	C16—C15—C14	118.8 (2)
C3—C2—H2	120.5	C16—C15—H15	120.6
C1—C2—H2	120.5	C14—C15—H15	120.6
C2—C3—C4	121.8 (3)	C15—C16—C17	121.5 (3)
C2—C3—H3	119.1	C15—C16—H16	119.3
C4—C3—H3	119.1	C17—C16—H16	119.3
C3—C4—C5	118.1 (3)	C18—C17—C16	118.1 (3)
C3—C4—C13	121.2 (3)	C18—C17—C26	121.5 (3)
C5—C4—C13	120.6 (3)	C16—C17—C26	120.4 (3)
C6—C5—C4	121.1 (3)	C19—C18—C17	121.5 (3)
C6—C5—H5	119.5	C19—C18—H18	119.3
C4—C5—H5	119.5	C17—C18—H18	119.3
C5—C6—C1	119.4 (3)	C18—C19—C14	118.9 (3)
C5—C6—H6	120.3	C18—C19—H19	120.6
C1—C6—H6	120.3	C14—C19—H19	120.6
C12—C7—C8	119.1 (3)	C21—C20—C25	118.5 (2)
C12—C7—N1	120.8 (3)	C21—C20—N2	119.3 (2)
C8—C7—N1	120.0 (3)	C25—C20—N2	122.1 (2)
C9—C8—C7	120.0 (3)	C20—C21—C22	120.5 (2)
C9—C8—C11	120.6 (2)	C20—C21—C13	119.61 (19)
C7—C8—C11	119.4 (2)	C22—C21—C13	119.9 (2)
C10—C9—C8	120.2 (3)	C23—C22—C21	120.6 (3)
C10—C9—C12	119.5 (2)	C23—C22—C14	119.1 (2)
C8—C9—C12	120.3 (2)	C21—C22—C14	120.2 (2)
C11—C10—C9	119.8 (3)	C24—C23—C22	118.9 (3)
C11—C10—H10	120.1	C24—C23—H23	120.5
C9—C10—H10	120.1	C22—C23—H23	120.5
C10—C11—C12	121.6 (3)	C23—C24—C25	121.7 (3)
C10—C11—H11	119.2	C23—C24—H24	119.2
C12—C11—H11	119.2	C25—C24—H24	119.2

C11—C12—C7	119.2 (3)	C20—C25—C24	119.7 (3)
C11—C12—H12	120.4	C20—C25—H25	120.1
C7—C12—H12	120.4	C24—C25—H25	120.1
C4—C13—H13A	109.5	C17—C26—H26A	109.5
C4—C13—H13B	109.5	C17—C26—H26B	109.5
H13A—C13—H13B	109.5	H26A—C26—H26B	109.5
C4—C13—H13C	109.5	C17—C26—H26C	109.5
H13A—C13—H13C	109.5	H26A—C26—H26C	109.5
H13B—C13—H13C	109.5	H26B—C26—H26C	109.5
C7—N1—S1	123.66 (19)	C20—N2—S2	124.27 (18)
C7—N1—H1N	117 (2)	C20—N2—H2N	114 (2)
S1—N1—H1N	107 (2)	S2—N2—H2N	113 (2)
O1—S1—O2	119.51 (13)	O4—S2—O3	119.53 (12)
O1—S1—N1	108.38 (13)	O4—S2—N2	108.34 (13)
O2—S1—N1	104.38 (12)	O3—S2—N2	104.28 (12)
O1—S1—C1	108.59 (13)	O4—S2—C14	108.40 (13)
O2—S1—C1	108.57 (12)	O3—S2—C14	109.14 (13)
N1—S1—C1	106.70 (12)	N2—S2—C14	106.39 (11)
C6—C1—C2—C3	0.7 (4)	C19—C14—C15—C16	0.1 (4)
S1—C1—C2—C3	-178.1 (2)	S2—C14—C15—C16	179.4 (2)
C1—C2—C3—C4	0.3 (4)	C14—C15—C16—C17	-0.5 (4)
C2—C3—C4—C5	-1.3 (4)	C15—C16—C17—C18	1.0 (4)
C2—C3—C4—C13	178.9 (3)	C15—C16—C17—C26	-179.2 (3)
C3—C4—C5—C6	1.4 (4)	C16—C17—C18—C19	-1.1 (4)
C13—C4—C5—C6	-178.8 (3)	C26—C17—C18—C19	179.1 (3)
C4—C5—C6—C1	-0.5 (4)	C17—C18—C19—C14	0.7 (4)
C2—C1—C6—C5	-0.5 (4)	C15—C14—C19—C18	-0.2 (4)
S1—C1—C6—C5	178.3 (2)	S2—C14—C19—C18	-179.5 (2)
C12—C7—C8—C9	0.8 (4)	C25—C20—C21—C22	-0.4 (4)
N1—C7—C8—C9	-176.2 (2)	N2—C20—C21—C22	177.4 (2)
C12—C7—C8—C11	-179.0 (2)	C25—C20—C21—C13	178.2 (2)
N1—C7—C8—C11	4.0 (3)	N2—C20—C21—C13	-4.0 (3)
C7—C8—C9—C10	-1.9 (4)	C20—C21—C22—C23	1.1 (4)
C11—C8—C9—C10	177.9 (2)	C13—C21—C22—C23	-177.6 (2)
C7—C8—C9—C12	178.22 (19)	C20—C21—C22—C14	179.30 (18)
C11—C8—C9—C12	-2.0 (3)	C13—C21—C22—C14	0.7 (3)
C8—C9—C10—C11	1.2 (5)	C21—C22—C23—C24	-0.3 (4)
C12—C9—C10—C11	-178.9 (3)	C14—C22—C23—C24	-178.5 (2)
C9—C10—C11—C12	0.5 (6)	C22—C23—C24—C25	-1.2 (5)
C10—C11—C12—C7	-1.7 (5)	C21—C20—C25—C24	-1.0 (4)
C8—C7—C12—C11	1.0 (4)	N2—C20—C25—C24	-178.8 (3)
N1—C7—C12—C11	177.9 (3)	C23—C24—C25—C20	1.9 (5)
C12—C7—N1—S1	47.9 (3)	C21—C20—N2—S2	150.1 (2)
C8—C7—N1—S1	-135.2 (2)	C25—C20—N2—S2	-32.2 (3)
C7—N1—S1—O1	-51.4 (2)	C20—N2—S2—O4	54.7 (2)
C7—N1—S1—O2	-179.8 (2)	C20—N2—S2—O3	-177.0 (2)
C7—N1—S1—C1	65.4 (2)	C20—N2—S2—C14	-61.7 (2)
C6—C1—S1—O1	-169.9 (2)	C19—C14—S2—O4	-10.5 (2)
C2—C1—S1—O1	8.9 (3)	C15—C14—S2—O4	170.1 (2)

supplementary materials

C6—C1—S1—O2	-38.5 (3)	C19—C14—S2—O3	-142.2 (2)
C2—C1—S1—O2	140.3 (2)	C15—C14—S2—O3	38.4 (2)
C6—C1—S1—N1	73.5 (2)	C19—C14—S2—N2	105.8 (2)
C2—C1—S1—N1	-107.8 (2)	C15—C14—S2—N2	-73.6 (2)

Hydrogen-bond geometry (Å, °)

<i>D—H...A</i>	<i>D—H</i>	<i>H...A</i>	<i>D...A</i>	<i>D—H...A</i>
N1—H1N...O3	0.84 (2)	2.29 (2)	3.086 (3)	160 (3)
N1—H1N...C11	0.84 (2)	2.54 (3)	2.957 (2)	112 (3)
N2—H2N...O2	0.85 (2)	2.32 (2)	3.104 (3)	154 (3)
N2—H2N...C13	0.85 (2)	2.46 (3)	2.944 (2)	117 (3)

Fig. 1




Fig. 2

