

1-[(3-Methylpiperidin-1-yl)(3-nitrophenyl)methyl]naphthalen-2-ol

Jin Mei Chen and Hong Zhao*

School of Chemistry and Chemical Engineering, Southeast University, Nanjing 210096, People's Republic of China

Correspondence e-mail: zhaozhong@seu.edu.cn


Received 3 November 2010; accepted 4 November 2010

Key indicators: single-crystal X-ray study; $T = 295\text{ K}$; mean $\sigma(\text{C}-\text{C}) = 0.005\text{ \AA}$; R factor = 0.083; wR factor = 0.174; data-to-parameter ratio = 15.2.

The title compound, $C_{23}H_{24}N_2O_3$, was synthesized from naphthalen-2-ol, 3-nitrobenzaldehyde and 3-methylpiperidine. The dihedral angles between the naphthalene system and the nitrobenzene and methylpiperidine rings are $78.53(13)$ and $64.14(15)^\circ$, respectively. The molecular conformation is stabilized by a strong intramolecular $O-\text{H}\cdots\text{N}$ hydrogen bond.

Related literature

For applications of naphthalen-2-ol derivatives in catalytic asymmetric synthesis, see: Szatmari & Fulop (2004). For related structures, see: Zhao & Sun (2005); Wang & Zhao (2009); Xiao & Zhao (2010);



Experimental

Crystal data

$C_{23}H_{24}N_2O_3$	$V = 3980(4)\text{ \AA}^3$
$M_r = 376.44$	$Z = 8$
Orthorhombic, $Pbca$	Mo $K\alpha$ radiation
$a = 11.980(2)\text{ \AA}$	$\mu = 0.08\text{ mm}^{-1}$
$b = 10.965(2)\text{ \AA}$	$T = 295\text{ K}$
$c = 30.30(3)\text{ \AA}$	$0.25 \times 0.22 \times 0.18\text{ mm}$

Data collection

Rigaku SCXmini diffractometer	34080 measured reflections
Absorption correction: multi-scan (<i>CrystalClear</i> ; Rigaku, 2005)	3877 independent reflections
$(\text{CrystalClear}; \text{Rigaku}, 2005)$	2409 reflections with $I > 2\sigma(I)$
$R_{\min} = 0.982$, $T_{\max} = 0.992$	$R_{\text{int}} = 0.110$

Refinement

$R[F^2 > 2\sigma(F^2)] = 0.083$	255 parameters
$wR(F^2) = 0.174$	H-atom parameters constrained
$S = 1.12$	$\Delta\rho_{\max} = 0.14\text{ e \AA}^{-3}$
3877 reflections	$\Delta\rho_{\min} = -0.15\text{ e \AA}^{-3}$

Table 1

Hydrogen-bond geometry (\AA , $^\circ$).

$D-\text{H}\cdots A$	$D-\text{H}$	$\text{H}\cdots A$	$D\cdots A$	$D-\text{H}\cdots A$
O1-H1 \cdots N1	0.82	1.86	2.579 (3)	147

Data collection: *CrystalClear* (Rigaku, 2005); cell refinement: *CrystalClear*; data reduction: *CrystalClear*; program(s) used to solve structure: *SHELXS97* (Sheldrick, 2008); program(s) used to refine structure: *SHELXL97* (Sheldrick, 2008); molecular graphics: *SHELXTL/PC* (Sheldrick, 2008); software used to prepare material for publication: *SHELXTL/PC*.

This work was supported financially by a Southeast University grant for young researchers (4007041027).

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: BX2327).

References

- Rigaku (2005). *CrystalClear*. Rigaku Corporation, Tokyo, Japan.
Sheldrick, G. M. (2008). *Acta Cryst. A* **64**, 112–122.
Szatmari, I. & Fulop, F. (2004). *Curr. Org. Synth.* **1**, 155–165.
Wang, W. X. & Zhao, H. (2009). *Acta Cryst. E* **65**, o1277.
Xiao, J. & Zhao, H. (2010). *Acta Cryst. E* **66**, o2938.
Zhao, B. & Sun, Y.-X. (2005). *Acta Cryst. E* **61**, m652–m653.

supplementary materials

Acta Cryst. (2010). E66, o3118 [doi:10.1107/S1600536810045149]

1-[(3-Methylpiperidin-1-yl)(3-nitrophenyl)methyl]naphthalen-2-ol

J. M. Chen and H. Zhao

Comment

Compounds derived from naphthalen-2-ol have been of great interest in organic chemistry due to their application in catalytic asymmetric synthesis (Szatmari & Fulop, 2004; Zhao & Sun, 2005). As an extension of our work on the structural characterization of naphthol compounds (Wang & Zhao, 2009; Xiao & Zhao, 2010), we report here the structure of (I). In the title compound (Fig. 1) bond lengths and angles have normal values. The dihedral angle between the naphthylen fragment with the nitrobenzene and methyl piperidine rings are 78.53 (13) and 64.14 (15) $^{\circ}$ respectively. The molecular conformation is stabilized by one strong intramolecular O—H \cdots N hydrogen bonding (Table 1).

Experimental

A dry 50 ml flask was charged with 3-nitrobenzaldehyde (10 mmol), naphthalen-2-ol (10 mmol) and 3-methylpiperidine (10 mmol). The mixture was stirred at 100°C for 12 h and then added ethanol (15 ml), after heated under reflux for 1 h, the precipitate was filtrated out and washed with ethanol for three times to give (I). Colourless crystals suitable for X-ray diffraction were obtained by slow evaporation of a dichloromethane solution.

Refinement

All H atoms were detected in a difference map, but were placed in calculated positions and refined using a riding motion approximation, with C—H=0.93–0.98 Å, with $U_{\text{iso}}(\text{H})=1.2U_{\text{eq}}(\text{C})$ and $U_{\text{iso}}(\text{Hmethyl})=1.2U_{\text{eq}}(\text{Cmethyl})$; O—H=0.82 Å, with $U_{\text{iso}}(\text{H})=1.5U_{\text{eq}}(\text{O})$.

Figures




Fig. 1. The molecular structure of the title compound, showing the atomic numbering scheme. The displacement ellipsoids are drawn at the 30% probability level.

1-[(3-Methylpiperidin-1-yl)(3-nitrophenyl)methyl]naphthalen-2-ol

Crystal data

C₂₃H₂₄N₂O₃

$M_r = 376.44$

$F(000) = 1600$

$D_x = 1.256 \text{ Mg m}^{-3}$

supplementary materials

Orthorhombic, <i>Pbca</i>	Mo $K\alpha$ radiation, $\lambda = 0.71073 \text{ \AA}$
Hall symbol: -P 2ac 2ab	Cell parameters from 4541 reflections
$a = 11.980(2) \text{ \AA}$	$\theta = 2.3\text{--}27.5^\circ$
$b = 10.965(2) \text{ \AA}$	$\mu = 0.08 \text{ mm}^{-1}$
$c = 30.30(3) \text{ \AA}$	$T = 295 \text{ K}$
$V = 3980(4) \text{ \AA}^3$	Prism, colourless
$Z = 8$	$0.25 \times 0.22 \times 0.18 \text{ mm}$

Data collection

Rigaku SCXmini diffractometer	3877 independent reflections
Radiation source: fine-focus sealed tube graphite	2409 reflections with $I > 2\sigma(I)$
Detector resolution: 13.6612 pixels mm^{-1}	$R_{\text{int}} = 0.110$
CCD_Profile_fitting scans	$\theta_{\text{max}} = 26.0^\circ, \theta_{\text{min}} = 2.6^\circ$
Absorption correction: multi-scan (<i>CrystalClear</i> ; Rigaku, 2005)	$h = -14 \rightarrow 14$
$T_{\text{min}} = 0.982, T_{\text{max}} = 0.992$	$k = -13 \rightarrow 13$
34080 measured reflections	$l = -37 \rightarrow 37$

Refinement

Refinement on F^2	Primary atom site location: structure-invariant direct methods
Least-squares matrix: full	Secondary atom site location: difference Fourier map
$R[F^2 > 2\sigma(F^2)] = 0.083$	Hydrogen site location: inferred from neighbouring sites
$wR(F^2) = 0.174$	H-atom parameters constrained
$S = 1.12$	$w = 1/[\sigma^2(F_o^2) + (0.0563P)^2 + 1.4193P]$ where $P = (F_o^2 + 2F_c^2)/3$
3877 reflections	$(\Delta/\sigma)_{\text{max}} < 0.001$
255 parameters	$\Delta\rho_{\text{max}} = 0.14 \text{ e \AA}^{-3}$
0 restraints	$\Delta\rho_{\text{min}} = -0.15 \text{ e \AA}^{-3}$

Special details

Geometry. All e.s.d.'s (except the e.s.d. in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell e.s.d.'s are taken into account individually in the estimation of e.s.d.'s in distances, angles and torsion angles; correlations between e.s.d.'s in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell e.s.d.'s is used for estimating e.s.d.'s involving l.s. planes.

Refinement. Refinement of F^2 against ALL reflections. The weighted R -factor wR and goodness of fit S are based on F^2 , conventional R -factors R are based on F , with F set to zero for negative F^2 . The threshold expression of $F^2 > \sigma(F^2)$ is used only for calculating R -factors(gt) etc. and is not relevant to the choice of reflections for refinement. R -factors based on F^2 are statistically about twice as large as those based on F , and R -factors based on ALL data will be even larger.

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (\AA^2)

	<i>x</i>	<i>y</i>	<i>z</i>	$U_{\text{iso}}^*/U_{\text{eq}}$
C1	0.7000 (2)	0.9558 (2)	0.65000 (10)	0.0433 (7)
C2	0.6194 (2)	1.0425 (3)	0.64171 (11)	0.0523 (8)
C3	0.5315 (3)	1.0634 (3)	0.67144 (13)	0.0659 (9)
H3	0.4769	1.1206	0.6646	0.079*
C4	0.5255 (3)	1.0015 (4)	0.70975 (13)	0.0707 (10)
H4	0.4661	1.0161	0.7288	0.085*
C5	0.6075 (3)	0.9147 (3)	0.72156 (11)	0.0584 (9)
C6	0.6961 (2)	0.8920 (3)	0.69134 (10)	0.0478 (7)
C7	0.7785 (3)	0.8061 (3)	0.70421 (10)	0.0552 (8)
H7	0.8381	0.7905	0.6854	0.066*
C8	0.7725 (3)	0.7459 (3)	0.74356 (11)	0.0708 (10)
H8	0.8279	0.6904	0.7513	0.085*
C9	0.6840 (3)	0.7671 (4)	0.77209 (12)	0.0820 (12)
H9	0.6796	0.7246	0.7986	0.098*
C10	0.6041 (3)	0.8493 (4)	0.76161 (12)	0.0774 (11)
H10	0.5457	0.8630	0.7812	0.093*
C11	0.7905 (2)	0.9244 (2)	0.61686 (9)	0.0422 (7)
H11	0.8033	0.8363	0.6189	0.051*
C12	0.6680 (3)	0.8576 (3)	0.55808 (11)	0.0599 (9)
H12A	0.6106	0.8529	0.5806	0.072*
H12B	0.7044	0.7787	0.5564	0.072*
C13	0.6143 (3)	0.8861 (3)	0.51400 (13)	0.0728 (11)
H13	0.5776	0.9657	0.5166	0.087*
C14	0.7021 (4)	0.8968 (4)	0.47882 (13)	0.0945 (14)
H14A	0.6680	0.9242	0.4515	0.113*
H14B	0.7353	0.8175	0.4736	0.113*
C15	0.7918 (3)	0.9860 (4)	0.49260 (11)	0.0828 (12)
H15A	0.8513	0.9851	0.4709	0.099*
H15B	0.7605	1.0676	0.4933	0.099*
C16	0.8395 (3)	0.9559 (3)	0.53738 (10)	0.0633 (9)
H16A	0.8776	0.8780	0.5361	0.076*
H16B	0.8936	1.0176	0.5457	0.076*
C17	0.5250 (4)	0.7925 (3)	0.50331 (16)	0.1061 (16)
H17A	0.4876	0.8153	0.4765	0.159*
H17B	0.4719	0.7892	0.5270	0.159*
H17C	0.5590	0.7139	0.4996	0.159*
C18	0.9014 (2)	0.9868 (3)	0.62676 (9)	0.0428 (7)
C19	1.0000 (2)	0.9224 (3)	0.62142 (9)	0.0472 (7)
H19	0.9988	0.8410	0.6128	0.057*
C20	1.0999 (3)	0.9808 (3)	0.62900 (10)	0.0553 (8)
C21	1.1060 (3)	1.1004 (4)	0.64166 (12)	0.0710 (10)
H21	1.1746	1.1379	0.6463	0.085*
C22	1.0079 (3)	1.1635 (3)	0.64731 (13)	0.0748 (10)
H22	1.0099	1.2448	0.6560	0.090*
C23	0.9067 (3)	1.1075 (3)	0.64020 (11)	0.0584 (8)

supplementary materials

H23	0.8410	1.1512	0.6445	0.070*
N1	0.7505 (2)	0.9505 (2)	0.57076 (7)	0.0469 (6)
N2	1.2039 (3)	0.9098 (4)	0.62304 (11)	0.0788 (9)
O1	0.61885 (19)	1.11174 (19)	0.60442 (8)	0.0645 (6)
H1	0.6631	1.0833	0.5865	0.097*
O2	1.1967 (2)	0.8052 (3)	0.61087 (11)	0.1051 (10)
O3	1.2920 (2)	0.9589 (3)	0.63113 (13)	0.1305 (14)

Atomic displacement parameters (\AA^2)

	U^{11}	U^{22}	U^{33}	U^{12}	U^{13}	U^{23}
C1	0.0363 (15)	0.0400 (15)	0.0536 (18)	-0.0022 (13)	-0.0002 (13)	-0.0087 (13)
C2	0.0470 (18)	0.0457 (17)	0.064 (2)	-0.0034 (15)	-0.0002 (15)	-0.0073 (15)
C3	0.047 (2)	0.066 (2)	0.085 (3)	0.0061 (17)	0.0059 (18)	-0.010 (2)
C4	0.048 (2)	0.084 (3)	0.079 (3)	-0.0049 (19)	0.0190 (18)	-0.019 (2)
C5	0.0454 (19)	0.068 (2)	0.062 (2)	-0.0110 (17)	0.0083 (16)	-0.0132 (17)
C6	0.0440 (17)	0.0445 (16)	0.0550 (19)	-0.0103 (14)	-0.0010 (14)	-0.0083 (14)
C7	0.057 (2)	0.0570 (19)	0.0515 (19)	-0.0080 (16)	0.0011 (15)	-0.0019 (15)
C8	0.072 (2)	0.086 (3)	0.054 (2)	-0.001 (2)	0.0013 (18)	0.0116 (19)
C9	0.080 (3)	0.109 (3)	0.057 (2)	-0.012 (3)	0.004 (2)	0.020 (2)
C10	0.064 (2)	0.110 (3)	0.058 (2)	-0.017 (2)	0.0178 (19)	-0.006 (2)
C11	0.0437 (17)	0.0387 (15)	0.0443 (16)	0.0001 (13)	-0.0026 (13)	-0.0012 (12)
C12	0.067 (2)	0.0439 (17)	0.069 (2)	-0.0014 (16)	-0.0200 (17)	-0.0041 (15)
C13	0.086 (3)	0.0478 (19)	0.085 (3)	0.0037 (19)	-0.040 (2)	-0.0030 (18)
C14	0.134 (4)	0.089 (3)	0.060 (3)	0.004 (3)	-0.028 (3)	-0.002 (2)
C15	0.105 (3)	0.090 (3)	0.053 (2)	-0.001 (2)	-0.009 (2)	0.006 (2)
C16	0.072 (2)	0.067 (2)	0.051 (2)	0.0067 (19)	-0.0002 (17)	-0.0001 (16)
C17	0.115 (4)	0.070 (3)	0.134 (4)	-0.009 (2)	-0.071 (3)	-0.004 (2)
C18	0.0425 (17)	0.0462 (17)	0.0397 (16)	-0.0016 (14)	-0.0008 (13)	-0.0007 (12)
C19	0.0503 (18)	0.0472 (17)	0.0441 (17)	0.0015 (15)	-0.0022 (14)	0.0004 (13)
C20	0.0444 (19)	0.069 (2)	0.0522 (19)	-0.0005 (17)	-0.0015 (15)	0.0060 (16)
C21	0.054 (2)	0.076 (2)	0.083 (3)	-0.022 (2)	-0.0081 (19)	-0.005 (2)
C22	0.071 (3)	0.058 (2)	0.095 (3)	-0.013 (2)	-0.002 (2)	-0.0178 (19)
C23	0.053 (2)	0.0474 (18)	0.075 (2)	-0.0015 (16)	0.0026 (17)	-0.0117 (16)
N1	0.0488 (14)	0.0447 (13)	0.0472 (14)	0.0003 (12)	-0.0068 (12)	-0.0026 (11)
N2	0.0454 (19)	0.101 (3)	0.090 (2)	0.009 (2)	-0.0059 (16)	0.011 (2)
O1	0.0670 (16)	0.0489 (13)	0.0775 (17)	0.0158 (11)	0.0021 (12)	0.0042 (12)
O2	0.0703 (19)	0.106 (2)	0.139 (3)	0.0348 (18)	-0.0131 (17)	-0.027 (2)
O3	0.0440 (17)	0.134 (3)	0.214 (4)	-0.0033 (18)	-0.020 (2)	0.008 (3)

Geometric parameters (\AA , $^\circ$)

C1—C2	1.379 (4)	C13—H13	0.9800
C1—C6	1.435 (4)	C14—C15	1.511 (5)
C1—C11	1.517 (4)	C14—H14A	0.9700
C2—O1	1.361 (4)	C14—H14B	0.9700
C2—C3	1.404 (4)	C15—C16	1.509 (5)
C3—C4	1.347 (5)	C15—H15A	0.9700
C3—H3	0.9300	C15—H15B	0.9700

C4—C5	1.413 (5)	C16—N1	1.471 (4)
C4—H4	0.9300	C16—H16A	0.9700
C5—C10	1.410 (5)	C16—H16B	0.9700
C5—C6	1.424 (4)	C17—H17A	0.9600
C6—C7	1.419 (4)	C17—H17B	0.9600
C7—C8	1.365 (4)	C17—H17C	0.9600
C7—H7	0.9300	C18—C19	1.386 (4)
C8—C9	1.387 (5)	C18—C23	1.387 (4)
C8—H8	0.9300	C19—C20	1.376 (4)
C9—C10	1.353 (5)	C19—H19	0.9300
C9—H9	0.9300	C20—C21	1.368 (5)
C10—H10	0.9300	C20—N2	1.481 (4)
C11—N1	1.505 (4)	C21—C22	1.375 (5)
C11—C18	1.524 (4)	C21—H21	0.9300
C11—H11	0.9800	C22—C23	1.376 (4)
C12—N1	1.471 (4)	C22—H22	0.9300
C12—C13	1.515 (5)	C23—H23	0.9300
C12—H12A	0.9700	N2—O2	1.207 (4)
C12—H12B	0.9700	N2—O3	1.210 (4)
C13—C14	1.502 (5)	O1—H1	0.8200
C13—C17	1.518 (5)		
C2—C1—C6	118.2 (3)	C13—C14—H14A	109.5
C2—C1—C11	122.5 (3)	C15—C14—H14A	109.5
C6—C1—C11	119.3 (2)	C13—C14—H14B	109.5
O1—C2—C1	122.6 (3)	C15—C14—H14B	109.5
O1—C2—C3	116.0 (3)	H14A—C14—H14B	108.1
C1—C2—C3	121.4 (3)	C16—C15—C14	112.1 (3)
C4—C3—C2	120.7 (3)	C16—C15—H15A	109.2
C4—C3—H3	119.6	C14—C15—H15A	109.2
C2—C3—H3	119.6	C16—C15—H15B	109.2
C3—C4—C5	121.3 (3)	C14—C15—H15B	109.2
C3—C4—H4	119.3	H15A—C15—H15B	107.9
C5—C4—H4	119.3	N1—C16—C15	110.6 (3)
C10—C5—C4	122.7 (3)	N1—C16—H16A	109.5
C10—C5—C6	119.0 (3)	C15—C16—H16A	109.5
C4—C5—C6	118.3 (3)	N1—C16—H16B	109.5
C7—C6—C5	117.3 (3)	C15—C16—H16B	109.5
C7—C6—C1	122.8 (3)	H16A—C16—H16B	108.1
C5—C6—C1	120.0 (3)	C13—C17—H17A	109.5
C8—C7—C6	121.7 (3)	C13—C17—H17B	109.5
C8—C7—H7	119.2	H17A—C17—H17B	109.5
C6—C7—H7	119.2	C13—C17—H17C	109.5
C7—C8—C9	120.2 (4)	H17A—C17—H17C	109.5
C7—C8—H8	119.9	H17B—C17—H17C	109.5
C9—C8—H8	119.9	C19—C18—C23	118.8 (3)
C10—C9—C8	120.4 (3)	C19—C18—C11	119.5 (3)
C10—C9—H9	119.8	C23—C18—C11	121.7 (3)
C8—C9—H9	119.8	C20—C19—C18	119.0 (3)
C9—C10—C5	121.4 (3)	C20—C19—H19	120.5

supplementary materials

C9—C10—H10	119.3	C18—C19—H19	120.5
C5—C10—H10	119.3	C21—C20—C19	122.6 (3)
N1—C11—C1	110.1 (2)	C21—C20—N2	119.6 (3)
N1—C11—C18	112.0 (2)	C19—C20—N2	117.8 (3)
C1—C11—C18	113.0 (2)	C20—C21—C22	118.1 (3)
N1—C11—H11	107.1	C20—C21—H21	120.9
C1—C11—H11	107.1	C22—C21—H21	120.9
C18—C11—H11	107.1	C21—C22—C23	120.6 (3)
N1—C12—C13	111.9 (3)	C21—C22—H22	119.7
N1—C12—H12A	109.2	C23—C22—H22	119.7
C13—C12—H12A	109.2	C22—C23—C18	120.8 (3)
N1—C12—H12B	109.2	C22—C23—H23	119.6
C13—C12—H12B	109.2	C18—C23—H23	119.6
H12A—C12—H12B	107.9	C12—N1—C16	109.6 (2)
C14—C13—C12	110.1 (3)	C12—N1—C11	109.0 (2)
C14—C13—C17	113.3 (4)	C16—N1—C11	114.5 (2)
C12—C13—C17	110.3 (3)	O2—N2—O3	123.2 (4)
C14—C13—H13	107.6	O2—N2—C20	118.5 (3)
C12—C13—H13	107.6	O3—N2—C20	118.3 (4)
C17—C13—H13	107.6	C2—O1—H1	109.5
C13—C14—C15	110.6 (3)		
C6—C1—C2—O1	176.9 (3)	C12—C13—C14—C15	−52.7 (4)
C11—C1—C2—O1	−4.2 (4)	C17—C13—C14—C15	−176.8 (3)
C6—C1—C2—C3	−3.8 (4)	C13—C14—C15—C16	53.0 (4)
C11—C1—C2—C3	175.1 (3)	C14—C15—C16—N1	−56.1 (4)
O1—C2—C3—C4	−178.7 (3)	N1—C11—C18—C19	−95.4 (3)
C1—C2—C3—C4	2.0 (5)	C1—C11—C18—C19	139.5 (3)
C2—C3—C4—C5	0.7 (5)	N1—C11—C18—C23	83.3 (3)
C3—C4—C5—C10	179.3 (3)	C1—C11—C18—C23	−41.7 (4)
C3—C4—C5—C6	−1.4 (5)	C23—C18—C19—C20	−0.9 (4)
C10—C5—C6—C7	−1.8 (4)	C11—C18—C19—C20	177.8 (3)
C4—C5—C6—C7	178.9 (3)	C18—C19—C20—C21	0.0 (5)
C10—C5—C6—C1	178.8 (3)	C18—C19—C20—N2	180.0 (3)
C4—C5—C6—C1	−0.5 (4)	C19—C20—C21—C22	0.6 (5)
C2—C1—C6—C7	−176.4 (3)	N2—C20—C21—C22	−179.4 (3)
C11—C1—C6—C7	4.7 (4)	C20—C21—C22—C23	−0.2 (6)
C2—C1—C6—C5	3.0 (4)	C21—C22—C23—C18	−0.7 (6)
C11—C1—C6—C5	−175.9 (2)	C19—C18—C23—C22	1.3 (5)
C5—C6—C7—C8	1.1 (4)	C11—C18—C23—C22	−177.5 (3)
C1—C6—C7—C8	−179.5 (3)	C13—C12—N1—C16	−60.4 (3)
C6—C7—C8—C9	0.4 (5)	C13—C12—N1—C11	173.6 (3)
C7—C8—C9—C10	−1.3 (6)	C15—C16—N1—C12	58.8 (3)
C8—C9—C10—C5	0.6 (6)	C15—C16—N1—C11	−178.4 (3)
C4—C5—C10—C9	−179.8 (4)	C1—C11—N1—C12	−72.9 (3)
C6—C5—C10—C9	1.0 (5)	C18—C11—N1—C12	160.5 (2)
C2—C1—C11—N1	−26.7 (3)	C1—C11—N1—C16	164.0 (2)
C6—C1—C11—N1	152.2 (2)	C18—C11—N1—C16	37.3 (3)
C2—C1—C11—C18	99.5 (3)	C21—C20—N2—O2	−178.3 (4)
C6—C1—C11—C18	−81.7 (3)	C19—C20—N2—O2	1.7 (5)

supplementary materials

N1—C12—C13—C14	57.6 (4)	C21—C20—N2—O3	2.6 (5)
N1—C12—C13—C17	-176.7 (3)	C19—C20—N2—O3	-177.4 (3)

Hydrogen-bond geometry (\AA , $^{\circ}$)

$D\cdots H$	$H\cdots A$	$D\cdots A$	$D\cdots H\cdots A$
O1—H1…N1	0.82	1.86	2.579 (3)

supplementary materials

Fig. 1

