

Acta Crystallographica Section E

Structure Reports

Online

ISSN 1600-5368

Diethyl 2-[(4-bromoanilino)methylidene]malonate

Zhi-Qiang Feng,* Xiao-Li Yang, Yuan-Feng Ye, Tao Dong and Huai-Qing Wang

College of Materials Engineering, Jinling Institute of Technology, No. 99 Hongjing Street, Nanjing 211169, People's Republic of China

Correspondence e-mail: fzq@jit.edu.cn


Received 1 November 2010; accepted 4 November 2010

Key indicators: single-crystal X-ray study; $T = 293$ K; mean $\sigma(\text{C}-\text{C}) = 0.012$ Å; R factor = 0.066; wR factor = 0.154; data-to-parameter ratio = 15.4.

In the title compound, $\text{C}_{14}\text{H}_{16}\text{BrNO}_4$, intermolecular $\text{C}-\text{H}\cdots\text{O}$ hydrogen bonds link the molecules, forming a stable structure. An intramolecular $\text{N}-\text{H}\cdots\text{O}$ hydrogen bond results in the formation of a six-membered ring and helps to establish the molecular conformation which is almost planar, with an r.m.s deviation of 0.0842 Å.

Related literature

For the preparation, see: Lager *et al.* (2006). For bond-length data, see: Allen *et al.* (1987).



Experimental

Crystal data

 $\text{C}_{14}\text{H}_{16}\text{BrNO}_4$ $M_r = 342.19$ Monoclinic, $P2_1$ $a = 9.2440$ (18) Å $b = 6.5000$ (13) Å $c = 13.448$ (3) Å $\beta = 110.10$ (3)° $V = 758.8$ (3) Å³ $Z = 2$ Mo $K\alpha$ radiation $\mu = 2.72$ mm⁻¹ $T = 293$ K $0.30 \times 0.10 \times 0.10$ mm

Data collection

Enraf-Nonius CAD-4 diffractometer
Absorption correction: ψ scan (North *et al.*, 1968)
 $T_{\min} = 0.496$, $T_{\max} = 0.773$
2851 measured reflections

2790 independent reflections
1606 reflections with $I > 2\sigma(I)$
 $R_{\text{int}} = 0.071$
3 standard reflections every 200 reflections
intensity decay: 1%

Refinement

 $R[F^2 > 2\sigma(F^2)] = 0.066$ $wR(F^2) = 0.154$ $S = 1.00$

2790 reflections

181 parameters

1 restraint

H-atom parameters constrained
 $\Delta\rho_{\text{max}} = 0.49$ e Å⁻³
 $\Delta\rho_{\text{min}} = -0.39$ e Å⁻³
Absolute structure: Flack (1983), 1253 Friedel pairs
Flack parameter: -0.01 (2)

Table 1

Hydrogen-bond geometry (Å, °).

$D-H\cdots A$	$D-H$	$H\cdots A$	$D\cdots A$	$D-H\cdots A$
$\text{N}-\text{H}0\text{A}\cdots\text{O}3$	0.86	1.95	2.615 (8)	134
$\text{C}1-\text{H}1\text{A}\cdots\text{O}3^{\text{i}}$	0.93	2.49	3.190 (10)	132
$\text{C}5-\text{H}5\text{A}\cdots\text{O}1^{\text{ii}}$	0.93	2.42	3.298 (9)	157

Symmetry codes: (i) $-x + 1, y + \frac{1}{2}, -z$; (ii) $-x + 2, y + \frac{1}{2}, -z + 1$.

Data collection: *CAD-4 EXPRESS* (Enraf-Nonius, 1989); cell refinement: *CAD-4 EXPRESS*; data reduction: *XCAD4* (Harms & Wocadlo, 1995); program(s) used to solve structure: *SHELXS97* (Sheldrick, 2008); program(s) used to refine structure: *SHELXL97* (Sheldrick, 2008); molecular graphics: *SHELXL97*; software used to prepare material for publication: *PLATON* (Spek, 2009).

The authors gratefully acknowledge Professor Hua-Qin Wang of the Analysis Center, Nanjing University, for providing the Enraf-Nonius CAD-4 diffractometer for this research project.

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: BQ2249).

References

- Allen, F. H., Kennard, O., Watson, D. G., Brammer, L., Orpen, A. G. & Taylor, R. (1987). *J. Chem. Soc. Perkin Trans. 2*, pp. S1–19.
- Enraf-Nonius (1989). *CAD-4 Software*. Enraf-Nonius, Delft, The Netherlands.
- Flack, H. D. (1983). *Acta Cryst.* **A39**, 876–881.
- Harms, K. & Wocadlo, S. (1995). *XCAD4*. University of Marburg, Germany.
- Lager, E., Andersson, P., Nilsson, J., Pettersson, I., Østergaard Nielsen, E., Nielsen, M., Sterner, O. & Liljefors, T. (2006). *J. Med. Chem.* **49**, 2526–2533.
- North, A. C. T., Phillips, D. C. & Mathews, F. S. (1968). *Acta Cryst.* **A24**, 351–359.
- Sheldrick, G. M. (2008). *Acta Cryst.* **A64**, 112–122.
- Spek, A. L. (2009). *Acta Cryst.* **D65**, 148–155.

supplementary materials

Acta Cryst. (2010). E66, o3119 [doi:10.1107/S1600536810045150]

Diethyl 2-[(4-bromoanilino)methylidene]malonate

Z.-Q. Feng, X.-L. Yang, Y.-F. Ye, T. Dong and H.-Q. Wang

Comment

We report herein the crystal structure of the title compound, diethyl 2-((4-bromophenylamino)methylene)malonate which is an important intermediate of synthesizing pyrazoloquinolinones. In the molecule of the title compound (Fig. 1), all bond lengths and angles (Allen *et al.*, 1987) are within normal ranges. The intramolecular N-H \cdots O hydrogen bond (Table 1) results in the formation of a six-membered ring (N/C7/C8/C12/O3/H0A). In the crystal structure, intermolecular weak C-H \cdots O hydrogen bonds link the molecules to form a stable structure (Fig. 2).

Experimental

The title compound, diethyl 2-((4-bromophenylamino)methylene)malonate was prepared by the literature method (Lager *et al.*, 2006). 4-bromoaniline (1.2 mmol) and diethyl ethoxymethylenemalonate (1.2 mmol) were mixed and heated at 403 K for 2 h. Low boiling components were evaporated at low pressure with a cold trap yielding diethyl 2-((4-bromophenylamino)methylene)malonate. The crude product was purified by recrystallization from diethyl ether yielding the title compound (73 % yield), as a white solid. Crystals suitable for X-ray analysis were obtained by slow evaporation of an methanol solution.

Refinement

H atoms were positioned geometrically, with N-H = 0.86 Å (for NH) and C-H = 0.93, 0.98 and 0.96 Å for aromatic, methine and methyl H, respectively, and constrained to ride on their parent atoms, with $U_{\text{iso}}(\text{H}) = xU_{\text{eq}}(\text{C}, \text{N})$, where $x = 1.5$ for methyl H and $x = 1.2$ for all other H atoms.

Figures




Fig. 1. The molecular structure of the title molecule, with the atom-numbering scheme. Displacement ellipsoids are drawn at the 50% probability level. Dashed lines show H-bonding.




Fig. 2. A packing diagram of the title compound. Hydrogen bonds are shown as dashed line.

Diethyl 2-[(4-bromoanilino)methylidene]malonate

Crystal data

$C_{14}H_{16}BrNO_4$	$F(000) = 348$
$M_r = 342.19$	$D_x = 1.498 \text{ Mg m}^{-3}$
Monoclinic, $P2_1$	Melting point: 367 K
Hall symbol: P 2yb	Mo $K\alpha$ radiation, $\lambda = 0.71073 \text{ \AA}$
$a = 9.2440 (18) \text{ \AA}$	Cell parameters from 25 reflections
$b = 6.5000 (13) \text{ \AA}$	$\theta = 9\text{--}14^\circ$
$c = 13.448 (3) \text{ \AA}$	$\mu = 2.72 \text{ mm}^{-1}$
$\beta = 110.10 (3)^\circ$	$T = 293 \text{ K}$
$V = 758.8 (3) \text{ \AA}^3$	Needle, colourless
$Z = 2$	$0.30 \times 0.10 \times 0.10 \text{ mm}$

Data collection

Enraf–Nonius CAD-4 diffractometer	1606 reflections with $I > 2\sigma(I)$
Radiation source: fine-focus sealed tube	$R_{\text{int}} = 0.071$
graphite	$\theta_{\text{max}} = 25.4^\circ$, $\theta_{\text{min}} = 1.6^\circ$
$\omega/2\theta$ scans	$h = -11 \rightarrow 11$
Absorption correction: ψ scan (North <i>et al.</i> , 1968)	$k = -7 \rightarrow 7$
$T_{\text{min}} = 0.496$, $T_{\text{max}} = 0.773$	$l = -5 \rightarrow 16$
2851 measured reflections	3 standard reflections every 200 reflections
2790 independent reflections	intensity decay: 1%

Refinement

Refinement on F^2	Secondary atom site location: difference Fourier map
Least-squares matrix: full	Hydrogen site location: inferred from neighbouring sites
$R[F^2 > 2\sigma(F^2)] = 0.066$	H-atom parameters constrained
$wR(F^2) = 0.154$	$w = 1/[\sigma^2(F_o^2) + (0.078P)^2 +]$
$S = 1.00$	where $P = (F_o^2 + 2F_c^2)/3$
2790 reflections	$(\Delta/\sigma)_{\text{max}} < 0.001$
181 parameters	$\Delta\rho_{\text{max}} = 0.49 \text{ e \AA}^{-3}$
1 restraint	$\Delta\rho_{\text{min}} = -0.39 \text{ e \AA}^{-3}$
Primary atom site location: structure-invariant direct methods	Absolute structure: Flack (1983), 1253 Friedel pairs
	Flack parameter: $-0.01 (2)$

Special details

Geometry. All esds (except the esd in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell esds are taken into account individually in the estimation of esds in distances, angles and torsion angles; correlations between esds in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell esds is used for estimating esds involving l.s. planes.

Refinement. Refinement of F^2 against ALL reflections. The weighted R-factor wR and goodness of fit S are based on F^2 , conventional R-factors R are based on F, with F set to zero for negative F^2 . The threshold expression of $F^2 > 2\sigma(F^2)$ is used only for calculating R-factors(gt) etc. and is not relevant to the choice of reflections for refinement. R-factors based on F^2 are statistically about twice as large as those based on F, and R- factors based on ALL data will be even larger.

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (\AA^2)

	x	y	z	$U_{\text{iso}}^*/U_{\text{eq}}$
Br	0.96900 (11)	0.78434 (19)	0.08931 (9)	0.1008 (4)
N	0.7057 (7)	0.0195 (9)	0.2183 (5)	0.0626 (16)
H0A	0.6242	-0.0249	0.1697	0.075*
C1	0.7111 (8)	0.2666 (15)	0.0881 (6)	0.0691 (19)
H1A	0.6318	0.1944	0.0387	0.083*
O1	0.8800 (8)	-0.2964 (10)	0.4901 (5)	0.116 (3)
O2	0.7014 (6)	-0.5254 (9)	0.4487 (4)	0.0768 (15)
C2	0.7701 (9)	0.4458 (13)	0.0594 (6)	0.068 (2)
H2A	0.7280	0.4973	-0.0091	0.082*
C3	0.8892 (9)	0.5442 (11)	0.1320 (8)	0.069 (2)
O3	0.4955 (6)	-0.2676 (8)	0.1562 (5)	0.0859 (18)
C4	0.9487 (9)	0.4794 (12)	0.2343 (7)	0.068 (2)
H4A	1.0276	0.5532	0.2832	0.082*
O4	0.4944 (6)	-0.5243 (8)	0.2634 (4)	0.0673 (13)
C5	0.8899 (8)	0.2992 (19)	0.2660 (5)	0.066 (2)
H5A	0.9301	0.2520	0.3354	0.079*
C6	0.7740 (8)	0.1980 (10)	0.1930 (6)	0.0558 (19)
C7	0.7510 (8)	-0.0856 (11)	0.3061 (6)	0.0521 (17)
H7A	0.8348	-0.0334	0.3608	0.062*
C8	0.6869 (8)	-0.2699 (11)	0.3269 (6)	0.062 (2)
C9	0.7662 (9)	-0.3565 (13)	0.4292 (6)	0.0594 (19)
C10	0.7736 (9)	-0.6258 (12)	0.5497 (6)	0.072 (2)
H10A	0.7952	-0.5273	0.6071	0.086*
H10B	0.8694	-0.6907	0.5524	0.086*
C11	0.6597 (12)	-0.7843 (17)	0.5578 (9)	0.124 (5)
H11A	0.7005	-0.8544	0.6244	0.186*
H11B	0.6411	-0.8817	0.5011	0.186*
H11C	0.5648	-0.7178	0.5530	0.186*
C12	0.5544 (9)	-0.3509 (12)	0.2415 (6)	0.0586 (18)
C13	0.3642 (9)	-0.6084 (13)	0.1829 (6)	0.079 (2)
H13A	0.3922	-0.6543	0.1233	0.094*
H13B	0.2832	-0.5063	0.1581	0.094*
C14	0.3121 (12)	-0.7836 (14)	0.2315 (8)	0.105 (4)

supplementary materials

H14A	0.2224	-0.8442	0.1805	0.157*
H14B	0.2872	-0.7363	0.2913	0.157*
H14C	0.3927	-0.8844	0.2543	0.157*

Atomic displacement parameters (\AA^2)

	U^{11}	U^{22}	U^{33}	U^{12}	U^{13}	U^{23}
Br	0.0966 (6)	0.0669 (5)	0.1532 (9)	-0.0125 (6)	0.0612 (6)	0.0184 (7)
N	0.051 (4)	0.065 (4)	0.065 (4)	-0.003 (3)	0.011 (3)	0.000 (3)
C1	0.072 (5)	0.056 (5)	0.079 (5)	0.008 (5)	0.024 (4)	0.012 (5)
O1	0.098 (5)	0.119 (6)	0.091 (4)	-0.047 (4)	-0.019 (4)	0.038 (4)
O2	0.056 (3)	0.080 (4)	0.085 (4)	-0.008 (3)	0.013 (3)	0.031 (3)
C2	0.060 (5)	0.073 (5)	0.074 (5)	0.008 (4)	0.026 (4)	0.026 (4)
C3	0.057 (5)	0.045 (4)	0.108 (7)	-0.014 (4)	0.034 (5)	-0.009 (5)
O3	0.078 (4)	0.071 (5)	0.088 (4)	-0.021 (3)	0.002 (3)	0.013 (3)
C4	0.058 (5)	0.060 (5)	0.087 (6)	-0.003 (4)	0.026 (5)	-0.006 (5)
O4	0.067 (3)	0.059 (3)	0.077 (3)	-0.010 (3)	0.026 (3)	0.000 (3)
C5	0.064 (4)	0.072 (5)	0.058 (4)	0.022 (6)	0.014 (4)	0.023 (5)
C6	0.059 (4)	0.042 (3)	0.076 (5)	-0.016 (3)	0.036 (4)	-0.015 (4)
C7	0.047 (4)	0.055 (4)	0.055 (5)	0.007 (3)	0.019 (4)	0.002 (3)
C8	0.052 (4)	0.058 (6)	0.088 (6)	-0.002 (3)	0.040 (4)	-0.008 (4)
C9	0.052 (5)	0.074 (5)	0.040 (4)	-0.009 (4)	0.001 (4)	0.000 (4)
C10	0.061 (5)	0.075 (5)	0.072 (5)	0.003 (4)	0.013 (4)	0.028 (4)
C11	0.087 (6)	0.132 (11)	0.157 (9)	0.002 (6)	0.048 (7)	0.075 (9)
C12	0.061 (5)	0.063 (5)	0.058 (5)	0.011 (4)	0.029 (4)	0.014 (4)
C13	0.065 (5)	0.086 (6)	0.073 (5)	-0.022 (5)	0.009 (4)	-0.022 (5)
C14	0.118 (8)	0.083 (7)	0.140 (9)	-0.035 (6)	0.080 (7)	-0.024 (6)

Geometric parameters (\AA , $^\circ$)

Br—C3	1.898 (7)	C5—C6	1.349 (11)
N—C7	1.302 (8)	C5—H5A	0.9300
N—C6	1.417 (9)	C7—C8	1.407 (9)
N—H0A	0.8600	C7—H7A	0.9300
C1—C2	1.396 (11)	C8—C9	1.432 (10)
C1—C6	1.401 (10)	C8—C12	1.459 (11)
C1—H1A	0.9300	C10—C11	1.503 (11)
O1—C9	1.157 (8)	C10—H10A	0.9700
O2—C9	1.320 (9)	C10—H10B	0.9700
O2—C10	1.446 (8)	C11—H11A	0.9600
C2—C3	1.356 (11)	C11—H11B	0.9600
C2—H2A	0.9300	C11—H11C	0.9600
C3—C4	1.361 (11)	C13—C14	1.474 (11)
O3—C12	1.215 (8)	C13—H13A	0.9700
C4—C5	1.417 (14)	C13—H13B	0.9700
C4—H4A	0.9300	C14—H14A	0.9600
O4—C12	1.333 (8)	C14—H14B	0.9600
O4—C13	1.423 (8)	C14—H14C	0.9600

C7—N—C6	128.0 (6)	O1—C9—C8	126.0 (8)
C7—N—H0A	116.0	O2—C9—C8	113.6 (7)
C6—N—H0A	116.0	O2—C10—C11	105.6 (7)
C2—C1—C6	118.6 (8)	O2—C10—H10A	110.6
C2—C1—H1A	120.7	C11—C10—H10A	110.6
C6—C1—H1A	120.7	O2—C10—H10B	110.6
C9—O2—C10	117.9 (6)	C11—C10—H10B	110.6
C3—C2—C1	119.5 (8)	H10A—C10—H10B	108.8
C3—C2—H2A	120.3	C10—C11—H11A	109.5
C1—C2—H2A	120.3	C10—C11—H11B	109.5
C2—C3—C4	121.9 (7)	H11A—C11—H11B	109.5
C2—C3—Br	118.3 (7)	C10—C11—H11C	109.5
C4—C3—Br	119.8 (6)	H11A—C11—H11C	109.5
C3—C4—C5	119.8 (7)	H11B—C11—H11C	109.5
C3—C4—H4A	120.1	O3—C12—O4	120.0 (7)
C5—C4—H4A	120.1	O3—C12—C8	124.4 (7)
C12—O4—C13	117.7 (6)	O4—C12—C8	115.6 (6)
C6—C5—C4	118.3 (7)	O4—C13—C14	106.2 (7)
C6—C5—H5A	120.8	O4—C13—H13A	110.5
C4—C5—H5A	120.8	C14—C13—H13A	110.5
C5—C6—C1	121.9 (7)	O4—C13—H13B	110.5
C5—C6—N	122.1 (7)	C14—C13—H13B	110.5
C1—C6—N	116.0 (7)	H13A—C13—H13B	108.7
N—C7—C8	126.9 (7)	C13—C14—H14A	109.5
N—C7—H7A	116.5	C13—C14—H14B	109.5
C8—C7—H7A	116.5	H14A—C14—H14B	109.5
C7—C8—C9	114.5 (7)	C13—C14—H14C	109.5
C7—C8—C12	116.6 (7)	H14A—C14—H14C	109.5
C9—C8—C12	128.9 (6)	H14B—C14—H14C	109.5
O1—C9—O2	120.3 (7)		
C6—C1—C2—C3	2.2 (11)	C10—O2—C9—O1	2.6 (12)
C1—C2—C3—C4	-3.4 (11)	C10—O2—C9—C8	179.5 (6)
C1—C2—C3—Br	178.8 (6)	C7—C8—C9—O1	-5.5 (11)
C2—C3—C4—C5	2.7 (12)	C12—C8—C9—O1	171.8 (9)
Br—C3—C4—C5	-179.5 (6)	C7—C8—C9—O2	177.9 (6)
C3—C4—C5—C6	-0.8 (12)	C12—C8—C9—O2	-4.9 (11)
C4—C5—C6—C1	-0.3 (11)	C9—O2—C10—C11	169.0 (7)
C4—C5—C6—N	-178.4 (7)	C13—O4—C12—O3	1.8 (9)
C2—C1—C6—C5	-0.4 (11)	C13—O4—C12—C8	179.5 (6)
C2—C1—C6—N	177.8 (6)	C7—C8—C12—O3	-1.7 (10)
C7—N—C6—C5	-8.8 (11)	C9—C8—C12—O3	-178.9 (7)
C7—N—C6—C1	173.0 (7)	C7—C8—C12—O4	-179.3 (5)
C6—N—C7—C8	-175.6 (6)	C9—C8—C12—O4	3.5 (10)
N—C7—C8—C9	176.9 (7)	C12—O4—C13—C14	-173.9 (6)
N—C7—C8—C12	-0.7 (9)		

Hydrogen-bond geometry (\AA , $^\circ$)

<i>D</i> —H \cdots <i>A</i>	<i>D</i> —H	H \cdots <i>A</i>	<i>D</i> \cdots <i>A</i>	<i>D</i> —H \cdots <i>A</i>
-------------------------------	-------------	---------------------	----------------------------	-------------------------------

supplementary materials

N—H0A···O3	0.86	1.95	2.615 (8)	134
C1—H1A···O3 ⁱ	0.93	2.49	3.190 (10)	132
C5—H5A···O1 ⁱⁱ	0.93	2.42	3.298 (9)	157

Symmetry codes: (i) $-x+1, y+1/2, -z$; (ii) $-x+2, y+1/2, -z+1$.

Fig. 1




Fig. 2

