

Pyridine-2-carboximidamide chloride monohydrate

Qifan Chen,^a Huidong Zhang,^a Fang Zhang^b and Fei Liu^{b*}

^aExperiment Center, Eastern Liaoning University, No. 325 Wenhua Road, Yuanbao District, Dandong City, Liaoning Province 118003, People's Republic of China, and

^bCollege of Chemical Engineering & Materials, Eastern Liaoning University, No. 325 Wenhua Road, Yuanbao District, Dandong City, Liaoning Province 118003, People's Republic of China

Correspondence e-mail: berylliu8090@sina.com


Received 26 October 2010; accepted 9 November 2010

Key indicators: single-crystal X-ray study; $T = 293\text{ K}$; mean $\sigma(\text{C}-\text{C}) = 0.003\text{ \AA}$; R factor = 0.035; wR factor = 0.117; data-to-parameter ratio = 14.2.

The title compound, $\text{C}_6\text{H}_8\text{N}_3^+\cdot\text{Cl}^-\cdot\text{H}_2\text{O}$, crystallizes with three formula units in the asymmetric unit. The cations are non-planar with the $-\text{C}(\text{NH}_2)_2$ groups twisted out of the ring planes. Each pyridine carboximidamide cation is linked to another cation through $\text{N}-\text{H}\cdots\text{N}$ hydrogen bonds, to chloride ions by $\text{N}-\text{H}\cdots\text{Cl}$ hydrogen bonds, and to water molecules by $\text{N}-\text{H}\cdots\text{O}$ hydrogen bonds. Water molecules and chloride ions are also linked together *via* $\text{O}-\text{H}\cdots\text{Cl}$ hydrogen bonds. In the crystal, all these intermolecular interactions result in a three-dimensional network.

Related literature

For related structures, see: Guo *et al.* (2005); Fan *et al.* (2009); Góker *et al.* (2005); Walther *et al.* (2006).



Experimental

Crystal data

$\text{C}_6\text{H}_8\text{N}_3^+\cdot\text{Cl}^-\cdot\text{H}_2\text{O}$
 $M_r = 175.62$
Triclinic, $P\bar{1}$
 $a = 7.2570(15)\text{ \AA}$
 $b = 11.146(2)\text{ \AA}$

$c = 16.862(3)\text{ \AA}$
 $\alpha = 79.24(3)^\circ$
 $\beta = 82.14(3)^\circ$
 $\gamma = 78.34(3)^\circ$
 $V = 1305.3(5)\text{ \AA}^3$

$Z = 6$
Mo $K\alpha$ radiation
 $\mu = 0.39\text{ mm}^{-1}$

$T = 293\text{ K}$
 $0.25 \times 0.24 \times 0.22\text{ mm}$

Data collection

Rigaku R-AXIS RAPID diffractometer
Absorption correction: multi-scan (*ABSCOR*; Higashi, 1995)
 $T_{\min} = 0.909$, $T_{\max} = 0.918$

12872 measured reflections
5919 independent reflections
4550 reflections with $I > 2\sigma(I)$
 $R_{\text{int}} = 0.024$

Refinement

$R[F^2 > 2\sigma(F^2)] = 0.035$
 $wR(F^2) = 0.117$
 $S = 1.11$
5919 reflections

418 parameters
All H-atom parameters refined
 $\Delta\rho_{\text{max}} = 0.23\text{ e \AA}^{-3}$
 $\Delta\rho_{\text{min}} = -0.26\text{ e \AA}^{-3}$

Table 1
Hydrogen-bond geometry (\AA , $^\circ$).

$D-\text{H}\cdots A$	$D-\text{H}$	$\text{H}\cdots A$	$D\cdots A$	$D-\text{H}\cdots A$
N23—H23B \cdots O1 ⁱ	0.87 (2)	1.97 (2)	2.829 (2)	173 (2)
N22—H22B \cdots C11	0.89 (3)	2.31 (3)	3.187 (2)	169 (2)
N12—H12B \cdots Cl3 ⁱⁱ	0.91 (2)	2.35 (2)	3.2522 (19)	170.2 (18)
O1—H1B \cdots Cl3	0.80 (3)	2.37 (3)	3.163 (2)	169 (2)
N2—H2A \cdots Cl2 ⁱⁱⁱ	0.85 (3)	2.49 (3)	3.326 (2)	167 (2)
O1—H10 \cdots Cl2 ^{iv}	0.98 (4)	2.22 (4)	3.1902 (19)	173 (3)
N3—H3A \cdots O2	0.87 (2)	1.94 (2)	2.808 (2)	174.9 (19)
N2—H2B \cdots C11	0.96 (2)	2.32 (2)	3.2754 (17)	170.9 (17)
N12—H12A \cdots Cl3 ^v	0.89 (2)	2.43 (2)	3.313 (2)	171 (2)
N23—H23A \cdots Cl3 ^{vi}	0.87 (2)	2.50 (2)	3.324 (2)	158 (2)
N13—H13B \cdots O3 ⁱⁱ	0.92 (2)	1.97 (2)	2.883 (2)	178 (2)
N22—H22A \cdots Cl2 ⁱⁱⁱ	0.91 (2)	2.43 (3)	3.3194 (18)	168 (2)
N13—H13A \cdots C11	0.91 (2)	2.48 (2)	3.335 (2)	156.9 (19)
O3—H03B \cdots Cl2 ^{iv}	0.84 (4)	2.45 (4)	3.286 (2)	174 (3)
O2—H0B \cdots C11	0.85 (3)	2.28 (3)	3.1171 (18)	172 (2)
O2—H0A \cdots Cl2	0.79 (3)	2.40 (3)	3.1833 (18)	175 (3)
O3—H03A \cdots Cl3	0.94 (4)	2.25 (4)	3.1586 (18)	164 (3)

Symmetry codes: (i) $x - 1, y, z$; (ii) $x, y - 1, z$; (iii) $-x, -y + 1, -z$; (iv) $-x + 1, -y + 1, -z$; (v) $-x + 1, -y + 1, -z + 1$; (vi) $-x, -y + 1, -z + 1$.

Data collection: *RAPID-AUTO* (Rigaku, 1998); cell refinement: *RAPID-AUTO*; data reduction: *RAPID-AUTO*; program(s) used to solve structure: *SHELXS97* (Sheldrick, 2008); program(s) used to refine structure: *SHELXL97* (Sheldrick, 2008); molecular graphics: *DIAMOND* (Brandenburg, 1999) and *SHELXP97* (Sheldrick, 2008); software used to prepare material for publication: *SHELXL97*.

The authors gratefully acknowledge financial support from the Education Department of Liaoning Province (2009 A 265) and Liaoning University.

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: ZQ2070).

References

- Brandenburg, K. (1999). *DIAMOND*. Crystal Impact GbR, Bonn, Germany.
- Fan, P., Wang, L. & Zhang, H. (2009). *Acta Cryst. E65*, o2408.
- Góker, H., Özden, S., Yıldız, S. & Boykin, D. W. (2005). *Eur. J. Med. Chem.* **40**, 1062–1069.
- Guo, J., Wong, W.-K. & Wong, W.-Y. (2005). *Polyhedron*, **24**, 927–939.
- Higashi, T. (1995). *ABSCOR*. Rigaku Corporation, Tokyo, Japan.
- Rigaku (1998). *RAPID-AUTO*. Rigaku Corporation, Tokyo, Japan.
- Sheldrick, G. M. (2008). *Acta Cryst. A64*, 112–122.
- Walther, M., Wermann, H., Lutsche, M., Gúnther, W., Görsls, H. & Anders, E. (2006). *J. Org. Chem.* **71**, 1399–1406.

supplementary materials

Acta Cryst. (2010). E66, o3169 [doi:10.1107/S1600536810046106]

Pyridine-2-carboximidamide chloride monohydrate

Q. Chen, H. Zhang, F. Zhang and F. Liu

Comment

The pyridinecarboximidamide derivatives have attracted a great attention due to their very potent antibacterial activities, interesting biological properties and applications in coordination chemistry, but a few samples have been reported (Guo *et al.*, 2005; Góker *et al.*, 2005; Walther *et al.*, 2006; Fan *et al.*, 2009). Here we report the structure of title compound $C_6H_8N_3^+ \cdot Cl^- \cdot H_2O$. The title compound is an organic salt and crystallizes with three formula units in the asymmetric unit (Fig. 1). The molecules are nonplanar with the $C(NH_2)_2$ groups twisted out of the ring planes. The bond lengths and angles exhibit normal values and are comparable with those in the related structures (Guo *et al.*, 2005; Fan *et al.*, 2009).

In the crystal structure, pyridine carboximidamide cations adopt three different connecting types (Fig. 2). Each pyridine carboximidamide cation is linked to another cation through N—H···N hydrogen bonds, to chloride ions by N—H···Cl hydrogen bonds, and to water molecules by N—H···O hydrogen bonds. Water molecules and chloride ions are also linked together *via* O—H···Cl hydrogen bonds (Table 1). In the crystal structure, all these intermolecular interactions result in a three-dimensional network (Fig. 3).

Experimental

To a solution of sodium methoxide (5.15 mmol) in methanol (50 mL) was added 2-cyanopyridine (5.2 g, 4.99 mmol). The mixture was stirred at room temperature for 4 h. Then ammonium chloride (2.9 g, 5.42 mmol) was slowly added to the resulting solution and the mixture was stirred at room temperature for 68 h. The resulting suspension was filtered and the solvent was removed from the filtrate under reduced pressure. Purification by wash with ethyl ether gave 2-amidinopyridine hydrochloride (5.62 g, 92%) as an off-white solid. Colorless crystals were formed after several days.

Refinement

The H atoms were initially found in a difference Fourier map and their coordinates were refined freely.

Figures



Fig. 1. The molecular structure of the title compound showing 30% probability displacement ellipsoids for non-H atoms.




Fig. 2. View of the hydrogen bonding interactions in the title compound.

supplementary materials




Fig. 3. Crystal packing of the title compound.

Pyridine-2-carboximidamide chloride monohydrate

Crystal data

$C_6H_8N_3^+\cdot Cl^- \cdot H_2O$	$Z = 6$
$M_r = 175.62$	$F(000) = 552$
Triclinic, $P\bar{1}$	$D_x = 1.340 \text{ Mg m}^{-3}$
Hall symbol: -P 1	Mo $K\alpha$ radiation, $\lambda = 0.71073 \text{ \AA}$
$a = 7.2570 (15) \text{ \AA}$	Cell parameters from 9445 reflections
$b = 11.146 (2) \text{ \AA}$	$\theta = 3.0\text{--}27.4^\circ$
$c = 16.862 (3) \text{ \AA}$	$\mu = 0.39 \text{ mm}^{-1}$
$\alpha = 79.24 (3)^\circ$	$T = 293 \text{ K}$
$\beta = 82.14 (3)^\circ$	Block, colourless
$\gamma = 78.34 (3)^\circ$	$0.25 \times 0.24 \times 0.22 \text{ mm}$
$V = 1305.3 (5) \text{ \AA}^3$	

Data collection

Rigaku R-AXIS RAPID diffractometer	5919 independent reflections
Radiation source: fine-focus sealed tube graphite	4550 reflections with $I > 2\sigma(I)$
Detector resolution: 10 pixels mm^{-1}	$R_{\text{int}} = 0.024$
ω scans	$\theta_{\text{max}} = 27.5^\circ, \theta_{\text{min}} = 3.0^\circ$
Absorption correction: multi-scan (<i>ABSCOR</i> ; Higashi, 1995)	$h = -9 \rightarrow 9$
$T_{\text{min}} = 0.909, T_{\text{max}} = 0.918$	$k = -14 \rightarrow 14$
12872 measured reflections	$l = -21 \rightarrow 21$

Refinement

Refinement on F^2	Primary atom site location: structure-invariant direct methods
Least-squares matrix: full	Secondary atom site location: difference Fourier map
$R[F^2 > 2\sigma(F^2)] = 0.035$	Hydrogen site location: inferred from neighbouring sites
$wR(F^2) = 0.117$	All H-atom parameters refined
$S = 1.11$	$w = 1/[\sigma^2(F_o^2) + (0.0684P)^2]$
5919 reflections	where $P = (F_o^2 + 2F_c^2)/3$
418 parameters	$(\Delta/\sigma)_{\text{max}} = 0.001$
0 restraints	$\Delta\rho_{\text{max}} = 0.23 \text{ e \AA}^{-3}$
	$\Delta\rho_{\text{min}} = -0.26 \text{ e \AA}^{-3}$

Special details

Geometry. All e.s.d.'s (except the e.s.d. in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell e.s.d.'s are taken into account individually in the estimation of e.s.d.'s in distances, angles and torsion angles; correlations between e.s.d.'s in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell e.s.d.'s is used for estimating e.s.d.'s involving l.s. planes.

Refinement. Refinement of F^2 against ALL reflections. The weighted R -factor wR and goodness of fit S are based on F^2 , conventional R -factors R are based on F , with F set to zero for negative F^2 . The threshold expression of $F^2 > \sigma(F^2)$ is used only for calculating R -factors(gt) etc. and is not relevant to the choice of reflections for refinement. R -factors based on F^2 are statistically about twice as large as those based on F , and R -factors based on ALL data will be even larger.

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (\AA^2)

	<i>x</i>	<i>y</i>	<i>z</i>	$U_{\text{iso}}^*/U_{\text{eq}}$
Cl1	0.17320 (7)	0.30230 (4)	0.10288 (2)	0.04346 (13)
O1	0.7888 (3)	0.60439 (14)	0.40146 (10)	0.0647 (4)
H1B	0.690 (4)	0.647 (3)	0.4141 (15)	0.073 (8)*
H10	0.821 (5)	0.635 (3)	0.344 (2)	0.120 (11)*
N1	0.3062 (2)	0.86483 (13)	-0.07194 (8)	0.0432 (3)
C1	0.3052 (3)	0.98536 (18)	-0.07462 (13)	0.0540 (5)
H1	0.336 (4)	1.031 (2)	-0.1243 (15)	0.078 (7)*
Cl2	0.13596 (7)	0.30724 (4)	-0.21061 (3)	0.04888 (14)
O2	0.3482 (2)	0.37193 (13)	-0.07519 (9)	0.0530 (4)
H0B	0.305 (4)	0.345 (2)	-0.0273 (16)	0.077 (8)*
H0A	0.297 (4)	0.351 (2)	-0.1073 (17)	0.075 (8)*
N2	0.1754 (2)	0.60101 (14)	0.06004 (10)	0.0429 (3)
H2A	0.111 (4)	0.629 (2)	0.1009 (15)	0.071 (7)*
H2B	0.180 (3)	0.513 (2)	0.0663 (12)	0.053 (6)*
C2	0.2604 (3)	1.04352 (17)	-0.00802 (14)	0.0561 (5)
H2	0.261 (3)	1.131 (2)	-0.0152 (12)	0.055 (6)*
Cl3	0.43514 (7)	0.80296 (4)	0.45146 (3)	0.05099 (14)
O3	0.4557 (3)	0.86529 (14)	0.26022 (10)	0.0627 (4)
H03B	0.561 (5)	0.826 (3)	0.24445 (19)	0.102 (11)*
H03A	0.448 (5)	0.831 (3)	0.316 (2)	0.121 (12)*
N3	0.3540 (2)	0.61884 (14)	-0.06282 (9)	0.0426 (3)
H3B	0.410 (3)	0.669 (2)	-0.1007 (13)	0.054 (6)*
H3A	0.350 (3)	0.544 (2)	-0.0696 (12)	0.051 (5)*
C3	0.2174 (3)	0.97567 (18)	0.06587 (13)	0.0564 (5)
H3	0.189 (3)	1.014 (2)	0.1127 (14)	0.070 (7)*
C4	0.2181 (3)	0.85015 (18)	0.07107 (11)	0.0478 (4)
H4	0.192 (3)	0.800 (2)	0.1214 (14)	0.067 (7)*
C5	0.2600 (2)	0.80020 (14)	0.00027 (9)	0.0343 (3)
C6	0.2627 (2)	0.66679 (14)	-0.00077 (9)	0.0338 (3)
N11	0.4303 (2)	0.37190 (13)	0.25233 (9)	0.0428 (3)
C11	0.4055 (3)	0.49449 (19)	0.24780 (13)	0.0540 (5)
H11	0.446 (3)	0.542 (2)	0.2001 (13)	0.058 (6)*
N12	0.4363 (3)	0.09814 (16)	0.39452 (10)	0.0502 (4)

supplementary materials

H12B	0.440 (3)	0.014 (2)	0.4040 (13)	0.055 (6)*
H12A	0.456 (3)	0.128 (2)	0.4371 (15)	0.065 (7)*
C12	0.3223 (3)	0.55573 (19)	0.31194 (14)	0.0604 (5)
H12	0.311 (4)	0.648 (2)	0.3024 (15)	0.081 (8)*
N13	0.3872 (2)	0.12940 (15)	0.25963 (10)	0.0447 (3)
H13B	0.413 (3)	0.045 (2)	0.2600 (13)	0.059 (6)*
H13A	0.360 (3)	0.187 (2)	0.2153 (13)	0.060 (6)*
C13	0.2613 (3)	0.4879 (2)	0.38308 (14)	0.0599 (5)
H13	0.207 (4)	0.528 (2)	0.4224 (16)	0.078 (8)*
C14	0.2851 (3)	0.36072 (18)	0.38966 (12)	0.0477 (4)
H14	0.241 (3)	0.3113 (19)	0.4350 (12)	0.048 (5)*
C15	0.3718 (2)	0.30783 (15)	0.32274 (10)	0.0368 (3)
C16	0.4002 (2)	0.17137 (16)	0.32577 (10)	0.0384 (4)
N21	-0.0617 (2)	0.10605 (13)	0.39607 (8)	0.0441 (3)
C21	-0.0357 (3)	-0.01675 (19)	0.40054 (13)	0.0563 (5)
H21	-0.005 (3)	-0.059 (2)	0.4527 (13)	0.062 (6)*
N22	-0.0909 (2)	0.38620 (15)	0.25737 (9)	0.0446 (3)
H22B	-0.032 (3)	0.359 (2)	0.2128 (15)	0.066 (7)*
H22A	-0.111 (3)	0.470 (2)	0.2526 (14)	0.069 (7)*
C22	-0.0480 (3)	-0.07293 (19)	0.33556 (13)	0.0559 (5)
H22	-0.030 (3)	-0.160 (2)	0.3423 (13)	0.061 (6)*
N23	-0.2026 (2)	0.35240 (16)	0.39229 (10)	0.0453 (4)
H23B	-0.215 (3)	0.430 (2)	0.3963 (13)	0.058 (6)*
H23A	-0.232 (3)	0.299 (2)	0.4342 (14)	0.057 (6)*
C23	-0.0891 (3)	-0.00047 (19)	0.26314 (12)	0.0526 (5)
H23	-0.103 (4)	-0.035 (2)	0.2163 (15)	0.077 (7)*
C24	-0.1160 (3)	0.12748 (18)	0.25641 (10)	0.0431 (4)
H24	-0.146 (3)	0.1764 (18)	0.2077 (12)	0.048 (5)*
C25	-0.1009 (2)	0.17539 (15)	0.32452 (9)	0.0338 (3)
C26	-0.1327 (2)	0.31196 (15)	0.32427 (9)	0.0352 (3)

Atomic displacement parameters (\AA^2)

	U^{11}	U^{22}	U^{33}	U^{12}	U^{13}	U^{23}
Cl1	0.0579 (3)	0.0378 (2)	0.0333 (2)	-0.01169 (18)	0.00298 (17)	-0.00477 (16)
O1	0.0845 (12)	0.0469 (8)	0.0521 (9)	0.0040 (8)	0.0066 (8)	-0.0091 (7)
N1	0.0551 (9)	0.0372 (8)	0.0361 (7)	-0.0117 (7)	-0.0026 (6)	-0.0009 (6)
C1	0.0739 (14)	0.0379 (10)	0.0502 (11)	-0.0197 (9)	-0.0090 (10)	0.0050 (9)
Cl2	0.0640 (3)	0.0430 (2)	0.0393 (2)	-0.0127 (2)	0.00567 (19)	-0.01046 (18)
O2	0.0740 (10)	0.0492 (8)	0.0412 (7)	-0.0272 (7)	0.0026 (7)	-0.0098 (6)
N2	0.0535 (9)	0.0310 (7)	0.0428 (8)	-0.0122 (7)	0.0073 (7)	-0.0061 (6)
C2	0.0651 (13)	0.0295 (9)	0.0749 (14)	-0.0119 (8)	-0.0056 (11)	-0.0094 (9)
Cl3	0.0642 (3)	0.0439 (2)	0.0370 (2)	-0.0035 (2)	0.00486 (19)	-0.00088 (18)
O3	0.0851 (12)	0.0472 (8)	0.0474 (8)	-0.0042 (8)	0.0026 (8)	-0.0020 (7)
N3	0.0529 (9)	0.0337 (8)	0.0407 (8)	-0.0116 (7)	0.0084 (7)	-0.0102 (7)
C3	0.0654 (13)	0.0476 (11)	0.0625 (12)	-0.0181 (10)	0.0094 (10)	-0.0280 (10)
C4	0.0608 (12)	0.0435 (10)	0.0397 (9)	-0.0164 (9)	0.0071 (8)	-0.0098 (8)
C5	0.0341 (8)	0.0301 (8)	0.0375 (8)	-0.0047 (6)	-0.0016 (6)	-0.0056 (6)

C6	0.0351 (8)	0.0304 (8)	0.0346 (8)	-0.0056 (6)	-0.0029 (6)	-0.0032 (6)
N11	0.0463 (8)	0.0403 (8)	0.0413 (8)	-0.0133 (7)	-0.0022 (6)	-0.0011 (6)
C11	0.0637 (13)	0.0431 (10)	0.0535 (11)	-0.0168 (9)	-0.0050 (10)	0.0036 (9)
N12	0.0662 (11)	0.0378 (9)	0.0454 (9)	-0.0115 (8)	-0.0133 (8)	0.0036 (7)
C12	0.0682 (14)	0.0369 (10)	0.0749 (15)	-0.0044 (9)	-0.0132 (11)	-0.0077 (10)
N13	0.0533 (9)	0.0357 (8)	0.0439 (8)	-0.0081 (7)	-0.0068 (7)	-0.0020 (7)
C13	0.0672 (14)	0.0520 (12)	0.0582 (12)	0.0050 (10)	-0.0078 (11)	-0.0191 (10)
C14	0.0487 (10)	0.0482 (10)	0.0398 (9)	-0.0024 (8)	-0.0011 (8)	-0.0003 (8)
C15	0.0332 (8)	0.0374 (9)	0.0389 (8)	-0.0072 (7)	-0.0064 (7)	-0.0013 (7)
C16	0.0322 (8)	0.0416 (9)	0.0394 (8)	-0.0090 (7)	-0.0025 (7)	0.0000 (7)
N21	0.0555 (9)	0.0392 (8)	0.0350 (7)	-0.0006 (7)	-0.0085 (6)	-0.0053 (6)
C21	0.0746 (14)	0.0421 (10)	0.0474 (10)	0.0028 (9)	-0.0149 (10)	-0.0041 (8)
N22	0.0531 (9)	0.0370 (8)	0.0395 (8)	-0.0089 (7)	0.0081 (7)	-0.0044 (7)
C22	0.0681 (14)	0.0391 (10)	0.0615 (12)	-0.0083 (9)	-0.0054 (10)	-0.0137 (9)
N23	0.0547 (10)	0.0402 (9)	0.0387 (8)	-0.0099 (7)	0.0087 (7)	-0.0088 (7)
C23	0.0634 (13)	0.0548 (11)	0.0470 (10)	-0.0201 (10)	-0.0012 (9)	-0.0202 (9)
C24	0.0499 (10)	0.0495 (10)	0.0328 (8)	-0.0184 (8)	-0.0036 (7)	-0.0046 (7)
C25	0.0287 (7)	0.0381 (8)	0.0337 (7)	-0.0066 (6)	0.0006 (6)	-0.0060 (6)
C26	0.0309 (8)	0.0383 (9)	0.0355 (8)	-0.0074 (7)	0.0001 (6)	-0.0053 (7)

Geometric parameters (Å, °)

O1—H1B	0.80 (3)	N12—H12A	0.89 (2)
O1—H10	0.98 (4)	C12—C13	1.363 (3)
N1—C5	1.331 (2)	C12—H12	1.00 (3)
N1—C1	1.334 (2)	N13—C16	1.309 (2)
C1—C2	1.369 (3)	N13—H13B	0.92 (2)
C1—H1	0.92 (2)	N13—H13A	0.91 (2)
O2—H0B	0.85 (3)	C13—C14	1.378 (3)
O2—H0A	0.79 (3)	C13—H13	0.87 (3)
N2—C6	1.311 (2)	C14—C15	1.386 (3)
N2—H2A	0.85 (3)	C14—H14	0.91 (2)
N2—H2B	0.96 (2)	C15—C16	1.485 (2)
C2—C3	1.364 (3)	N21—C21	1.333 (2)
C2—H2	0.96 (2)	N21—C25	1.338 (2)
O3—H03B	0.84 (4)	C21—C22	1.381 (3)
O3—H03A	0.94 (4)	C21—H21	0.95 (2)
N3—C6	1.302 (2)	N22—C26	1.307 (2)
N3—H3B	0.88 (2)	N22—H22B	0.89 (3)
N3—H3A	0.87 (2)	N22—H22A	0.91 (2)
C3—C4	1.384 (3)	C22—C23	1.366 (3)
C3—H3	0.95 (2)	C22—H22	0.94 (2)
C4—C5	1.381 (2)	N23—C26	1.311 (2)
C4—H4	0.94 (2)	N23—H23B	0.87 (2)
C5—C6	1.486 (2)	N23—H23A	0.87 (2)
N11—C15	1.330 (2)	C23—C24	1.385 (3)
N11—C11	1.331 (2)	C23—H23	0.97 (2)
C11—C12	1.390 (3)	C24—C25	1.380 (2)
C11—H11	0.92 (2)	C24—H24	0.924 (19)

supplementary materials

N12—C16	1.312 (2)	C25—C26	1.492 (2)
N12—H12B	0.91 (2)		
H1B—O1—H10	106 (3)	C16—N13—H13A	116.6 (13)
C5—N1—C1	116.70 (15)	H13B—N13—H13A	124.4 (19)
N1—C1—C2	123.59 (18)	C12—C13—C14	119.3 (2)
N1—C1—H1	117.3 (16)	C12—C13—H13	117.7 (17)
C2—C1—H1	119.1 (16)	C14—C13—H13	123.0 (17)
H0B—O2—H0A	111 (2)	C13—C14—C15	117.69 (19)
C6—N2—H2A	125.2 (16)	C13—C14—H14	122.7 (13)
C6—N2—H2B	124.9 (13)	C15—C14—H14	119.5 (12)
H2A—N2—H2B	110 (2)	N11—C15—C14	124.30 (16)
C3—C2—C1	119.11 (17)	N11—C15—C16	115.39 (15)
C3—C2—H2	122.4 (12)	C14—C15—C16	120.29 (16)
C1—C2—H2	118.5 (12)	N13—C16—N12	122.82 (17)
H03B—O3—H03A	100 (3)	N13—C16—C15	118.41 (15)
C6—N3—H3B	116.3 (14)	N12—C16—C15	118.77 (16)
C6—N3—H3A	122.4 (14)	C21—N21—C25	116.99 (15)
H3B—N3—H3A	120.9 (19)	N21—C21—C22	122.95 (18)
C2—C3—C4	118.91 (18)	N21—C21—H21	111.6 (13)
C2—C3—H3	120.1 (14)	C22—C21—H21	125.4 (13)
C4—C3—H3	121.0 (14)	C26—N22—H22B	123.0 (15)
C5—C4—C3	117.91 (18)	C26—N22—H22A	124.0 (15)
C5—C4—H4	121.0 (13)	H22B—N22—H22A	113 (2)
C3—C4—H4	121.1 (13)	C23—C22—C21	119.16 (18)
N1—C5—C4	123.74 (15)	C23—C22—H22	121.3 (13)
N1—C5—C6	114.14 (13)	C21—C22—H22	119.5 (13)
C4—C5—C6	122.09 (15)	C26—N23—H23B	121.2 (15)
N3—C6—N2	122.05 (15)	C26—N23—H23A	118.2 (14)
N3—C6—C5	118.14 (15)	H23B—N23—H23A	121 (2)
N2—C6—C5	119.81 (15)	C22—C23—C24	119.30 (17)
C15—N11—C11	116.63 (16)	C22—C23—H23	122.4 (15)
N11—C11—C12	123.22 (19)	C24—C23—H23	118.3 (15)
N11—C11—H11	119.1 (14)	C25—C24—C23	117.51 (17)
C12—C11—H11	117.7 (14)	C25—C24—H24	123.3 (12)
C16—N12—H12B	124.8 (13)	C23—C24—H24	119.2 (12)
C16—N12—H12A	121.0 (15)	N21—C25—C24	124.10 (15)
H12B—N12—H12A	114.1 (19)	N21—C25—C26	114.20 (14)
C13—C12—C11	118.88 (19)	C24—C25—C26	121.68 (15)
C13—C12—H12	124.1 (15)	N22—C26—N23	122.63 (16)
C11—C12—H12	117.0 (15)	N22—C26—C25	119.59 (15)
C16—N13—H13B	119.0 (13)	N23—C26—C25	117.78 (15)

Hydrogen-bond geometry (\AA , $^\circ$)

$D—\text{H}\cdots A$	$D—\text{H}$	$\text{H}\cdots A$	$D\cdots A$	$D—\text{H}\cdots A$
N23—H23B···O1 ⁱ	0.87 (2)	1.97 (2)	2.829 (2)	173 (2)
N22—H22B···Cl1	0.89 (3)	2.31 (3)	3.187 (2)	169 (2)
N12—H12B···Cl3 ⁱⁱ	0.91 (2)	2.35 (2)	3.2522 (19)	170.2 (18)

supplementary materials

O1—H1B···Cl3	0.80 (3)	2.37 (3)	3.163 (2)	169 (2)
N2—H2A···Cl2 ⁱⁱⁱ	0.85 (3)	2.49 (3)	3.326 (2)	167 (2)
O1—H10···Cl2 ^{iv}	0.98 (4)	2.22 (4)	3.1902 (19)	173 (3)
N3—H3A···O2	0.87 (2)	1.94 (2)	2.808 (2)	174.9 (19)
N2—H2B···Cl1	0.96 (2)	2.32 (2)	3.2754 (17)	170.9 (17)
N12—H12A···Cl3 ^v	0.89 (2)	2.43 (2)	3.313 (2)	171 (2)
N23—H23A···Cl3 ^{vi}	0.87 (2)	2.50 (2)	3.324 (2)	158 (2)
N13—H13B···O3 ⁱⁱ	0.92 (2)	1.97 (2)	2.883 (2)	178 (2)
N22—H22A···Cl2 ⁱⁱⁱ	0.91 (2)	2.43 (3)	3.3194 (18)	168 (2)
N13—H13A···Cl1	0.91 (2)	2.48 (2)	3.335 (2)	156.9 (19)
O3—H03B···Cl2 ^{iv}	0.84 (4)	2.45 (4)	3.286 (2)	174 (3)
O2—H0B···Cl1	0.85 (3)	2.28 (3)	3.1171 (18)	172 (2)
O2—H0A···Cl2	0.79 (3)	2.40 (3)	3.1833 (18)	175 (3)
O3—H03A···Cl3	0.94 (4)	2.25 (4)	3.1586 (18)	164 (3)

Symmetry codes: (i) $x-1, y, z$; (ii) $x, y-1, z$; (iii) $-x, -y+1, -z$; (iv) $-x+1, -y+1, -z$; (v) $-x+1, -y+1, -z+1$; (vi) $-x, -y+1, -z+1$.

supplementary materials

Fig. 1





Fig. 2



supplementary materials

Fig. 3

