

(3*R*,6*R*,12*R*,20*S*,24*S*)-20,24-Epoxy-dammarane-3,6,12,25-tetraol dihydrate

Qing-Guo Meng,^{a*} Lian-Dong Liu,^b Huan-Mei Guo,^c Yi Bi^a and Liang Wang^a

^aSchool of Pharmacy, Yantai University, Yantai 264005, People's Republic of China,

^bCollege of Chemistry, Chemical Engineering and Materials Science, Shandong Normal University, Jinan 250014, People's Republic of China, and ^cMicroscale Science Institute, Weifang University, Weifang 261061, People's Republic of China
Correspondence e-mail: mqg@ytu.edu.cn


Received 15 October 2010; accepted 10 November 2010

Key indicators: single-crystal X-ray study; $T = 298\text{ K}$; mean $\sigma(\text{C}-\text{C}) = 0.004\text{ \AA}$; R factor = 0.042; wR factor = 0.108; data-to-parameter ratio = 8.9.

The title compound, $\text{C}_{30}\text{H}_{52}\text{O}_5 \cdot 2\text{H}_2\text{O}$, was degraded from pseudoginsenoside F11 which was extracted and separated from *Panax quinquefolium saponin*. The three six-membered rings are in chair conformations. The five-membered ring is in an envelope conformation and the tetrahydrofuran ring has a conformation intermediate between half-chair and envelope. In the crystal, intermolecular $\text{O}-\text{H} \cdots \text{O}$ hydrogen bonds link molecules into a three-dimensional network. Intramolecular $\text{O}-\text{H} \cdots \text{O}$ hydrogen bonds also occur.

Related literature

For background and the medicinal properties of *Panax ginseng* and *Panax quinquefolium*, see: Iljin *et al.* (1982); Shi *et al.* (1992); Shibata *et al.* (1985); Takano *et al.* (1999); Yu *et al.* (2007).



Experimental

Crystal data

$\text{C}_{30}\text{H}_{52}\text{O}_5 \cdot 2\text{H}_2\text{O}$

$M_r = 528.75$

Orthorhombic, $P2_12_12_1$
 $a = 11.4575(15)\text{ \AA}$
 $b = 15.457(2)\text{ \AA}$
 $c = 16.726(2)\text{ \AA}$
 $V = 2962.2(7)\text{ \AA}^3$

$Z = 4$
 Mo $K\alpha$ radiation
 $\mu = 0.08\text{ mm}^{-1}$
 $T = 298\text{ K}$
 $0.51 \times 0.38 \times 0.32\text{ mm}$

Data collection

Bruker SMART CCD
diffractometer
15595 measured reflections

3095 independent reflections
2768 reflections with $I > 2\sigma(I)$
 $R_{\text{int}} = 0.032$

Refinement

$R[F^2 > 2\sigma(F^2)] = 0.042$
 $wR(F^2) = 0.108$
 $S = 1.05$
 3095 reflections
 346 parameters

6 restraints
 H-atom parameters constrained
 $\Delta\rho_{\text{max}} = 0.18\text{ e \AA}^{-3}$
 $\Delta\rho_{\text{min}} = -0.16\text{ e \AA}^{-3}$

Table 1
Hydrogen-bond geometry (\AA , $^\circ$).

$D-\text{H} \cdots A$	$D-\text{H}$	$\text{H} \cdots A$	$D \cdots A$	$D-\text{H} \cdots A$
O7—H7B \cdots O4	0.84	2.05	2.836 (3)	155
O6—H6C \cdots O5 ⁱ	0.83	2.24	2.791 (3)	124
O7—H7A \cdots O1 ⁱⁱ	0.84	2.02	2.829 (3)	160
O6—H6D \cdots O3 ⁱⁱⁱ	0.85	2.04	2.879 (3)	167
O5—H5 \cdots O7 ^{iv}	0.82	1.92	2.734 (3)	174
O4—H4 \cdots O6 ^v	0.82	2.00	2.795 (3)	164
O3—H3 \cdots O2	0.82	1.82	2.627 (2)	170
O1—H1 \cdots O3	0.82	2.14	2.938 (3)	164

Symmetry codes: (i) $x - \frac{1}{2}, -y + \frac{3}{2}, -z$; (ii) $-x + 1, y + \frac{1}{2}, -z + \frac{1}{2}$; (iii) $x - 1, y + 1, z$; (iv) $-x + \frac{3}{2}, -y + 1, z - \frac{1}{2}$; (v) $-x + 1, y - \frac{1}{2}, -z + \frac{1}{2}$.

Data collection: *SMART* (Bruker, 1997); cell refinement: *SAINT* (Bruker, 1997); data reduction: *SAINT*; program(s) used to solve structure: *SHELXS97* (Sheldrick, 2008); program(s) used to refine structure: *SHELXL97* (Sheldrick, 2008); molecular graphics: *SHELXTL* (Sheldrick, 2008); software used to prepare material for publication: *SHELXTL*.

The authors would like to thank the Shandong Provincial Natural Science Foundation, China (Y2007C138), the Scientific Research Foundation of the Higher Education Institutions of Shandong Province, China (J07WE26) and the National Natural Science Foundation of China (No.81001358) for research grants.

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: LH5154).

References

- Bruker (1997). *SMART* and *SAINT*. Bruker AXS Inc., Madison, Wisconsin, USA.
- Iljin, S. G., Mallnovskaya, G. V., Uvarova, N. I., Elyakov, G. B., Antipin, M. Yu & Struchkov, Yu. T. (1982). *Tetrahedron Lett.* **23**, 5067–5070.
- Sheldrick, G. M. (2008). *Acta Cryst. A* **64**, 112–122.
- Shi, Q., Hen, K., Jioka, T. & Miwada, Y. (1992). *J. Nat. Prod.* **55**, 1488–1497.
- Shibata, S., Tanaka, L., Shoji, L. & Saito, H. (1985). *Econ. Med. Res.* pp. 217–284.
- Takano, K., Midori, T., Eiichiro, I. & Teruo, M. (1999). *Cancer Lett.* **143**, 11–16.
- Yu, C., Fu, H., Yu, X., Han, B. & Zhu, M. (2007). *Arzneimittelforschung*, **111**, 568–572.

supplementary materials

Acta Cryst. (2010). E66, o3210 [doi:10.1107/S1600536810046362]

(3R,6R,12R,20S,24S)-20,24-Epoxydammarane-3,6,12,25-tetraol dihydrate

Q.-G. Meng, L.-D. Liu, H.-M. Guo, Y. Bi and L. Wang

Comment

Both Panax ginseng and Panax quinquefolium, belonging to Araliaceae, are well known traditional medicinal herbs. They are used as tonics and for the treatment for diseases, such as tumor and myocardial ischemia. Panax ginseng contains a number of saponins, namely ginsenoside and an oleanolic acid-type saponin in addition to the major protopanaxadiol and protopanaxatriol-type saponins (Shibata *et al.*, 1985). Panax quinquefolium contains an ocotillol-type (20S, 24R-epoxyside) saponin with high anti-tumor activity (Takano *et al.*, 1999), as well as oleanolic acid-type saponin, protopanaxadiol and protopanaxatriol-type saponins. (3R,6R,12R,20S,24S)-20,24-epoxy-dammarane-3,6,12,25-tetraol is found to possess cardioprotective effects on myocardial injury induced by isoproterenol in rats (Yu *et al.*, 2007). As part of our ongoing investigation of ocotillol-type compounds and their cardioprotective effect on myocardial injury, we report here the crystal structure of the title compound, (I).

The molecular structure of (I) is shown in Fig. 1. In the molecule, all bond lengths and angles are within normal ranges (Shi *et al.*, 1992; Iljin *et al.*, 1982). The six-membered rings A(C3/C4/C5/C6/C7/C8), B(C7/C8/C10/C11/C12/C14), and C(C12/C14/C15/C16/C17/C18) are in chair conformations that depart significantly from ideal geometry to accommodate for intramolecular non-bonded 1,3-diaxial interactions involving the methyl groups. The five-membered ring D(C13/C14/C15/C16/C17) has an envelope form with C18 as the out-of-plane atom. The tetrahydrofuran ring has a conformation intermediate between half-chair and envelope forms. In the crystal structure intermolecular O-H \cdots O hydrogen bonds link molecules into a three-dimensional network.

Experimental

Pseudoginsenoside F11 was extracted and separated from Panax quinquefolium saponin. (3R,6R,12R,20S,24S)-20,24-epoxy-dammarane-3,6,12,25-tetraol was degraded from Pseudoginsenoside F11 with sodium in glycerine and separated by flash chromatography. Finally, the crystals were dried at room temperature. Single crystals of compound (I) suitable for X-ray measurements were obtained by recrystallization from ethyl acetate at room temperature.

Refinement

H atoms bonded to C atoms were fixed geometrically and allowed to ride on their attached atoms, with C—H distances in the range 0.96–0.98 Å, and with $U_{\text{iso}}(\text{H}) = 1.2U_{\text{eq}}(\text{C})$ or $1.5U_{\text{eq}}(\text{C}_{\text{methyl}})$. The H atoms bonded to hydroxy O atoms were included with O-H = 0.82 Å and $U_{\text{iso}}(\text{H}) = 1.5U_{\text{eq}}(\text{O})$ and water H atoms were included in 'as-found' positions with $U_{\text{iso}}(\text{H}) = 0.12\text{\AA}^3$. In the absence of anomalous dispersion effects Friedel pairs were merged. The absolute configuration is based on unchanging stereochemical centers in the synthesis.

supplementary materials

Figures




Fig. 1. The molecular structure of the title compound with the atom-labeling scheme. Displacement ellipsoids are drawn at the 30% probability level. H atoms are not shown.

(3*R*,6*R*,12*R*,20*S*,24*S*)-20,24-Epoxydammarane-3,6,12,25-tetraol dihydrate

Crystal data

C ₃₀ H ₅₂ O ₅ ·2H ₂ O	<i>F</i> (000) = 1168
<i>M_r</i> = 528.75	<i>D_x</i> = 1.186 Mg m ⁻³
Orthorhombic, <i>P</i> 2 ₁ 2 ₁ 2 ₁	Mo <i>Kα</i> radiation, λ = 0.71073 Å
Hall symbol: P 2ac 2ab	Cell parameters from 5972 reflections
<i>a</i> = 11.4575 (15) Å	θ = 2.4–28.0°
<i>b</i> = 15.457 (2) Å	μ = 0.08 mm ⁻¹
<i>c</i> = 16.726 (2) Å	<i>T</i> = 298 K
<i>V</i> = 2962.2 (7) Å ³	Block, colourless
<i>Z</i> = 4	0.51 × 0.38 × 0.32 mm

Data collection

Bruker SMART CCD diffractometer	2768 reflections with $I > 2\sigma(I)$
Radiation source: fine-focus sealed tube	R_{int} = 0.032
graphite	$\theta_{\text{max}} = 25.5^\circ$, $\theta_{\text{min}} = 2.2^\circ$
φ and ω scans	$h = -13 \rightarrow 13$
15595 measured reflections	$k = -18 \rightarrow 18$
3095 independent reflections	$l = -10 \rightarrow 20$

Refinement

Refinement on F^2	Primary atom site location: structure-invariant direct methods
Least-squares matrix: full	Secondary atom site location: difference Fourier map
$R[F^2 > 2\sigma(F^2)]$ = 0.042	Hydrogen site location: inferred from neighbouring sites
$wR(F^2)$ = 0.108	H-atom parameters constrained
S = 1.05	$w = 1/[\sigma^2(F_o^2) + (0.0592P)^2 + 0.5608P]$
3095 reflections	where $P = (F_o^2 + 2F_c^2)/3$
346 parameters	$(\Delta/\sigma)_{\text{max}} < 0.001$
6 restraints	$\Delta\rho_{\text{max}} = 0.18 \text{ e \AA}^{-3}$
	$\Delta\rho_{\text{min}} = -0.16 \text{ e \AA}^{-3}$

Special details

Geometry. All e.s.d.'s (except the e.s.d. in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell e.s.d.'s are taken into account individually in the estimation of e.s.d.'s in distances, angles and torsion angles; correlations between e.s.d.'s in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell e.s.d.'s is used for estimating e.s.d.'s involving l.s. planes.

Refinement. Refinement of F^2 against ALL reflections. The weighted R -factor wR and goodness of fit S are based on F^2 , conventional R -factors R are based on F , with F set to zero for negative F^2 . The threshold expression of $F^2 > \sigma(F^2)$ is used only for calculating R -factors(gt) etc. and is not relevant to the choice of reflections for refinement. R -factors based on F^2 are statistically about twice as large as those based on F , and R -factors based on ALL data will be even larger.

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (\AA^2)

	<i>x</i>	<i>y</i>	<i>z</i>	$U_{\text{iso}}^*/U_{\text{eq}}$
C1	0.7657 (3)	0.50304 (19)	0.18423 (19)	0.0600 (8)
H1A	0.7841	0.5541	0.1541	0.090*
H1B	0.8326	0.4859	0.2149	0.090*
H1C	0.7017	0.5149	0.2196	0.090*
C2	0.6080 (2)	0.44716 (19)	0.09599 (19)	0.0546 (7)
H2A	0.6027	0.5058	0.0774	0.082*
H2B	0.5531	0.4384	0.1386	0.082*
H2C	0.5904	0.4083	0.0528	0.082*
C3	0.7318 (2)	0.42959 (16)	0.12652 (17)	0.0436 (6)
C4	0.8223 (2)	0.43365 (17)	0.05835 (17)	0.0453 (6)
H4A	0.8999	0.4327	0.0831	0.054*
C5	0.8168 (3)	0.35892 (18)	0.00041 (16)	0.0485 (7)
H5A	0.8769	0.3658	-0.0400	0.058*
H5B	0.7417	0.3588	-0.0262	0.058*
C6	0.8343 (3)	0.27323 (17)	0.04358 (15)	0.0437 (6)
H6A	0.9126	0.2716	0.0654	0.052*
H6B	0.8276	0.2265	0.0051	0.052*
C7	0.7463 (2)	0.25757 (16)	0.11176 (14)	0.0367 (5)
C8	0.7455 (2)	0.33919 (15)	0.16833 (15)	0.0370 (5)
H8	0.8237	0.3405	0.1920	0.044*
C9	0.6270 (2)	0.23805 (18)	0.07120 (16)	0.0452 (6)
H9A	0.6200	0.2718	0.0232	0.068*
H9B	0.5648	0.2527	0.1072	0.068*
H9C	0.6227	0.1777	0.0581	0.068*
C10	0.6633 (2)	0.32106 (15)	0.23917 (16)	0.0402 (6)
H10	0.5864	0.3064	0.2173	0.048*
C11	0.7058 (2)	0.24394 (16)	0.28745 (15)	0.0431 (6)
H11A	0.6510	0.2339	0.3307	0.052*
H11B	0.7802	0.2589	0.3114	0.052*
C12	0.7209 (2)	0.15891 (15)	0.24068 (14)	0.0363 (5)
C13	0.5974 (2)	0.12287 (18)	0.22259 (17)	0.0451 (6)
H13A	0.5619	0.1035	0.2714	0.068*
H13B	0.6036	0.0752	0.1860	0.068*

supplementary materials

H13C	0.5504	0.1676	0.1992	0.068*
C14	0.7921 (2)	0.17999 (15)	0.16346 (15)	0.0359 (5)
H14	0.8678	0.2000	0.1836	0.043*
C15	0.8217 (2)	0.09726 (16)	0.11639 (15)	0.0416 (6)
H15A	0.7500	0.0713	0.0970	0.050*
H15B	0.8690	0.1124	0.0704	0.050*
C16	0.8869 (2)	0.03177 (16)	0.16698 (15)	0.0397 (6)
H16	0.9597	0.0583	0.1856	0.048*
C17	0.8132 (2)	0.00865 (15)	0.23932 (14)	0.0367 (5)
H17	0.7367	-0.0091	0.2190	0.044*
C18	0.7928 (2)	0.09115 (16)	0.29063 (15)	0.0404 (6)
C19	0.9103 (3)	0.12856 (18)	0.32081 (17)	0.0514 (7)
H19A	0.8969	0.1624	0.3681	0.077*
H19B	0.9439	0.1645	0.2801	0.077*
H19C	0.9628	0.0820	0.3330	0.077*
C20	0.7324 (3)	0.05094 (18)	0.36378 (15)	0.0518 (7)
H20A	0.7330	0.0905	0.4088	0.062*
H20B	0.6523	0.0353	0.3516	0.062*
C21	0.8061 (3)	-0.02966 (19)	0.38146 (16)	0.0542 (7)
H21A	0.7589	-0.0739	0.4068	0.065*
H21B	0.8705	-0.0153	0.4167	0.065*
C22	0.8527 (2)	-0.06215 (16)	0.29903 (15)	0.0414 (6)
H22	0.9382	-0.0629	0.3008	0.050*
C23	0.8091 (2)	-0.15422 (16)	0.28113 (15)	0.0406 (6)
C24	0.8690 (3)	-0.2194 (2)	0.33484 (18)	0.0605 (8)
H24A	0.8404	-0.2763	0.3230	0.091*
H24B	0.8529	-0.2057	0.3897	0.091*
H24C	0.9517	-0.2174	0.3258	0.091*
C25	0.6758 (2)	-0.16592 (19)	0.28239 (19)	0.0530 (7)
H25A	0.6509	-0.1916	0.3325	0.064*
H25B	0.6368	-0.1106	0.2759	0.064*
C26	0.6484 (3)	-0.2245 (3)	0.2139 (2)	0.0717 (10)
H26A	0.5886	-0.1991	0.1803	0.086*
H26B	0.6202	-0.2798	0.2334	0.086*
C27	0.7600 (2)	-0.23660 (17)	0.16702 (16)	0.0441 (6)
H27	0.7902	-0.2948	0.1777	0.053*
C28	0.7516 (3)	-0.22421 (18)	0.07639 (16)	0.0462 (6)
C29	0.6604 (3)	-0.2842 (2)	0.0421 (2)	0.0706 (9)
H29A	0.5848	-0.2681	0.0619	0.106*
H29B	0.6775	-0.3426	0.0577	0.106*
H29C	0.6609	-0.2800	-0.0152	0.106*
C30	0.8687 (3)	-0.2400 (2)	0.03777 (18)	0.0629 (9)
H30A	0.8627	-0.2304	-0.0188	0.094*
H30B	0.8925	-0.2986	0.0475	0.094*
H30C	0.9253	-0.2011	0.0600	0.094*
O1	0.71201 (17)	-0.13798 (12)	0.05889 (13)	0.0524 (5)
H1	0.7640	-0.1033	0.0693	0.079*
O2	0.84059 (14)	-0.17448 (10)	0.19926 (10)	0.0380 (4)
O3	0.91541 (18)	-0.04074 (11)	0.11696 (11)	0.0491 (5)

H3	0.8984	-0.0858	0.1401	0.074*
O4	0.64852 (19)	0.39272 (12)	0.29256 (12)	0.0552 (5)
H4	0.7100	0.4020	0.3161	0.083*
O5	0.81123 (18)	0.51420 (12)	0.01735 (13)	0.0559 (5)
H5	0.8679	0.5214	-0.0118	0.084*
O6	0.1664 (2)	0.9510 (2)	0.11289 (16)	0.1016 (10)
H6D	0.0926	0.9519	0.1058	0.122*
H7A	0.4471	0.4180	0.4365	0.122*
H6C	0.1989	0.9933	0.0921	0.122*
H7B	0.5268	0.4294	0.3756	0.122*
O7	0.5037 (2)	0.44817 (18)	0.42014 (16)	0.0821 (8)

Atomic displacement parameters (\AA^2)

	U^{11}	U^{22}	U^{33}	U^{12}	U^{13}	U^{23}
C1	0.077 (2)	0.0443 (14)	0.0584 (17)	-0.0088 (15)	0.0047 (17)	0.0012 (14)
C2	0.0428 (15)	0.0536 (16)	0.0674 (19)	0.0066 (13)	0.0042 (14)	0.0095 (15)
C3	0.0435 (14)	0.0405 (13)	0.0469 (15)	0.0016 (11)	0.0028 (12)	0.0015 (12)
C4	0.0392 (14)	0.0479 (14)	0.0489 (15)	-0.0018 (11)	-0.0019 (12)	0.0084 (12)
C5	0.0513 (16)	0.0557 (15)	0.0385 (14)	0.0009 (13)	0.0068 (12)	0.0082 (12)
C6	0.0470 (15)	0.0483 (14)	0.0359 (13)	0.0034 (12)	0.0053 (12)	-0.0002 (11)
C7	0.0348 (12)	0.0427 (13)	0.0326 (12)	-0.0015 (10)	-0.0013 (11)	-0.0019 (10)
C8	0.0348 (12)	0.0408 (13)	0.0354 (12)	-0.0015 (10)	-0.0010 (11)	-0.0029 (11)
C9	0.0441 (15)	0.0518 (15)	0.0399 (14)	-0.0022 (12)	-0.0043 (12)	-0.0057 (13)
C10	0.0409 (13)	0.0398 (12)	0.0398 (14)	-0.0012 (11)	0.0048 (12)	-0.0081 (11)
C11	0.0495 (15)	0.0466 (14)	0.0334 (13)	-0.0027 (12)	0.0083 (12)	-0.0058 (11)
C12	0.0376 (13)	0.0400 (12)	0.0312 (12)	0.0008 (11)	0.0044 (10)	-0.0047 (10)
C13	0.0391 (13)	0.0484 (14)	0.0477 (15)	-0.0028 (11)	0.0068 (12)	-0.0014 (12)
C14	0.0329 (12)	0.0419 (13)	0.0329 (12)	-0.0004 (10)	0.0018 (10)	-0.0022 (10)
C15	0.0482 (15)	0.0442 (13)	0.0325 (12)	0.0013 (11)	0.0102 (12)	-0.0013 (11)
C16	0.0392 (13)	0.0406 (13)	0.0392 (13)	-0.0002 (10)	0.0073 (11)	-0.0032 (11)
C17	0.0353 (12)	0.0402 (12)	0.0346 (12)	-0.0016 (10)	0.0010 (10)	-0.0024 (10)
C18	0.0450 (13)	0.0440 (13)	0.0322 (12)	-0.0006 (11)	0.0014 (11)	-0.0038 (11)
C19	0.0578 (16)	0.0504 (15)	0.0462 (16)	-0.0017 (13)	-0.0124 (14)	-0.0044 (13)
C20	0.0724 (19)	0.0519 (15)	0.0312 (13)	0.0021 (14)	0.0074 (13)	-0.0040 (12)
C21	0.072 (2)	0.0576 (16)	0.0334 (13)	0.0012 (15)	0.0019 (14)	0.0001 (13)
C22	0.0397 (13)	0.0500 (14)	0.0345 (12)	-0.0008 (11)	-0.0013 (11)	0.0006 (11)
C23	0.0420 (13)	0.0451 (13)	0.0349 (13)	0.0025 (11)	0.0070 (11)	0.0045 (11)
C24	0.077 (2)	0.0574 (17)	0.0470 (16)	0.0101 (16)	-0.0033 (16)	0.0092 (15)
C25	0.0437 (15)	0.0515 (15)	0.0639 (19)	-0.0012 (13)	0.0115 (14)	0.0044 (15)
C26	0.0505 (17)	0.106 (3)	0.0582 (19)	-0.0272 (18)	0.0041 (16)	0.001 (2)
C27	0.0484 (15)	0.0363 (12)	0.0478 (15)	-0.0050 (11)	-0.0073 (13)	0.0052 (11)
C28	0.0482 (15)	0.0462 (14)	0.0442 (14)	0.0077 (12)	-0.0062 (13)	0.0008 (12)
C29	0.075 (2)	0.070 (2)	0.067 (2)	-0.0067 (18)	-0.0228 (19)	-0.0056 (18)
C30	0.068 (2)	0.075 (2)	0.0459 (16)	0.0217 (17)	0.0027 (15)	-0.0087 (16)
O1	0.0506 (11)	0.0523 (11)	0.0543 (11)	0.0097 (9)	-0.0071 (10)	0.0098 (10)
O2	0.0353 (8)	0.0418 (9)	0.0370 (9)	0.0006 (7)	0.0022 (8)	0.0003 (8)
O3	0.0587 (11)	0.0409 (9)	0.0476 (10)	0.0016 (9)	0.0195 (10)	-0.0005 (9)

supplementary materials

O4	0.0682 (13)	0.0476 (10)	0.0500 (12)	0.0035 (10)	0.0078 (10)	-0.0130 (9)
O5	0.0558 (13)	0.0523 (11)	0.0595 (12)	-0.0010 (9)	0.0062 (10)	0.0165 (10)
O6	0.0586 (15)	0.175 (3)	0.0714 (16)	0.0095 (18)	0.0082 (13)	0.021 (2)
O7	0.0564 (13)	0.1091 (19)	0.0807 (16)	-0.0227 (13)	0.0062 (12)	-0.0208 (16)

Geometric parameters (\AA , $^\circ$)

C1—C3	1.540 (4)	C17—C18	1.555 (3)
C1—H1A	0.9600	C17—H17	0.9800
C1—H1B	0.9600	C18—C20	1.537 (4)
C1—H1C	0.9600	C18—C19	1.549 (4)
C2—C3	1.532 (4)	C19—H19A	0.9600
C2—H2A	0.9600	C19—H19B	0.9600
C2—H2B	0.9600	C19—H19C	0.9600
C2—H2C	0.9600	C20—C21	1.534 (4)
C3—C4	1.543 (4)	C20—H20A	0.9700
C3—C8	1.570 (3)	C20—H20B	0.9700
C4—O5	1.427 (3)	C21—C22	1.561 (4)
C4—C5	1.509 (4)	C21—H21A	0.9700
C4—H4A	0.9800	C21—H21B	0.9700
C5—C6	1.522 (4)	C22—C23	1.538 (4)
C5—H5A	0.9700	C22—H22	0.9800
C5—H5B	0.9700	C23—O2	1.450 (3)
C6—C7	1.541 (3)	C23—C24	1.514 (4)
C6—H6A	0.9700	C23—C25	1.538 (4)
C6—H6B	0.9700	C24—H24A	0.9600
C7—C9	1.555 (3)	C24—H24B	0.9600
C7—C14	1.569 (3)	C24—H24C	0.9600
C7—C8	1.577 (3)	C25—C26	1.494 (5)
C8—C10	1.539 (3)	C25—H25A	0.9700
C8—H8	0.9800	C25—H25B	0.9700
C9—H9A	0.9600	C26—C27	1.511 (4)
C9—H9B	0.9600	C26—H26A	0.9700
C9—H9C	0.9600	C26—H26B	0.9700
C10—O4	1.433 (3)	C27—O2	1.437 (3)
C10—C11	1.520 (4)	C27—C28	1.531 (4)
C10—H10	0.9800	C27—H27	0.9800
C11—C12	1.539 (3)	C28—O1	1.438 (3)
C11—H11A	0.9700	C28—C30	1.509 (4)
C11—H11B	0.9700	C28—C29	1.511 (4)
C12—C13	1.550 (4)	C29—H29A	0.9600
C12—C14	1.562 (3)	C29—H29B	0.9600
C12—C18	1.573 (4)	C29—H29C	0.9600
C13—H13A	0.9600	C30—H30A	0.9600
C13—H13B	0.9600	C30—H30B	0.9600
C13—H13C	0.9600	C30—H30C	0.9600
C14—C15	1.539 (3)	O1—H1	0.8200
C14—H14	0.9800	O3—H3	0.8200
C15—C16	1.516 (4)	O4—H4	0.8200

C15—H15A	0.9700	O5—H5	0.8200
C15—H15B	0.9700	O6—H6D	0.8537
C16—O3	1.436 (3)	O6—H6C	0.8296
C16—C17	1.518 (3)	O7—H7A	0.8443
C16—H16	0.9800	O7—H7B	0.8426
C17—C22	1.549 (3)		
C3—C1—H1A	109.5	C15—C16—H16	108.5
C3—C1—H1B	109.5	C17—C16—H16	108.5
H1A—C1—H1B	109.5	C16—C17—C22	121.2 (2)
C3—C1—H1C	109.5	C16—C17—C18	109.29 (19)
H1A—C1—H1C	109.5	C22—C17—C18	105.51 (18)
H1B—C1—H1C	109.5	C16—C17—H17	106.7
C3—C2—H2A	109.5	C22—C17—H17	106.7
C3—C2—H2B	109.5	C18—C17—H17	106.7
H2A—C2—H2B	109.5	C20—C18—C19	106.4 (2)
C3—C2—H2C	109.5	C20—C18—C17	100.10 (19)
H2A—C2—H2C	109.5	C19—C18—C17	110.8 (2)
H2B—C2—H2C	109.5	C20—C18—C12	117.1 (2)
C2—C3—C1	108.2 (2)	C19—C18—C12	112.3 (2)
C2—C3—C4	111.7 (2)	C17—C18—C12	109.37 (19)
C1—C3—C4	105.3 (2)	C18—C19—H19A	109.5
C2—C3—C8	113.5 (2)	C18—C19—H19B	109.5
C1—C3—C8	110.6 (2)	H19A—C19—H19B	109.5
C4—C3—C8	107.3 (2)	C18—C19—H19C	109.5
O5—C4—C5	110.8 (2)	H19A—C19—H19C	109.5
O5—C4—C3	109.3 (2)	H19B—C19—H19C	109.5
C5—C4—C3	114.5 (2)	C21—C20—C18	103.5 (2)
O5—C4—H4A	107.3	C21—C20—H20A	111.1
C5—C4—H4A	107.3	C18—C20—H20A	111.1
C3—C4—H4A	107.3	C21—C20—H20B	111.1
C4—C5—C6	110.9 (2)	C18—C20—H20B	111.1
C4—C5—H5A	109.5	H20A—C20—H20B	109.0
C6—C5—H5A	109.5	C20—C21—C22	106.2 (2)
C4—C5—H5B	109.5	C20—C21—H21A	110.5
C6—C5—H5B	109.5	C22—C21—H21A	110.5
H5A—C5—H5B	108.1	C20—C21—H21B	110.5
C5—C6—C7	113.7 (2)	C22—C21—H21B	110.5
C5—C6—H6A	108.8	H21A—C21—H21B	108.7
C7—C6—H6A	108.8	C23—C22—C17	115.7 (2)
C5—C6—H6B	108.8	C23—C22—C21	111.0 (2)
C7—C6—H6B	108.8	C17—C22—C21	104.0 (2)
H6A—C6—H6B	107.7	C23—C22—H22	108.6
C6—C7—C9	106.4 (2)	C17—C22—H22	108.6
C6—C7—C14	108.0 (2)	C21—C22—H22	108.6
C9—C7—C14	112.7 (2)	O2—C23—C24	107.7 (2)
C6—C7—C8	108.8 (2)	O2—C23—C22	107.6 (2)
C9—C7—C8	114.3 (2)	C24—C23—C22	110.7 (2)
C14—C7—C8	106.43 (19)	O2—C23—C25	103.6 (2)
C10—C8—C3	116.3 (2)	C24—C23—C25	111.3 (2)

supplementary materials

C10—C8—C7	108.65 (19)	C22—C23—C25	115.4 (2)
C3—C8—C7	116.4 (2)	C23—C24—H24A	109.5
C10—C8—H8	104.7	C23—C24—H24B	109.5
C3—C8—H8	104.7	H24A—C24—H24B	109.5
C7—C8—H8	104.7	C23—C24—H24C	109.5
C7—C9—H9A	109.5	H24A—C24—H24C	109.5
C7—C9—H9B	109.5	H24B—C24—H24C	109.5
H9A—C9—H9B	109.5	C26—C25—C23	105.6 (2)
C7—C9—H9C	109.5	C26—C25—H25A	110.6
H9A—C9—H9C	109.5	C23—C25—H25A	110.6
H9B—C9—H9C	109.5	C26—C25—H25B	110.6
O4—C10—C11	108.2 (2)	C23—C25—H25B	110.6
O4—C10—C8	114.3 (2)	H25A—C25—H25B	108.7
C11—C10—C8	110.8 (2)	C25—C26—C27	107.1 (2)
O4—C10—H10	107.8	C25—C26—H26A	110.3
C11—C10—H10	107.8	C27—C26—H26A	110.3
C8—C10—H10	107.8	C25—C26—H26B	110.3
C10—C11—C12	115.8 (2)	C27—C26—H26B	110.3
C10—C11—H11A	108.3	H26A—C26—H26B	108.5
C12—C11—H11A	108.3	O2—C27—C26	105.5 (2)
C10—C11—H11B	108.3	O2—C27—C28	109.2 (2)
C12—C11—H11B	108.3	C26—C27—C28	116.4 (3)
H11A—C11—H11B	107.4	O2—C27—H27	108.5
C11—C12—C13	107.6 (2)	C26—C27—H27	108.5
C11—C12—C14	107.51 (19)	C28—C27—H27	108.5
C13—C12—C14	113.0 (2)	O1—C28—C30	110.1 (2)
C11—C12—C18	111.0 (2)	O1—C28—C29	105.9 (2)
C13—C12—C18	110.0 (2)	C30—C28—C29	110.6 (3)
C14—C12—C18	107.74 (19)	O1—C28—C27	109.7 (2)
C12—C13—H13A	109.5	C30—C28—C27	110.4 (2)
C12—C13—H13B	109.5	C29—C28—C27	110.1 (3)
H13A—C13—H13B	109.5	C28—C29—H29A	109.5
C12—C13—H13C	109.5	C28—C29—H29B	109.5
H13A—C13—H13C	109.5	H29A—C29—H29B	109.5
H13B—C13—H13C	109.5	C28—C29—H29C	109.5
C15—C14—C12	111.37 (19)	H29A—C29—H29C	109.5
C15—C14—C7	115.3 (2)	H29B—C29—H29C	109.5
C12—C14—C7	116.13 (19)	C28—C30—H30A	109.5
C15—C14—H14	104.1	C28—C30—H30B	109.5
C12—C14—H14	104.1	H30A—C30—H30B	109.5
C7—C14—H14	104.1	C28—C30—H30C	109.5
C16—C15—C14	112.2 (2)	H30A—C30—H30C	109.5
C16—C15—H15A	109.2	H30B—C30—H30C	109.5
C14—C15—H15A	109.2	C28—O1—H1	109.5
C16—C15—H15B	109.2	C27—O2—C23	109.80 (19)
C14—C15—H15B	109.2	C16—O3—H3	109.5
H15A—C15—H15B	107.9	C10—O4—H4	109.5
O3—C16—C15	107.9 (2)	C4—O5—H5	109.5
O3—C16—C17	114.01 (19)	H6D—O6—H6C	111.9

C15—C16—C17	109.1 (2)	H7A—O7—H7B	109.7
O3—C16—H16	108.5		
C2—C3—C4—O5	53.7 (3)	O3—C16—C17—C18	177.5 (2)
C1—C3—C4—O5	−63.5 (3)	C15—C16—C17—C18	−61.8 (3)
C8—C3—C4—O5	178.7 (2)	C16—C17—C18—C20	−173.0 (2)
C2—C3—C4—C5	−71.3 (3)	C22—C17—C18—C20	−41.2 (3)
C1—C3—C4—C5	171.5 (2)	C16—C17—C18—C19	−60.9 (3)
C8—C3—C4—C5	53.7 (3)	C22—C17—C18—C19	70.9 (2)
O5—C4—C5—C6	177.4 (2)	C16—C17—C18—C12	63.4 (3)
C3—C4—C5—C6	−58.4 (3)	C22—C17—C18—C12	−164.77 (19)
C4—C5—C6—C7	56.6 (3)	C11—C12—C18—C20	70.8 (3)
C5—C6—C7—C9	72.3 (3)	C13—C12—C18—C20	−48.2 (3)
C5—C6—C7—C14	−166.4 (2)	C14—C12—C18—C20	−171.7 (2)
C5—C6—C7—C8	−51.3 (3)	C11—C12—C18—C19	−52.8 (3)
C2—C3—C8—C10	−56.4 (3)	C13—C12—C18—C19	−171.8 (2)
C1—C3—C8—C10	65.4 (3)	C14—C12—C18—C19	64.6 (2)
C4—C3—C8—C10	179.7 (2)	C11—C12—C18—C17	−176.3 (2)
C2—C3—C8—C7	73.7 (3)	C13—C12—C18—C17	64.7 (2)
C1—C3—C8—C7	−164.5 (2)	C14—C12—C18—C17	−58.9 (2)
C4—C3—C8—C7	−50.2 (3)	C19—C18—C20—C21	−71.0 (3)
C6—C7—C8—C10	−176.5 (2)	C17—C18—C20—C21	44.4 (3)
C9—C7—C8—C10	64.7 (3)	C12—C18—C20—C21	162.5 (2)
C14—C7—C8—C10	−60.4 (2)	C18—C20—C21—C22	−32.0 (3)
C6—C7—C8—C3	49.8 (3)	C16—C17—C22—C23	−91.4 (3)
C9—C7—C8—C3	−69.0 (3)	C18—C17—C22—C23	143.9 (2)
C14—C7—C8—C3	165.9 (2)	C16—C17—C22—C21	146.6 (2)
C3—C8—C10—O4	−42.3 (3)	C18—C17—C22—C21	21.9 (3)
C7—C8—C10—O4	−176.0 (2)	C20—C21—C22—C23	−119.1 (3)
C3—C8—C10—C11	−164.9 (2)	C20—C21—C22—C17	6.0 (3)
C7—C8—C10—C11	61.4 (3)	C17—C22—C23—O2	53.1 (3)
O4—C10—C11—C12	177.1 (2)	C21—C22—C23—O2	171.4 (2)
C8—C10—C11—C12	−56.9 (3)	C17—C22—C23—C24	170.6 (2)
C10—C11—C12—C13	−73.3 (3)	C21—C22—C23—C24	−71.2 (3)
C10—C11—C12—C14	48.7 (3)	C17—C22—C23—C25	−61.9 (3)
C10—C11—C12—C18	166.3 (2)	C21—C22—C23—C25	56.3 (3)
C11—C12—C14—C15	174.7 (2)	O2—C23—C25—C26	22.3 (3)
C13—C12—C14—C15	−66.7 (3)	C24—C23—C25—C26	−93.1 (3)
C18—C12—C14—C15	55.1 (3)	C22—C23—C25—C26	139.7 (3)
C11—C12—C14—C7	−50.7 (3)	C23—C25—C26—C27	−7.6 (4)
C13—C12—C14—C7	68.0 (3)	C25—C26—C27—O2	−10.2 (3)
C18—C12—C14—C7	−170.30 (19)	C25—C26—C27—C28	−131.4 (3)
C6—C7—C14—C15	−52.5 (3)	O2—C27—C28—O1	−59.5 (3)
C9—C7—C14—C15	64.7 (3)	C26—C27—C28—O1	59.7 (3)
C8—C7—C14—C15	−169.2 (2)	O2—C27—C28—C30	61.9 (3)
C6—C7—C14—C12	174.6 (2)	C26—C27—C28—C30	−178.8 (3)
C9—C7—C14—C12	−68.1 (3)	O2—C27—C28—C29	−175.7 (2)
C8—C7—C14—C12	58.0 (3)	C26—C27—C28—C29	−56.4 (3)
C12—C14—C15—C16	−56.1 (3)	C26—C27—O2—C23	25.7 (3)
C7—C14—C15—C16	168.9 (2)	C28—C27—O2—C23	151.5 (2)

supplementary materials

C14—C15—C16—O3	−177.17 (19)	C24—C23—O2—C27	87.9 (2)
C14—C15—C16—C17	58.4 (3)	C22—C23—O2—C27	−152.78 (19)
O3—C16—C17—C22	54.6 (3)	C25—C23—O2—C27	−30.1 (2)
C15—C16—C17—C22	175.3 (2)		

Hydrogen-bond geometry (Å, °)

<i>D</i> —H··· <i>A</i>	<i>D</i> —H	H··· <i>A</i>	<i>D</i> ··· <i>A</i>	<i>D</i> —H··· <i>A</i>
O7—H7B···O4	0.84	2.05	2.836 (3)	155
O6—H6C···O5 ⁱ	0.83	2.24	2.791 (3)	124
O7—H7A···O1 ⁱⁱ	0.84	2.02	2.829 (3)	160
O6—H6D···O3 ⁱⁱⁱ	0.85	2.04	2.879 (3)	167
O5—H5···O7 ^{iv}	0.82	1.92	2.734 (3)	174
O4—H4···O6 ^v	0.82	2.00	2.795 (3)	164
O3—H3···O2	0.82	1.82	2.627 (2)	170
O1—H1···O3	0.82	2.14	2.938 (3)	164

Symmetry codes: (i) $x-1/2, -y+3/2, -z$; (ii) $-x+1, y+1/2, -z+1/2$; (iii) $x-1, y+1, z$; (iv) $-x+3/2, -y+1, z-1/2$; (v) $-x+1, y-1/2, -z+1/2$.

Fig. 1

