

1-Acetyl-2-*r*,6-*c*-bis(4-chlorophenyl)-3-methyl-1,2,5,6-tetrahydropyridin-4-yl acetate

V. Vimalraj and K. Pandiarajan*

Department of Chemistry, Annamalai University, Annamalai Nagar 608 002, Tamilnadu, India
Correspondence e-mail: profkprajan@yahoo.co.in


Received 17 October 2010; accepted 16 November 2010

Key indicators: single-crystal X-ray study; $T = 293\text{ K}$; mean $\sigma(\text{C}-\text{C}) = 0.003\text{ \AA}$; R factor = 0.037; wR factor = 0.100; data-to-parameter ratio = 21.7.

In the title compound, $\text{C}_{22}\text{H}_{21}\text{Cl}_2\text{NO}_3$, the pyridine ring adopts a half-chair conformation and the 4-chlorophenyl groups occupy axial positions. The 4-chlorophenyl groups are almost perpendicular to the plane of the tetrahydropyridine ring forming dihedral angles 84.62 (6) and 85.55 (5) $^\circ$; the dihedral angle between the two 4-chlorophenyl rings is 12.16 (4) $^\circ$. The crystal structure is stabilized by intermolecular C—H \cdots O interactions.

Related literature

For a related structure, see: Subha Nandhini *et al.* (2003).



Experimental

Crystal data

$\text{C}_{22}\text{H}_{21}\text{Cl}_2\text{NO}_3$
 $M_r = 418.30$

Monoclinic, Cc
 $a = 16.560 (3)\text{ \AA}$

$b = 14.809 (3)\text{ \AA}$
 $c = 10.241 (2)\text{ \AA}$
 $\beta = 124.27 (3)^\circ$
 $V = 2075.5 (10)\text{ \AA}^3$
 $Z = 4$

Mo $K\alpha$ radiation
 $\mu = 0.34\text{ mm}^{-1}$
 $T = 293\text{ K}$
 $0.30 \times 0.25 \times 0.20\text{ mm}$

Data collection

Bruker Kappa APEXII CCD diffractometer
Absorption correction: multi-scan (*SADABS*; Bruker, 1999)
 $T_{\min} = 0.866$, $T_{\max} = 0.936$

13520 measured reflections
5546 independent reflections
4578 reflections with $I > 2\sigma(I)$
 $R_{\text{int}} = 0.020$

Refinement

$R[F^2 > 2\sigma(F^2)] = 0.037$
 $wR(F^2) = 0.100$
 $S = 1.04$
5546 reflections
256 parameters
2 restraints

H-atom parameters constrained
 $\Delta\rho_{\max} = 0.27\text{ e \AA}^{-3}$
 $\Delta\rho_{\min} = -0.17\text{ e \AA}^{-3}$
Absolute structure: Flack (1983),
2649 Friedel pairs
Flack parameter: 0.02 (5)

Table 1
Hydrogen-bond geometry (\AA , $^\circ$).

$D-\text{H}\cdots A$	$D-\text{H}$	$\text{H}\cdots A$	$D\cdots A$	$D-\text{H}\cdots A$
$\text{C}3-\text{H}3\cdots\text{O}3^i$	0.98	2.44	3.341 (3)	152
$\text{C}4-\text{H}4B\cdots\text{O}1^{ii}$	0.97	2.35	3.308 (3)	169

Symmetry codes: (i) $x + \frac{1}{2}, -y + \frac{1}{2}, z + \frac{1}{2}$, (ii) $x, -y + 1, z + \frac{1}{2}$.

Data collection: *APEX2* (Bruker, 2004); cell refinement: *APPEX2* and *SAINT* (Bruker, 2004); data reduction: *SAINT*; program(s) used to solve structure: *SIR92* (Altomare *et al.*, 1993); program(s) used to refine structure: *SHELXL97* (Sheldrick, 2008); molecular graphics: *ORTEP-3* (Farrugia, 1997) and *Mercury* (Bruno *et al.*, 2002); software used to prepare material for publication: *PLATON* (Spek, 2009).

The authors are grateful to Dr Babu Varghese, Senior Scientist, Indian Institute of Technology Madras, for his valuable suggestions and for the data collection.

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: PV2343).

References

- Altomare, A., Cascarano, G., Giacovazzo, C. & Guagliardi, A. (1993). *J. Appl. Cryst.* **26**, 343–350.
- Bruker (1999). *SADABS*. Bruker AXS Inc., Madison, Wisconsin, USA.
- Bruker (2004). *APPEX2* and *SAINT*. Bruker AXS Inc., Madison, Wisconsin, USA.
- Bruno, I. J., Cole, J. C., Edgington, P. R., Kessler, M., Macrae, C. F., McCabe, P., Pearson, J. & Taylor, R. (2002). *Acta Cryst.* **B58**, 389–397.
- Farrugia, L. J. (1997). *J. Appl. Cryst.* **30**, 565.
- Flack, H. D. (1983). *Acta Cryst.* **A39**, 876–881.
- Sheldrick, G. M. (2008). *Acta Cryst.* **A64**, 112–122.
- Spek, A. L. (2009). *Acta Cryst.* **D65**, 148–155.
- Subha Nandhini, M., Vijayakumar, V., Mostad, A., Sundaravadivelu, M. & Natarajan, S. (2003). *Acta Cryst.* **E59**, o1672–o1674.

supplementary materials

Acta Cryst. (2010). E66, o3259 [doi:10.1107/S1600536810047586]

1-Acetyl-2-*r*,6-*c*-bis(4-chlorophenyl)-3-methyl-1,2,5,6-tetrahydropyridin-4-yl acetate

V. Vimalraj and K. Pandiarajan

Comment

The X-ray crystal structure determination of the title compound was undertaken to determine the effect of substitution of acetyl and acetoxy groups at positions 1 and 4, respectively, on the conformation of the tetrahydropyridine ring. The tetrahydropyridine ring adopts a half chair conformation with N1 and C3 atoms 0.324 (3) and -0.328 (3) Å, respectively, out of the basal plane formed by the remaining ring atoms (C4/C5/C6/C7) and the aryl groups occupy axial positions. The 4-chlorophenyl groups, C11–C16/C11 and C17–C22/C12, are almost perpendicular to the tetrahydropyridine ring forming dihedral angles 84.62 (6) and 85.55 (5)°, respectively; the dihedral angle between the two 4-chlorophenyl rings is 12.16 (4)°. The aryl groups take axial positions to avoid A1,3 strain. The acetoxy group O2/O3/C9/C10 is almost perpendicular (88.05 (6)°) to the tetrahydropyridine ring. The crystal structure is stabilized by intermolecular C—H···O interactions. The bond distances and angles in the title compound are comparable to a similar structure reported earlier (Subha Nandhini *et al.*, (2003)).

Experimental

A mixture of 3 t-methyl-2*r*,6*c*-bis(4-chlorophenyl)-piperidin-4-one (0.01 mol) and hippuric acid in acetic anhydride (20 ml) was refluxed for about 2 h. After the completion of reaction, excess of acetic anhydride was removed by distillation and water (50 ml) was added. The title compound thus obtained as a solid product was separated and colourless crystals were grown by slow evaporation method using ethanol as solvent.

Refinement

The H atoms were included in the refinement at geometrically idealized positions with C—H distances 0.93, 0.96, 0.97 and 0.99 Å for aryl, methyl, methylene and methyne type H-atoms in riding mode allowing $U_{\text{iso}}(\text{H}) = 1.5$ or $1.2 U_{\text{eq}}$ of the carrier methyl and non-methyl C-atoms, respectively.

Figures




Fig. 1. The molecular structure of the title compound, showing 50% probability displacement ellipsoids.

supplementary materials




Fig. 2. Part of the crystal structure showing the formation of the possible three C—H···O hydrogen bonds C4—H4B···O1ⁱ, C3—H3···O3ⁱⁱ and C21—H21···O1ⁱⁱⁱ [symmetry code: (i) x, -y+1, z+1/2, (ii) x+1/2, -y+1/2, z+1/2 /and (iii) x,-y+1,+z-1/2] within the unit cell.

1-Acetyl-2-r,6-c-bis(4-chlorophenyl)-3-methyl-1,2,5,6-tetrahydropyridin-4-yl acetate

Crystal data

C ₂₂ H ₂₁ Cl ₂ NO ₃	$F(000) = 872$
$M_r = 418.30$	$D_x = 1.339 \text{ Mg m}^{-3}$
Monoclinic, <i>Cc</i>	Melting point: 411 K
Hall symbol: C -2yc	Mo $K\alpha$ radiation, $\lambda = 0.71073 \text{ \AA}$
$a = 16.560 (3) \text{ \AA}$	Cell parameters from 6663 reflections
$b = 14.809 (3) \text{ \AA}$	$\theta = 2.8\text{--}59.2^\circ$
$c = 10.241 (2) \text{ \AA}$	$\mu = 0.34 \text{ mm}^{-1}$
$\beta = 124.27 (3)^\circ$	$T = 293 \text{ K}$
$V = 2075.5 (10) \text{ \AA}^3$	Block, colourless
$Z = 4$	$0.30 \times 0.25 \times 0.20 \text{ mm}$

Data collection

Bruker Kappa APEXII CCD diffractometer	5546 independent reflections
Radiation source: fine-focus sealed tube graphite	4578 reflections with $I > 2\sigma(I)$
ω and φ scans	$R_{\text{int}} = 0.020$
Absorption correction: multi-scan (<i>SADABS</i> ; Bruker, 1999)	$\theta_{\text{max}} = 29.6^\circ, \theta_{\text{min}} = 2.0^\circ$
$T_{\text{min}} = 0.866, T_{\text{max}} = 0.936$	$h = -22\text{--}22$
13520 measured reflections	$k = -20\text{--}20$
	$l = -14\text{--}14$

Refinement

Refinement on F^2	Secondary atom site location: difference Fourier map
Least-squares matrix: full	Hydrogen site location: inferred from neighbouring sites
$R[F^2 > 2\sigma(F^2)] = 0.037$	H-atom parameters constrained
$wR(F^2) = 0.100$	$w = 1/[\sigma^2(F_o^2) + (0.0528P)^2 + 0.2729P]$
$S = 1.04$	where $P = (F_o^2 + 2F_c^2)/3$
5546 reflections	$(\Delta/\sigma)_{\text{max}} = 0.011$
256 parameters	$\Delta\rho_{\text{max}} = 0.27 \text{ e \AA}^{-3}$
	$\Delta\rho_{\text{min}} = -0.17 \text{ e \AA}^{-3}$

2 restraints
 Primary atom site location: structure-invariant direct methods Absolute structure: Flack (1983), 2649 Friedel pairs
 Flack parameter: 0.02 (5)

Special details

Geometry. All e.s.d.'s (except the e.s.d. in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell e.s.d.'s are taken into account individually in the estimation of e.s.d.'s in distances, angles and torsion angles; correlations between e.s.d.'s in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell e.s.d.'s is used for estimating e.s.d.'s involving l.s. planes.

Refinement. Refinement of F^2 against ALL reflections. The weighted R -factor wR and goodness of fit S are based on F^2 , conventional R -factors R are based on F , with F set to zero for negative F^2 . The threshold expression of $F^2 > \sigma(F^2)$ is used only for calculating R -factors(gt) etc. and is not relevant to the choice of reflections for refinement. R -factors based on F^2 are statistically about twice as large as those based on F , and R -factors based on ALL data will be even larger.

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (\AA^2)

	<i>x</i>	<i>y</i>	<i>z</i>	$U_{\text{iso}}^*/U_{\text{eq}}$
Cl2	0.12844 (5)	0.18128 (4)	-0.47636 (7)	0.07210 (17)
Cl1	0.25477 (6)	-0.00598 (5)	-0.08332 (9)	0.0858 (2)
O2	0.03552 (10)	0.37134 (10)	0.21164 (17)	0.0576 (4)
N1	0.26481 (10)	0.40891 (9)	0.16833 (17)	0.0389 (3)
C2	0.33771 (12)	0.46887 (12)	0.2128 (2)	0.0478 (4)
O1	0.32636 (10)	0.53254 (10)	0.1281 (2)	0.0664 (4)
C20	0.13277 (14)	0.25613 (13)	-0.3423 (2)	0.0488 (4)
C17	0.14951 (12)	0.36840 (12)	-0.1132 (2)	0.0395 (3)
O3	-0.01433 (13)	0.24011 (13)	0.0838 (2)	0.0784 (5)
C19	0.08085 (13)	0.23746 (14)	-0.2795 (2)	0.0496 (4)
H19	0.0403	0.1871	-0.3131	0.059*
C3	0.27461 (12)	0.33000 (11)	0.2627 (2)	0.0398 (3)
H3	0.3404	0.3321	0.3604	0.048*
C7	0.16844 (11)	0.42629 (12)	0.0235 (2)	0.0398 (3)
H7	0.1680	0.4894	-0.0058	0.048*
C4	0.20259 (13)	0.33908 (12)	0.3092 (2)	0.0457 (4)
H4A	0.1924	0.2803	0.3393	0.055*
H4B	0.2301	0.3785	0.4003	0.055*
C11	0.26760 (11)	0.24311 (10)	0.17759 (19)	0.0391 (3)
C18	0.08892 (12)	0.29388 (13)	-0.1653 (2)	0.0442 (4)
H18	0.0531	0.2815	-0.1229	0.053*
C21	0.19096 (16)	0.33134 (13)	-0.2975 (3)	0.0539 (5)
H21	0.2246	0.3444	-0.3435	0.065*
C16	0.32496 (13)	0.23387 (13)	0.1197 (2)	0.0492 (4)
H16	0.3668	0.2807	0.1349	0.059*
C15	0.32233 (15)	0.15862 (14)	0.0411 (3)	0.0557 (5)
H15	0.3611	0.1545	0.0022	0.067*
C5	0.10804 (13)	0.37564 (13)	0.1803 (2)	0.0446 (4)
C12	0.20837 (14)	0.17175 (12)	0.1571 (2)	0.0462 (4)
H12	0.1699	0.1757	0.1966	0.055*

supplementary materials

C22	0.19884 (14)	0.38696 (13)	-0.1840 (2)	0.0497 (4)
H22	0.2380	0.4382	-0.1537	0.060*
C6	0.08920 (12)	0.41668 (12)	0.0530 (2)	0.0437 (4)
C14	0.26133 (15)	0.08872 (13)	0.0203 (2)	0.0518 (4)
C13	0.20498 (15)	0.09451 (14)	0.0792 (2)	0.0536 (4)
H13	0.1648	0.0467	0.0666	0.064*
C9	-0.02127 (14)	0.29704 (16)	0.1591 (3)	0.0569 (5)
C8	-0.00692 (15)	0.45758 (17)	-0.0687 (3)	0.0646 (5)
H8A	-0.0584	0.4164	-0.0930	0.097*
H8B	-0.0091	0.4697	-0.1627	0.097*
H8C	-0.0152	0.5130	-0.0289	0.097*
C1	0.43369 (16)	0.45659 (18)	0.3692 (3)	0.0773 (7)
H1A	0.4787	0.5024	0.3819	0.116*
H1B	0.4598	0.3981	0.3727	0.116*
H1C	0.4239	0.4615	0.4528	0.116*
C10	-0.0915 (2)	0.2967 (2)	0.2046 (4)	0.0833 (8)
H10A	-0.1459	0.3350	0.1341	0.125*
H10B	-0.0599	0.3187	0.3108	0.125*
H10C	-0.1143	0.2363	0.1985	0.125*

Atomic displacement parameters (\AA^2)

	U^{11}	U^{22}	U^{33}	U^{12}	U^{13}	U^{23}
Cl2	0.0966 (4)	0.0705 (3)	0.0709 (3)	-0.0064 (3)	0.0603 (3)	-0.0127 (3)
Cl1	0.1101 (5)	0.0630 (3)	0.0896 (4)	0.0059 (3)	0.0595 (4)	-0.0197 (3)
O2	0.0616 (8)	0.0641 (9)	0.0680 (9)	-0.0104 (7)	0.0490 (8)	-0.0149 (7)
N1	0.0354 (6)	0.0352 (6)	0.0391 (7)	-0.0014 (6)	0.0166 (6)	0.0038 (6)
C2	0.0403 (9)	0.0400 (9)	0.0554 (11)	-0.0051 (7)	0.0222 (9)	-0.0003 (8)
O1	0.0557 (8)	0.0515 (8)	0.0809 (11)	-0.0070 (6)	0.0317 (8)	0.0186 (8)
C20	0.0537 (10)	0.0509 (10)	0.0419 (9)	0.0050 (9)	0.0269 (9)	0.0031 (8)
C17	0.0355 (7)	0.0441 (9)	0.0363 (8)	0.0034 (7)	0.0186 (7)	0.0069 (7)
O3	0.0740 (10)	0.0830 (11)	0.0867 (12)	-0.0323 (9)	0.0504 (10)	-0.0365 (10)
C19	0.0419 (9)	0.0561 (11)	0.0468 (10)	-0.0072 (8)	0.0226 (8)	-0.0041 (8)
C3	0.0401 (8)	0.0393 (8)	0.0349 (8)	-0.0014 (7)	0.0179 (7)	0.0031 (6)
C7	0.0383 (8)	0.0390 (8)	0.0402 (9)	0.0005 (7)	0.0211 (8)	0.0046 (7)
C4	0.0556 (10)	0.0439 (9)	0.0417 (9)	-0.0039 (8)	0.0299 (9)	-0.0022 (7)
C11	0.0364 (8)	0.0363 (8)	0.0362 (8)	0.0031 (6)	0.0154 (7)	0.0054 (6)
C18	0.0396 (8)	0.0535 (10)	0.0425 (9)	-0.0022 (7)	0.0249 (8)	0.0005 (8)
C21	0.0649 (11)	0.0550 (11)	0.0584 (11)	0.0003 (9)	0.0447 (10)	0.0088 (9)
C16	0.0416 (9)	0.0492 (9)	0.0576 (11)	0.0014 (8)	0.0284 (9)	0.0054 (8)
C15	0.0552 (11)	0.0560 (11)	0.0627 (12)	0.0108 (9)	0.0372 (10)	0.0038 (9)
C5	0.0480 (9)	0.0459 (9)	0.0488 (10)	-0.0057 (7)	0.0326 (8)	-0.0114 (8)
C12	0.0522 (9)	0.0416 (9)	0.0519 (10)	0.0010 (8)	0.0335 (9)	0.0043 (8)
C22	0.0562 (10)	0.0459 (10)	0.0548 (11)	-0.0036 (8)	0.0360 (10)	0.0056 (8)
C6	0.0387 (8)	0.0449 (9)	0.0479 (10)	-0.0021 (7)	0.0246 (8)	-0.0063 (8)
C14	0.0596 (11)	0.0439 (9)	0.0473 (10)	0.0126 (8)	0.0273 (9)	0.0026 (8)
C13	0.0600 (11)	0.0434 (10)	0.0539 (11)	-0.0042 (8)	0.0299 (10)	-0.0004 (8)
C9	0.0465 (10)	0.0690 (13)	0.0547 (11)	-0.0071 (9)	0.0283 (10)	-0.0039 (10)

C8	0.0464 (10)	0.0793 (15)	0.0663 (13)	0.0130 (10)	0.0305 (10)	0.0061 (11)
C1	0.0444 (11)	0.0672 (15)	0.0794 (17)	-0.0147 (10)	0.0100 (11)	0.0094 (12)
C10	0.0653 (14)	0.105 (2)	0.100 (2)	-0.0069 (15)	0.0590 (16)	0.0042 (17)

Geometric parameters (Å, °)

Cl2—C20	1.7343 (19)	C11—C16	1.381 (2)
Cl1—C14	1.725 (2)	C18—H18	0.9300
O2—C9	1.347 (3)	C21—C22	1.369 (3)
O2—C5	1.409 (2)	C21—H21	0.9300
N1—C2	1.354 (2)	C16—C15	1.361 (3)
N1—C7	1.465 (2)	C16—H16	0.9300
N1—C3	1.466 (2)	C15—C14	1.377 (3)
C2—O1	1.222 (2)	C15—H15	0.9300
C2—C1	1.503 (3)	C5—C6	1.307 (3)
C20—C19	1.361 (2)	C12—C13	1.378 (3)
C20—C21	1.372 (3)	C12—H12	0.9300
C17—C18	1.381 (3)	C22—H22	0.9300
C17—C22	1.391 (2)	C6—C8	1.490 (3)
C17—C7	1.516 (2)	C14—C13	1.369 (3)
O3—C9	1.191 (3)	C13—H13	0.9300
C19—C18	1.381 (3)	C9—C10	1.475 (3)
C19—H19	0.9300	C8—H8A	0.9600
C3—C4	1.519 (2)	C8—H8B	0.9600
C3—C11	1.521 (2)	C8—H8C	0.9600
C3—H3	0.9800	C1—H1A	0.9600
C7—C6	1.510 (2)	C1—H1B	0.9600
C7—H7	0.9800	C1—H1C	0.9600
C4—C5	1.470 (3)	C10—H10A	0.9600
C4—H4A	0.9700	C10—H10B	0.9600
C4—H4B	0.9700	C10—H10C	0.9600
C11—C12	1.375 (2)		
C9—O2—C5	116.06 (15)	C15—C16—H16	118.9
C2—N1—C7	118.79 (14)	C11—C16—H16	118.9
C2—N1—C3	123.66 (14)	C16—C15—C14	118.94 (18)
C7—N1—C3	117.42 (13)	C16—C15—H15	120.5
O1—C2—N1	121.05 (17)	C14—C15—H15	120.5
O1—C2—C1	119.84 (18)	C6—C5—O2	119.56 (17)
N1—C2—C1	119.11 (17)	C6—C5—C4	127.06 (15)
C19—C20—C21	121.22 (17)	O2—C5—C4	113.09 (16)
C19—C20—Cl2	119.27 (15)	C11—C12—C13	121.28 (17)
C21—C20—Cl2	119.48 (14)	C11—C12—H12	119.4
C18—C17—C22	117.96 (17)	C13—C12—H12	119.4
C18—C17—C7	122.41 (14)	C21—C22—C17	121.33 (18)
C22—C17—C7	119.52 (16)	C21—C22—H22	119.3
C20—C19—C18	119.45 (17)	C17—C22—H22	119.3
C20—C19—H19	120.3	C5—C6—C8	124.46 (16)
C18—C19—H19	120.3	C5—C6—C7	119.81 (15)
N1—C3—C4	108.74 (14)	C8—C6—C7	115.73 (16)

supplementary materials

N1—C3—C11	110.61 (13)	C13—C14—C15	120.48 (18)
C4—C3—C11	115.75 (14)	C13—C14—Cl1	119.96 (17)
N1—C3—H3	107.1	C15—C14—Cl1	119.55 (15)
C4—C3—H3	107.1	C14—C13—C12	119.44 (18)
C11—C3—H3	107.1	C14—C13—H13	120.3
N1—C7—C6	110.78 (13)	C12—C13—H13	120.3
N1—C7—C17	112.11 (14)	O3—C9—O2	122.43 (18)
C6—C7—C17	112.35 (14)	O3—C9—C10	125.6 (2)
N1—C7—H7	107.1	O2—C9—C10	111.9 (2)
C6—C7—H7	107.1	C6—C8—H8A	109.5
C17—C7—H7	107.1	C6—C8—H8B	109.5
C5—C4—C3	112.22 (14)	H8A—C8—H8B	109.5
C5—C4—H4A	109.2	C6—C8—H8C	109.5
C3—C4—H4A	109.2	H8A—C8—H8C	109.5
C5—C4—H4B	109.2	H8B—C8—H8C	109.5
C3—C4—H4B	109.2	C2—C1—H1A	109.5
H4A—C4—H4B	107.9	C2—C1—H1B	109.5
C12—C11—C16	117.53 (16)	H1A—C1—H1B	109.5
C12—C11—C3	123.71 (15)	C2—C1—H1C	109.5
C16—C11—C3	118.74 (15)	H1A—C1—H1C	109.5
C17—C18—C19	120.89 (15)	H1B—C1—H1C	109.5
C17—C18—H18	119.6	C9—C10—H10A	109.5
C19—C18—H18	119.6	C9—C10—H10B	109.5
C22—C21—C20	119.07 (16)	H10A—C10—H10B	109.5
C22—C21—H21	120.5	C9—C10—H10C	109.5
C20—C21—H21	120.5	H10A—C10—H10C	109.5
C15—C16—C11	122.30 (17)	H10B—C10—H10C	109.5
C7—N1—C2—O1	-5.9 (3)	C12—C20—C21—C22	-175.81 (16)
C3—N1—C2—O1	178.46 (18)	C12—C11—C16—C15	1.7 (3)
C7—N1—C2—C1	173.89 (19)	C3—C11—C16—C15	-179.45 (18)
C3—N1—C2—C1	-1.8 (3)	C11—C16—C15—C14	-0.8 (3)
C21—C20—C19—C18	-1.8 (3)	C9—O2—C5—C6	93.2 (2)
C12—C20—C19—C18	175.93 (15)	C9—O2—C5—C4	-92.5 (2)
C2—N1—C3—C4	117.77 (18)	C3—C4—C5—C6	-15.7 (3)
C7—N1—C3—C4	-57.97 (18)	C3—C4—C5—O2	170.56 (14)
C2—N1—C3—C11	-114.08 (18)	C16—C11—C12—C13	-1.2 (3)
C7—N1—C3—C11	70.18 (18)	C3—C11—C12—C13	-179.91 (18)
C2—N1—C7—C6	-130.70 (16)	C20—C21—C22—C17	0.3 (3)
C3—N1—C7—C6	45.2 (2)	C18—C17—C22—C21	-2.6 (3)
C2—N1—C7—C17	102.91 (18)	C7—C17—C22—C21	173.71 (17)
C3—N1—C7—C17	-81.14 (17)	O2—C5—C6—C8	-2.5 (3)
C18—C17—C7—N1	103.99 (18)	C4—C5—C6—C8	-175.8 (2)
C22—C17—C7—N1	-72.1 (2)	O2—C5—C6—C7	176.44 (15)
C18—C17—C7—C6	-21.5 (2)	C4—C5—C6—C7	3.1 (3)
C22—C17—C7—C6	162.36 (16)	N1—C7—C6—C5	-16.0 (2)
N1—C3—C4—C5	39.76 (19)	C17—C7—C6—C5	110.29 (17)
C11—C3—C4—C5	-85.43 (19)	N1—C7—C6—C8	163.06 (16)
N1—C3—C11—C12	-131.11 (17)	C17—C7—C6—C8	-70.7 (2)
C4—C3—C11—C12	-6.9 (2)	C16—C15—C14—C13	-0.7 (3)

N1—C3—C11—C16	50.2 (2)	C16—C15—C14—Cl1	178.36 (16)
C4—C3—C11—C16	174.38 (15)	C15—C14—C13—C12	1.2 (3)
C22—C17—C18—C19	2.7 (3)	C11—C14—C13—C12	-177.82 (16)
C7—C17—C18—C19	-173.48 (16)	C11—C12—C13—C14	-0.3 (3)
C20—C19—C18—C17	-0.6 (3)	C5—O2—C9—O3	-3.6 (3)
C19—C20—C21—C22	1.9 (3)	C5—O2—C9—C10	177.1 (2)

Hydrogen-bond geometry (\AA , $^\circ$)

$D\text{—H}\cdots A$	$D\text{—H}$	$H\cdots A$	$D\cdots A$	$D\text{—H}\cdots A$
C3—H3 \cdots O3 ⁱ	0.98	2.44	3.341 (3)	152
C4—H4B \cdots O1 ⁱⁱ	0.97	2.35	3.308 (3)	169
C7—H7 \cdots O1	0.98	2.26	2.701 (2)	106

Symmetry codes: (i) $x+1/2, -y+1/2, z+1/2$; (ii) $x, -y+1, z+1/2$.

supplementary materials

Fig. 1




Fig. 2

