

7-Phenylsulfonyl-2,3-dihydro-7H-1,4-benzodioxino[6,7-b]carbazole

J. Kanchanadevi,^a V. Dhayalan,^b A. K. Mohanakrishnan,^b G. Anbalagan,^c G. Chakkaravarthi^d and V. Manivannan^{e*}

^aDepartment of Physics, Velammal Institute of Technology, Panchetty, Chennai 601 204, India, ^bDepartment of Organic Chemistry, University of Madras, Guindy Campus, Chennai 600 025, India, ^cDepartment of Physics, Presidency College (Autonomous), Chennai 600 005, India, ^dDepartment of Physics, CPCL Polytechnic College, Chennai 600 068, India, and ^eDepartment of Research and Development, PRIST University, Vallam, Thanjavur 613 403, Tamil Nadu, India
Correspondence e-mail: crystallography2010@gmail.com


Received 6 November 2010; accepted 15 November 2010

Key indicators: single-crystal X-ray study; $T = 295$ K; mean $\sigma(\text{C}-\text{C}) = 0.003$ Å; disorder in main residue; R factor = 0.041; wR factor = 0.124; data-to-parameter ratio = 16.6.

In the title compound, $\text{C}_{24}\text{H}_{17}\text{NO}_4\text{S}$, the phenyl ring makes a dihedral angle of $88.12(5)^\circ$ with the carbazole unit. The molecular structure is stabilized by weak intramolecular $\text{C}-\text{H}\cdots\text{O}$ interactions and the crystal packing exhibits weak intermolecular $\text{C}-\text{H}\cdots\text{O}$ and $\text{C}-\text{H}\cdots\pi$ interactions. Two C atoms of the 2,3-dihydro-1,4-dioxine fragment are disordered over two positions with site-occupancy factors of 0.718 (11) and 0.282 (11).

Related literature

For the biological activity of carbazole derivatives, see: Ramsewak *et al.* (1999); Tachibana *et al.* (2001). For the structures of closely related compounds, see: Chakkaravarthi *et al.* (2008a,b).



Experimental

Crystal data

$\text{C}_{24}\text{H}_{17}\text{NO}_4\text{S}$
 $M_r = 415.45$
Orthorhombic, $Pbca$
 $a = 13.189(5)$ Å
 $b = 16.363(6)$ Å
 $c = 18.039(5)$ Å
 $V = 3893(2)$ Å³
 $Z = 8$
Mo $K\alpha$ radiation
 $\mu = 0.20$ mm⁻¹
 $T = 295$ K
 $0.26 \times 0.22 \times 0.20$ mm

Data collection

Bruker Kappa APEXII diffractometer
Absorption correction: multi-scan (SADABS; Sheldrick, 1996)
 $T_{\min} = 0.950$, $T_{\max} = 0.961$
19551 measured reflections
4813 independent reflections
3469 reflections with $I > 2\sigma(I)$
 $R_{\text{int}} = 0.032$

Refinement

$R[F^2 > 2\sigma(F^2)] = 0.041$
 $wR(F^2) = 0.124$
 $S = 1.03$
4813 reflections
290 parameters
1 restraint
H-atom parameters constrained
 $\Delta\rho_{\max} = 0.28$ e Å⁻³
 $\Delta\rho_{\min} = -0.37$ e Å⁻³

Table 1

Hydrogen-bond geometry (Å, °).

$\text{Cg}1$, $\text{Cg}4$, $\text{Cg}5$ and $\text{Cg}7$ are the centroids of the $\text{N}1/\text{C}7/\text{C}18/\text{C}19/\text{C}24$, $\text{C}1-\text{C}6$, $\text{C}7-\text{C}9/\text{C}16-\text{C}18$ and $\text{C}19-\text{C}24$ rings, respectively.

$D-\text{H}\cdots A$	$D-\text{H}$	$\text{H}\cdots A$	$D\cdots A$	$D-\text{H}\cdots A$
$\text{C}8-\text{H}8\cdots\text{O}2$	0.93	2.42	3.005 (2)	120
$\text{C}23-\text{H}23\cdots\text{O}1$	0.93	2.32	2.908 (3)	121
$\text{C}10-\text{H}10\cdots\text{O}1^i$	0.93	2.52	3.411 (2)	161
$\text{C}12-\text{H}12A\cdots\text{Cg}1^{ii}$	0.97	2.98	3.670 (5)	129
$\text{C}12-\text{H}12B\cdots\text{Cg}7^{ii}$	0.97	2.78	3.411 (5)	124
$\text{C}13-\text{H}13A\cdots\text{Cg}5^{iii}$	0.97	2.77	3.680 (5)	157
$\text{C}20-\text{H}20\cdots\text{Cg}4^{iv}$	0.93	2.94	3.689 (2)	138
$\text{C}12A-\text{H}12C\cdots\text{Cg}5^{iii}$	0.97	2.53	3.446 (14)	158
$\text{C}12A-\text{H}12D\cdots\text{Cg}7^{ii}$	0.97	2.69	3.585 (14)	153
$\text{C}13A-\text{H}13D\cdots\text{Cg}1^{ii}$	0.97	2.92	3.585 (14)	127

Symmetry codes: (i) $x + \frac{1}{2}, -y + \frac{1}{2}, -z + 1$; (ii) $-x + 2, y + \frac{1}{2}, -z + \frac{3}{2}$; (iii) $x - \frac{1}{2}, y, -z + \frac{1}{2}$; (iv) $x, -y - \frac{1}{2}, z - \frac{1}{2}$.

Data collection: APEX2 (Bruker, 2004); cell refinement: SAINT (Bruker, 2004); data reduction: SAINT; program(s) used to solve structure: SHELXS97 (Sheldrick, 2008); program(s) used to refine structure: SHELXL97 (Sheldrick, 2008); molecular graphics: PLATON (Spek, 2009); software used to prepare material for publication: SHELXL97.

The authors wish to acknowledge DV University of Madras for the data collection.

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: GK2320).

References

- Bruker (2004). APEX2 and SAINT. Bruker AXS Inc., Madison, Wisconsin, USA.
Chakkaravarthi, G., Dhayalan, V., Mohanakrishnan, A. K. & Manivannan, V. (2008a). *Acta Cryst.* E64, o1667–o1668.
Chakkaravarthi, G., Dhayalan, V., Mohanakrishnan, A. K. & Manivannan, V. (2008b). *Acta Cryst.* E64, o1712–o1713.
Ramsewak, R. S., Nair, M. G., Strasburg, G. M., DeWitt, D. L. & Nitiss, J. L. (1999). *J. Agric. Food Chem.* 47, 444–447.

Sheldrick, G. M. (1996). *SADABS*, University of Göttingen, Germany.
Sheldrick, G. M. (2008). *Acta Cryst. A* **64**, 112–122.
Spek, A. L. (2009). *Acta Cryst. D* **65**, 148–155.

Tachibana, Y., Kikuzaki, H., Lajis, N. H. & Nakatani, N. (2001). *J. Agric. Food Chem.* **49**, 5589–5594.

supplementary materials

Acta Cryst. (2010). E66, o3264–o3265 [doi:10.1107/S1600536810047343]

7-Phenylsulfonyl-2,3-dihydro-7H-1,4-benzodioxino[6,7-*b*]carbazole

J. Kanchanadevi, V. Dhayalan, A. K. Mohanakrishnan, G. Anbalagan, G. Chakkaravarthi and V. Manivannan

Comment

Carbazole derivatives possess antioxidative (Tachibana *et al.*, 2001) and anti-inflammatory and antimutagenic (Ramsewak *et al.*, 1999) activities. The geometric parameters of the molecule (Fig. 1) agree well with the reported similar structures (Chakkaravarthi *et al.* 2008*a,b*).

The phenyl ring makes the dihedral angle of 88.12 (5)° with the carbazole ring system. The sum of the bond angles around N1 [357.29 (11)°] indicates that N1 atom is sp^2 hybridized. The molecular structure is stabilized by weak intramolecular C—H···O interactions and the crystal packing exhibits weak intermolecular C—H···O (Fig. 2) and C—H··· π interactions [C12—H12A···Cg1(2 - *x*, 1/2 + *y*, 3/2 - *z*) distance of 3.670 (5) Å; C12—H12B···Cg7 (2 - *x*, 1/2 + *y*, 3/2 - *z*) distance of 3.411 (5) Å; C13—H13A···Cg5 (1/2 + *x*, *y*, 3/2 - *z*) distance of 3.680 (5) Å; C20—H20··· Cg4 (*x*, 1/2 - *y*, 1/2 + *z*) distance of 3.689 (2) Å; C12A—H12C···Cg5 (1/2 + *x*, *y*, 3/2 - *z*) distance of 3.446 (14) Å; C12A—H12D···Cg7 (2 - *x*, 1/2 + *y*, 3/2 - *z*) distance of 3.585 (14) Å; C13A—H13D···Cg1 (2 - *x*, 1/2 + *y*, 3/2 - *z*) distance of 3.585 (14) Å; Cg1, Cg4, Cg5 and Cg7 are the centroids of the rings N1/C7/C18/C19/C24, C1—C6, C7/C8/C9/C16/C17/C18 and C19—C24, respectively].

Experimental

To a solution of diethyl-2-((2-(bromomethyl)-1-(phenylsulfonyl)-1*H*-indol-3-yl) methylene)malonate (0.5 g, 0.96 mmol) in dry DCE (15 ml), anhydrous ZnBr₂ (0.43 g, 1.90 mmol) and 1,4-benzodioxane (0.14 ml, 1.17 mmol) were added. It was stirred at room temperature for 2 h and then refluxed for 0.5 h under N₂ atmosphere. The solvent was removed and the residue was quenched with ice-water (50 ml) containing 1 ml of conc. HCl, extracted with chloroform (2 x 10 ml) and dried (Na₂SO₄). Removal of solvent followed by flash column chromatographic purification (n-hexane/ethyl acetate 99:1) led to the isolation of colourless crystal of the title compound.

Refinement

H atoms were positioned geometrically and refined using riding model with C—H = 0.93 Å and $U_{\text{iso}}(\text{H}) = 1.2U_{\text{eq}}(\text{C})$ for aromatic C—H and C—H = 0.97 Å and $U_{\text{iso}}(\text{H}) = 1.2U_{\text{eq}}(\text{C})$ for CH₂. The site occupancy factors of disordered C12 and C13 atoms refined at 0.718 (11) for major position and 0.282 (11) for minor position. The anisotropic displacement parameters of C3 and C4 were restrained with DELU in the final cycles of refinement.

Figures




Fig. 1. The molecular structure of the title compound with atomic labels and 30% probability displacement ellipsoids for non-H atoms.




Fig. 2. The crystal packing viewed down the *a* axis. Intermolecular hydrogen bonds are shown as dashed lines. H atoms not involved in hydrogen bonding have been omitted for clarity.

7-Phenylsulfonyl-2,3-dihydro-7H-1,4-benzodioxino[6,7-*b*]carbazole

Crystal data

$C_{24}H_{17}NO_4S$

$M_r = 415.45$

Orthorhombic, *Pbca*

Hall symbol: -P 2ac 2ab

$a = 13.189$ (5) Å

$b = 16.363$ (6) Å

$c = 18.039$ (5) Å

$V = 3893$ (2) Å³

$Z = 8$

$F(000) = 1728$

$D_x = 1.418$ Mg m⁻³

Mo $K\alpha$ radiation, $\lambda = 0.71073$ Å

Cell parameters from 5557 reflections

$\theta = 2.3$ – 27.8°

$\mu = 0.20$ mm⁻¹

$T = 295$ K

Block, colourless

$0.26 \times 0.22 \times 0.20$ mm

Data collection

Bruker Kappa APEXII
diffractometer

Radiation source: fine-focus sealed tube
graphite

ω and φ scans

Absorption correction: multi-scan
(*SADABS*; Sheldrick, 1996)

$T_{\min} = 0.950$, $T_{\max} = 0.961$

19551 measured reflections

4813 independent reflections

3469 reflections with $I > 2\sigma(I)$

$R_{\text{int}} = 0.032$

$\theta_{\max} = 28.3^\circ$, $\theta_{\min} = 2.3^\circ$

$h = -17 \rightarrow 17$

$k = -16 \rightarrow 21$

$l = -23 \rightarrow 17$

Refinement

Refinement on F^2	Primary atom site location: structure-invariant direct methods
Least-squares matrix: full	Secondary atom site location: difference Fourier map
$R[F^2 > 2\sigma(F^2)] = 0.041$	Hydrogen site location: inferred from neighbouring sites
$wR(F^2) = 0.124$	H-atom parameters constrained
$S = 1.03$	$w = 1/[\sigma^2(F_o^2) + (0.0637P)^2 + 0.6676P]$
4813 reflections	where $P = (F_o^2 + 2F_c^2)/3$
290 parameters	$(\Delta/\sigma)_{\max} < 0.001$
1 restraint	$\Delta\rho_{\max} = 0.28 \text{ e } \text{\AA}^{-3}$
	$\Delta\rho_{\min} = -0.36 \text{ e } \text{\AA}^{-3}$

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (\AA^2)

	x	y	z	$U_{\text{iso}}^*/U_{\text{eq}}$	Occ. (<1)
S1	0.75886 (3)	0.20738 (3)	0.50676 (2)	0.04410 (14)	
O1	0.68946 (12)	0.14774 (9)	0.48039 (8)	0.0645 (4)	
O2	0.86322 (10)	0.20175 (9)	0.48726 (7)	0.0609 (4)	
O3	1.18355 (9)	0.45554 (8)	0.71672 (8)	0.0547 (3)	
O4	1.10235 (10)	0.45229 (9)	0.86231 (7)	0.0614 (4)	
N1	0.75360 (10)	0.20284 (8)	0.59811 (8)	0.0412 (3)	
C1	0.71502 (13)	0.30481 (11)	0.48204 (9)	0.0429 (4)	
C2	0.61251 (14)	0.32135 (13)	0.48674 (10)	0.0541 (5)	
H2	0.5677	0.2819	0.5041	0.065*	
C3	0.57779 (19)	0.39686 (16)	0.46545 (12)	0.0728 (6)	
H3	0.5089	0.4087	0.4678	0.087*	
C4	0.6445 (3)	0.45467 (16)	0.44081 (14)	0.0850 (7)	
H4	0.6205	0.5057	0.4266	0.102*	
C5	0.7463 (2)	0.43862 (16)	0.43669 (15)	0.0840 (8)	
H5	0.7908	0.4788	0.4200	0.101*	
C6	0.78291 (17)	0.36299 (14)	0.45719 (12)	0.0636 (6)	
H6	0.8518	0.3514	0.4544	0.076*	
C7	0.81876 (11)	0.24936 (10)	0.64551 (8)	0.0348 (3)	
C8	0.90603 (11)	0.29052 (10)	0.62968 (8)	0.0379 (3)	
H8	0.9320	0.2914	0.5818	0.046*	
C9	0.95611 (11)	0.33194 (9)	0.68816 (8)	0.0343 (3)	
C10	1.04706 (12)	0.37570 (10)	0.67599 (9)	0.0397 (4)	
H10	1.0742	0.3780	0.6284	0.048*	
C11	1.09563 (11)	0.41443 (10)	0.73221 (9)	0.0396 (4)	
C12	1.2233 (3)	0.5044 (3)	0.7720 (3)	0.0511 (10)	0.718 (11)
H12A	1.1901	0.5573	0.7710	0.061*	0.718 (11)
H12B	1.2950	0.5128	0.7631	0.061*	0.718 (11)
C12A	1.2432 (9)	0.4744 (9)	0.7890 (7)	0.054 (3)	0.282 (11)
H12C	1.2765	0.4249	0.8057	0.064*	0.282 (11)
H12D	1.2956	0.5143	0.7780	0.064*	0.282 (11)

supplementary materials

C13	1.2089 (3)	0.4664 (3)	0.8461 (2)	0.0523 (11)	0.718 (11)
H13A	1.2450	0.4147	0.8478	0.063*	0.718 (11)
H13B	1.2375	0.5018	0.8838	0.063*	0.718 (11)
C13A	1.1788 (7)	0.5065 (9)	0.8505 (5)	0.054 (3)	0.282 (11)
H13C	1.2189	0.5128	0.8951	0.065*	0.282 (11)
H13D	1.1512	0.5595	0.8371	0.065*	0.282 (11)
C14	1.05574 (12)	0.41171 (10)	0.80512 (9)	0.0424 (4)	
C15	0.96920 (13)	0.36957 (10)	0.81897 (9)	0.0428 (4)	
H15	0.9442	0.3674	0.8671	0.051*	
C16	0.91622 (11)	0.32886 (10)	0.76166 (8)	0.0355 (3)	
C17	0.82565 (11)	0.28589 (10)	0.77520 (9)	0.0385 (3)	
H17	0.7989	0.2837	0.8229	0.046*	
C18	0.77697 (11)	0.24729 (9)	0.71804 (8)	0.0345 (3)	
C19	0.68351 (11)	0.20045 (9)	0.71441 (9)	0.0369 (3)	
C20	0.61522 (12)	0.17749 (11)	0.76943 (10)	0.0456 (4)	
H20	0.6249	0.1938	0.8183	0.055*	
C21	0.53295 (14)	0.13020 (12)	0.75009 (12)	0.0569 (5)	
H21	0.4869	0.1139	0.7862	0.068*	
C22	0.51865 (14)	0.10687 (13)	0.67716 (13)	0.0599 (5)	
H22	0.4619	0.0759	0.6650	0.072*	
C23	0.58604 (14)	0.12802 (12)	0.62164 (11)	0.0540 (5)	
H23	0.5763	0.1112	0.5729	0.065*	
C24	0.66883 (12)	0.17543 (10)	0.64165 (9)	0.0406 (4)	

Atomic displacement parameters (\AA^2)

	U^{11}	U^{22}	U^{33}	U^{12}	U^{13}	U^{23}
S1	0.0514 (2)	0.0484 (3)	0.0325 (2)	0.00302 (18)	-0.00712 (17)	-0.00640 (17)
O1	0.0849 (10)	0.0577 (9)	0.0509 (8)	-0.0113 (7)	-0.0220 (7)	-0.0092 (7)
O2	0.0564 (8)	0.0862 (11)	0.0400 (7)	0.0196 (7)	0.0003 (6)	-0.0129 (7)
O3	0.0468 (6)	0.0591 (8)	0.0582 (8)	-0.0183 (6)	0.0014 (6)	0.0065 (6)
O4	0.0638 (8)	0.0722 (9)	0.0484 (8)	-0.0254 (7)	-0.0039 (6)	-0.0135 (7)
N1	0.0462 (7)	0.0441 (8)	0.0334 (7)	-0.0060 (6)	-0.0073 (5)	0.0000 (6)
C1	0.0489 (9)	0.0498 (10)	0.0300 (8)	0.0005 (8)	-0.0015 (7)	0.0029 (7)
C2	0.0510 (10)	0.0658 (12)	0.0456 (10)	0.0058 (9)	0.0012 (8)	0.0040 (9)
C3	0.0784 (14)	0.0827 (16)	0.0572 (13)	0.0317 (11)	0.0006 (11)	0.0054 (11)
C4	0.129 (2)	0.0613 (15)	0.0647 (15)	0.0263 (12)	0.0033 (14)	0.0147 (11)
C5	0.113 (2)	0.0669 (16)	0.0722 (17)	-0.0160 (15)	0.0085 (14)	0.0251 (13)
C6	0.0626 (11)	0.0692 (14)	0.0588 (13)	-0.0094 (10)	0.0055 (10)	0.0160 (10)
C7	0.0390 (7)	0.0341 (8)	0.0313 (8)	0.0031 (6)	-0.0038 (6)	-0.0002 (6)
C8	0.0414 (8)	0.0449 (9)	0.0275 (7)	-0.0010 (7)	0.0033 (6)	0.0006 (7)
C9	0.0365 (7)	0.0359 (8)	0.0304 (8)	0.0018 (6)	0.0033 (6)	0.0012 (6)
C10	0.0427 (8)	0.0435 (9)	0.0329 (8)	-0.0023 (7)	0.0065 (6)	0.0039 (7)
C11	0.0379 (7)	0.0352 (8)	0.0456 (9)	-0.0025 (6)	0.0013 (6)	0.0051 (7)
C12	0.0400 (17)	0.047 (2)	0.066 (3)	-0.0101 (15)	-0.0075 (15)	-0.0009 (18)
C12A	0.053 (5)	0.051 (6)	0.057 (6)	-0.002 (4)	-0.019 (4)	-0.005 (5)
C13	0.0498 (17)	0.049 (2)	0.058 (2)	-0.0055 (16)	-0.0143 (15)	-0.0046 (17)
C13A	0.045 (4)	0.048 (6)	0.070 (5)	-0.003 (4)	-0.010 (3)	-0.007 (4)

C14	0.0490 (9)	0.0400 (9)	0.0383 (9)	-0.0039 (7)	-0.0035 (7)	-0.0036 (7)
C15	0.0494 (9)	0.0476 (10)	0.0314 (8)	-0.0075 (7)	0.0058 (7)	-0.0049 (7)
C16	0.0389 (7)	0.0356 (8)	0.0321 (8)	0.0010 (6)	0.0040 (6)	-0.0014 (6)
C17	0.0427 (8)	0.0420 (9)	0.0308 (8)	-0.0030 (7)	0.0075 (6)	-0.0014 (7)
C18	0.0359 (7)	0.0311 (8)	0.0364 (8)	0.0017 (6)	0.0027 (6)	0.0013 (6)
C19	0.0357 (7)	0.0312 (8)	0.0438 (9)	0.0023 (6)	-0.0007 (6)	0.0037 (7)
C20	0.0417 (8)	0.0439 (9)	0.0513 (10)	-0.0004 (7)	0.0059 (7)	0.0032 (8)
C21	0.0424 (9)	0.0562 (12)	0.0721 (13)	-0.0059 (8)	0.0055 (9)	0.0094 (10)
C22	0.0437 (9)	0.0555 (12)	0.0805 (15)	-0.0141 (8)	-0.0105 (9)	0.0095 (10)
C23	0.0534 (10)	0.0525 (11)	0.0561 (11)	-0.0090 (8)	-0.0173 (9)	0.0040 (9)
C24	0.0407 (8)	0.0345 (8)	0.0466 (10)	0.0011 (7)	-0.0075 (7)	0.0063 (7)

Geometric parameters (Å, °)

S1—O1	1.4201 (14)	C10—H10	0.9300
S1—O2	1.4237 (15)	C11—C14	1.417 (2)
S1—N1	1.6509 (15)	C12—C13	1.486 (7)
S1—C1	1.7535 (19)	C12—H12A	0.9700
O3—C11	1.3695 (19)	C12—H12B	0.9700
O3—C12	1.381 (4)	C12A—C13A	1.492 (19)
O3—C12A	1.555 (11)	C12A—H12C	0.9700
O4—C13A	1.360 (8)	C12A—H12D	0.9700
O4—C14	1.372 (2)	C13—H13A	0.9700
O4—C13	1.454 (4)	C13—H13B	0.9700
N1—C7	1.4315 (19)	C13A—H13C	0.9700
N1—C24	1.438 (2)	C13A—H13D	0.9700
C1—C2	1.381 (2)	C14—C15	1.357 (2)
C1—C6	1.382 (3)	C15—C16	1.414 (2)
C2—C3	1.373 (3)	C15—H15	0.9300
C2—H2	0.9300	C16—C17	1.407 (2)
C3—C4	1.367 (4)	C17—C18	1.369 (2)
C3—H3	0.9300	C17—H17	0.9300
C4—C5	1.370 (4)	C18—C19	1.453 (2)
C4—H4	0.9300	C19—C24	1.388 (2)
C5—C6	1.379 (3)	C19—C20	1.392 (2)
C5—H5	0.9300	C20—C21	1.378 (3)
C6—H6	0.9300	C20—H20	0.9300
C7—C8	1.364 (2)	C21—C22	1.383 (3)
C7—C18	1.420 (2)	C21—H21	0.9300
C8—C9	1.417 (2)	C22—C23	1.383 (3)
C8—H8	0.9300	C22—H22	0.9300
C9—C10	1.414 (2)	C23—C24	1.387 (2)
C9—C16	1.427 (2)	C23—H23	0.9300
C10—C11	1.357 (2)		
O1—S1—O2	119.73 (9)	C13A—C12A—O3	113.9 (9)
O1—S1—N1	106.04 (8)	C13A—C12A—H12C	108.8
O2—S1—N1	106.52 (7)	O3—C12A—H12C	108.8
O1—S1—C1	109.09 (9)	C13A—C12A—H12D	108.8
O2—S1—C1	108.35 (9)	O3—C12A—H12D	108.8

supplementary materials

N1—S1—C1	106.32 (8)	H12C—C12A—H12D	107.7
C11—O3—C12	117.3 (2)	O4—C13—C12	111.7 (4)
C11—O3—C12A	110.8 (5)	O4—C13—H13A	109.3
C13A—O4—C14	122.0 (4)	C12—C13—H13A	109.3
C14—O4—C13	111.0 (2)	O4—C13—H13B	109.3
C7—N1—C24	107.84 (13)	C12—C13—H13B	109.3
C7—N1—S1	123.16 (11)	H13A—C13—H13B	107.9
C24—N1—S1	126.29 (11)	O4—C13A—C12A	108.0 (10)
C2—C1—C6	121.28 (19)	O4—C13A—H13C	110.1
C2—C1—S1	119.03 (15)	C12A—C13A—H13C	110.1
C6—C1—S1	119.69 (15)	O4—C13A—H13D	110.1
C3—C2—C1	119.1 (2)	C12A—C13A—H13D	110.1
C3—C2—H2	120.5	H13C—C13A—H13D	108.4
C1—C2—H2	120.5	C15—C14—O4	118.97 (15)
C4—C3—C2	119.9 (2)	C15—C14—C11	119.94 (15)
C4—C3—H3	120.0	O4—C14—C11	121.08 (15)
C2—C3—H3	120.0	C14—C15—C16	121.36 (15)
C3—C4—C5	121.1 (2)	C14—C15—H15	119.3
C3—C4—H4	119.5	C16—C15—H15	119.3
C5—C4—H4	119.5	C17—C16—C15	121.85 (14)
C4—C5—C6	120.0 (2)	C17—C16—C9	119.45 (14)
C4—C5—H5	120.0	C15—C16—C9	118.70 (14)
C6—C5—H5	120.0	C18—C17—C16	119.86 (14)
C5—C6—C1	118.6 (2)	C18—C17—H17	120.1
C5—C6—H6	120.7	C16—C17—H17	120.1
C1—C6—H6	120.7	C17—C18—C7	120.05 (14)
C8—C7—C18	122.16 (14)	C17—C18—C19	132.47 (14)
C8—C7—N1	130.11 (14)	C7—C18—C19	107.48 (13)
C18—C7—N1	107.73 (13)	C24—C19—C20	120.28 (15)
C7—C8—C9	118.27 (14)	C24—C19—C18	108.45 (14)
C7—C8—H8	120.9	C20—C19—C18	131.24 (15)
C9—C8—H8	120.9	C21—C20—C19	118.73 (18)
C10—C9—C8	121.46 (14)	C21—C20—H20	120.6
C10—C9—C16	118.34 (14)	C19—C20—H20	120.6
C8—C9—C16	120.18 (14)	C20—C21—C22	120.22 (18)
C11—C10—C9	121.40 (15)	C20—C21—H21	119.9
C11—C10—H10	119.3	C22—C21—H21	119.9
C9—C10—H10	119.3	C21—C22—C23	122.16 (17)
C10—C11—O3	118.47 (15)	C21—C22—H22	118.9
C10—C11—C14	120.25 (14)	C23—C22—H22	118.9
O3—C11—C14	121.27 (15)	C22—C23—C24	117.21 (18)
O3—C12—C13	111.0 (4)	C22—C23—H23	121.4
O3—C12—H12A	109.4	C24—C23—H23	121.4
C13—C12—H12A	109.4	C23—C24—C19	121.38 (16)
O3—C12—H12B	109.4	C23—C24—N1	130.11 (16)
C13—C12—H12B	109.4	C19—C24—N1	108.42 (13)
H12A—C12—H12B	108.0		
O1—S1—N1—C7	174.92 (13)	O3—C12A—C13A—O4	55.0 (17)
O2—S1—N1—C7	46.36 (15)	C13A—O4—C14—C15	-168.7 (8)

C1—S1—N1—C7	-69.05 (14)	C13—O4—C14—C15	158.0 (3)
O1—S1—N1—C24	-26.04 (16)	C13A—O4—C14—C11	10.2 (8)
O2—S1—N1—C24	-154.61 (14)	C13—O4—C14—C11	-23.1 (3)
C1—S1—N1—C24	89.99 (15)	C10—C11—C14—C15	0.6 (3)
O1—S1—C1—C2	39.85 (17)	O3—C11—C14—C15	-179.12 (16)
O2—S1—C1—C2	171.74 (14)	C10—C11—C14—O4	-178.27 (16)
N1—S1—C1—C2	-74.10 (15)	O3—C11—C14—O4	2.0 (3)
O1—S1—C1—C6	-139.10 (16)	O4—C14—C15—C16	177.91 (16)
O2—S1—C1—C6	-7.21 (17)	C11—C14—C15—C16	-1.0 (3)
N1—S1—C1—C6	106.94 (16)	C14—C15—C16—C17	-179.13 (16)
C6—C1—C2—C3	0.7 (3)	C14—C15—C16—C9	0.7 (2)
S1—C1—C2—C3	-178.21 (16)	C10—C9—C16—C17	179.83 (15)
C1—C2—C3—C4	-0.7 (3)	C8—C9—C16—C17	-1.4 (2)
C2—C3—C4—C5	0.2 (4)	C10—C9—C16—C15	0.0 (2)
C3—C4—C5—C6	0.4 (4)	C8—C9—C16—C15	178.78 (15)
C4—C5—C6—C1	-0.3 (4)	C15—C16—C17—C18	-179.92 (15)
C2—C1—C6—C5	-0.2 (3)	C9—C16—C17—C18	0.3 (2)
S1—C1—C6—C5	178.73 (18)	C16—C17—C18—C7	1.1 (2)
C24—N1—C7—C8	-177.15 (16)	C16—C17—C18—C19	-178.74 (16)
S1—N1—C7—C8	-14.8 (2)	C8—C7—C18—C17	-1.5 (2)
C24—N1—C7—C18	2.68 (17)	N1—C7—C18—C17	178.69 (14)
S1—N1—C7—C18	165.04 (11)	C8—C7—C18—C19	178.43 (14)
C18—C7—C8—C9	0.3 (2)	N1—C7—C18—C19	-1.41 (16)
N1—C7—C8—C9	-179.87 (15)	C17—C18—C19—C24	179.46 (17)
C7—C8—C9—C10	179.83 (15)	C7—C18—C19—C24	-0.42 (17)
C7—C8—C9—C16	1.1 (2)	C17—C18—C19—C20	-2.7 (3)
C8—C9—C10—C11	-179.13 (15)	C7—C18—C19—C20	177.44 (16)
C16—C9—C10—C11	-0.4 (2)	C24—C19—C20—C21	-0.4 (2)
C9—C10—C11—O3	179.82 (14)	C18—C19—C20—C21	-178.10 (16)
C9—C10—C11—C14	0.1 (2)	C19—C20—C21—C22	-0.5 (3)
C12—O3—C11—C10	170.5 (3)	C20—C21—C22—C23	1.4 (3)
C12A—O3—C11—C10	-164.0 (6)	C21—C22—C23—C24	-1.1 (3)
C12—O3—C11—C14	-9.7 (3)	C22—C23—C24—C19	0.1 (3)
C12A—O3—C11—C14	15.8 (6)	C22—C23—C24—N1	176.05 (17)
C11—O3—C12—C13	37.2 (6)	C20—C19—C24—C23	0.7 (2)
C12A—O3—C12—C13	-42.9 (14)	C18—C19—C24—C23	178.81 (15)
C11—O3—C12A—C13A	-44.9 (14)	C20—C19—C24—N1	-176.06 (14)
C12—O3—C12A—C13A	65.6 (16)	C18—C19—C24—N1	2.09 (17)
C13A—O4—C13—C12	-66.6 (9)	C7—N1—C24—C23	-179.30 (17)
C14—O4—C13—C12	50.7 (5)	S1—N1—C24—C23	19.0 (3)
O3—C12—C13—O4	-59.1 (7)	C7—N1—C24—C19	-2.97 (17)
C14—O4—C13A—C12A	-37.8 (15)	S1—N1—C24—C19	-164.63 (12)
C13—O4—C13A—C12A	40.2 (10)		

Hydrogen-bond geometry (Å, °)

Cg1, Cg4, Cg5 and Cg7 are the centroids of the N1/C7/C18/C19/C24, C1—C6, C7—C9/C16—C18 and C19—C24 rings, respectively.

D—H...A	D—H	H...A	D...A	D—H...A
---------	-----	-------	-------	---------

supplementary materials

C6—H6···O2	0.93	2.52	2.894 (3)	104
C8—H8···O2	0.93	2.42	3.005 (2)	120
C23—H23···O1	0.93	2.32	2.908 (3)	121
C10—H10···O1 ⁱ	0.93	2.52	3.411 (2)	161
C12—H12A···Cg1 ⁱⁱ	0.97	2.98	3.670 (5)	129
C12—H12B···Cg7 ⁱⁱ	0.97	2.78	3.411 (5)	124
C13—H13A···Cg5 ⁱⁱⁱ	0.97	2.77	3.680 (5)	157
C20—H20···Cg4 ^{iv}	0.93	2.94	3.689 (2)	138
C12A—H12C···Cg5 ⁱⁱⁱ	0.97	2.53	3.446 (14)	158
C12A—H12D···Cg7 ⁱⁱ	0.97	2.69	3.585 (14)	153
C13A—H13D···Cg1 ⁱⁱ	0.97	2.92	3.585 (14)	127

Symmetry codes: (i) $x+1/2, -y+1/2, -z+1$; (ii) $-x+2, y+1/2, -z+3/2$; (iii) $x-1/2, y, -z+1/2$; (iv) $x, -y-1/2, z-1/2$.

Fig. 1




Fig. 2

