

Orthorhombic, $Pna2_1$
 $a = 8.1456(3)$ Å
 $b = 18.4472(5)$ Å
 $c = 10.3618(4)$ Å
 $V = 1557.00(9)$ Å³

$Z = 4$
Mo $K\alpha$ radiation
 $\mu = 0.23$ mm⁻¹
 $T = 173$ K
 $0.30 \times 0.25 \times 0.12$ mm

3-(4-Fluorophenylsulfonyl)-2-methyl-naphtho[1,2-*b*]furan

Hong Dae Choi,^a Pil Ja Seo,^a Byeng Wha Son^b and Uk Lee^{b*}

^aDepartment of Chemistry, Dongeui University, San 24 Kaya-dong Busanjin-gu, Busan 614-714, Republic of Korea, and ^bDepartment of Chemistry, Pukyong National University, 599-1 Daeyeon 3-dong, Nam-gu, Busan 608-737, Republic of Korea
Correspondence e-mail: uklee@pknu.ac.kr


Received 19 November 2010; accepted 19 November 2010

Key indicators: single-crystal X-ray study; $T = 173$ K; mean $\sigma(C-C) = 0.003$ Å; R factor = 0.030; wR factor = 0.075; data-to-parameter ratio = 12.5.

In the title compound, $C_{19}H_{13}FO_3S$, the 4-fluorophenyl ring makes a dihedral angle of 68.59 (5)° with the mean plane of the naphthofuran fragment. In the crystal, molecules are linked by weak intermolecular C—H···O, C—H···F and C—H···π interactions. The crystal structure also exhibits aromatic π—π interactions between the central benzene and the outer benzene rings of neighbouring molecules [centroid–centroid distance = 3.650 (3) Å].

Related literature

For the pharmacological activity of naphthofuran compounds, see: Einhorn *et al.* (1984); Hranjec *et al.* (2003); Mahadevan & Vaidya (2003). For our previous structural studies of related 3-arylsulfonyl-2-methylnaphtho[1,2-*b*]furan derivatives, see: Choi *et al.* (2008a,b).



Experimental

Crystal data

$C_{19}H_{13}FO_3S$

$M_r = 340.35$

Data collection

Bruker SMART APEXII CCD diffractometer
Absorption correction: multi-scan (*SADABS*; Bruker, 2009)
 $T_{\min} = 0.664$, $T_{\max} = 0.746$

7603 measured reflections
2719 independent reflections
2502 reflections with $I > 2\sigma(I)$
 $R_{\text{int}} = 0.026$

Refinement

$R[F^2 > 2\sigma(F^2)] = 0.030$
 $wR(F^2) = 0.075$
 $S = 1.05$
2719 reflections
218 parameters
1 restraint

H-atom parameters constrained
 $\Delta\rho_{\max} = 0.25$ e Å⁻³
 $\Delta\rho_{\min} = -0.23$ e Å⁻³
Absolute structure: Flack (1983),
830 Friedel pairs
Flack parameter: 0.09 (7)

Table 1

Hydrogen-bond geometry (Å, °).

$Cg1$ is the centroid of the C14—C19 4-fluorophenyl ring.

| $D-H \cdots A$ | $D-H$ | $H \cdots A$ | $D \cdots A$ | $D-H \cdots A$ |
|-----------------------------|-------|--------------|--------------|----------------|
| C8—H8···O3 ⁱ | 0.95 | 2.46 | 3.379 (2) | 163 |
| C15—H15···F1 ⁱⁱ | 0.95 | 2.53 | 3.150 (3) | 123 |
| C16—H16···O3 ⁱⁱⁱ | 0.95 | 2.44 | 3.380 (2) | 170 |
| C4—H4···Cg1 ^{iv} | 0.95 | 2.75 | 3.625 (3) | 154 |

Symmetry codes: (i) $-x + \frac{1}{2}, y - \frac{1}{2}, z - \frac{1}{2}$; (ii) $-x + 2, -y + 1, z + \frac{1}{2}$; (iii) $x + 1, y, z$; (iv) $-x + 1, -y + 1, z - \frac{1}{2}$.

Data collection: *APEX2* (Bruker, 2009); cell refinement: *SAINT* (Bruker, 2009); data reduction: *SAINT*; program(s) used to solve structure: *SHELXS97* (Sheldrick, 2008); program(s) used to refine structure: *SHELXL97* (Sheldrick, 2008); molecular graphics: *ORTEP-3* (Farrugia, 1997) and *DIAMOND* (Brandenburg, 1998); software used to prepare material for publication: *SHELXL97*.

This work was supported by the Blue-Bio Industry RIC at Dongeui University as an RIC programme under the Ministry of Knowledge Economy and Busan City.

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: SJ5062).

References

- Brandenburg, K. (1998). *DIAMOND*. Crystal Impact GbR, Bonn, Germany.
- Bruker (2009). *APEX2*, *SADABS* and *SAINT*. Bruker AXS Inc., Madison, Wisconsin, USA.
- Choi, H. D., Seo, P. J., Son, B. W. & Lee, U. (2008a). *Acta Cryst. E64*, o452.
- Choi, H. D., Seo, P. J., Son, B. W. & Lee, U. (2008b). *Acta Cryst. E64*, o837.
- Einhorn, J., Demerseman, P., Royer, R., Cavier, R. & Gayral, P. (1984). *Eur. J. Med. Chem.* **19**, 405–410.
- Farrugia, L. J. (1997). *J. Appl. Cryst.* **30**, 565.
- Flack, H. D. (1983). *Acta Cryst. A39*, 876–881.
- Hranjec, M., Grdisa, M., Pavelic, K., Boykin, D. W. & Karminski-Zamola, G. (2003). *Farmac.* **58**, 1319–1324.
- Mahadevan, K. M. & Vaidya, V. P. (2003). *Indian J. Pharm. Sci.* **65**, 128–134.
- Sheldrick, G. M. (2008). *Acta Cryst. A64*, 112–122.

supplementary materials

Acta Cryst. (2010). E66, o3302 [doi:10.1107/S1600536810048361]

3-(4-Fluorophenylsulfonyl)-2-methylnaphtho[1,2-*b*]furan

H. D. Choi, P. J. Seo, B. W. Son and U. Lee

Comment

Many compounds containing a naphthofuran moiety show diverse pharmacological properties such as antibacterial, antitumor and anthelmintic activities (Einhorn *et al.*, 1984, Hranjec *et al.*, 2003, Mahadevan & Vaidya, 2003). As a part of our ongoing studies of the substituent effect on the solid state structures of 3-arylsulfonyl-2-methylnaphtho[1,2-*b*]furan analogues (Choi *et al.*, 2008*a,b*), we report herein on the crystal structure of the title compound.

In the title molecule (Fig. 1), the naphthofuran unit is essentially planar, with a mean deviation of 0.008 (2) Å from the least-squares plane defined by the thirteen constituent atoms. The dihedral angle formed by the mean plane of the naphthofuran ring and the 4-fluorophenyl ring is 68.59 (5)°. The crystal packing (Fig. 2) is stabilized by weak intermolecular C–H···O and C–H···F hydrogen bonds; the first one between a benzene H atom and the oxygen of the O=S=O unit (Table 1; C8–H8···O3^j), and the second one between the 4-fluorophenyl H atom and the oxygen of the O=S=O unit (Table 1; C16–H16···O3ⁱⁱⁱ), and the third one between the 4-fluorophenyl H atom and the fluorine (Table 1; C15–H15···F1ⁱⁱ). The crystal packing (Fig. 3) is also exhibits an intermolecular C–H..π interaction between a benzene H atom and the 4-fluorophenyl ring (Table 1; C4–H4···Cg1^{iv}, Cg1 is the centroid of the C14–C19 4-fluorophenyl ring). The crystal packing (Fig. 3) is further stabilized by an aromatic π–π interaction between the central benzene and the outer benzene rings of neighbouring molecules. The Cg2···Cg3^{viii} distance is 3.650 (3) Å (Cg2 and Cg3 are the centroids of the C2/C3/C4/C5/C10/C11 benzene ring and the C5–C10 benzene ring, respectively).

Experimental

77% 3-chloroperoxybenzoic acid (493 mg, 2.2 mmol) was added in small portions to a stirred solution of 3-(4-fluorophenylsulfonyl)-2-methylnaphtho[1,2-*b*]furan (339 mg, 1.1 mmol) in dichloromethane (30 mL) at 273 K. After being stirred at room temperature for 8 h, the mixture was washed with saturated sodium bicarbonate solution and the organic layer was separated, dried over magnesium sulfate, filtered and concentrated at reduced pressure. The residue was purified by column chromatography (hexane-ethyl acetate, 4:1 v/v) to afford the title compound as a colorless solid [yield 71%, m.p. 439–440 K; R_f = 0.55 (hexane-ethyl acetate, 4:1 v/v)]. Single crystals suitable for X-ray diffraction were obtained by slow evaporation of a solution of the title compound in acetone at room temperature.

Refinement

All H atoms were positioned geometrically and refined using a riding model, with C–H = 0.95 Å for aryl and 0.98 Å for methyl H atoms. $U_{\text{iso}}(\text{H}) = 1.2U_{\text{eq}}(\text{C})$ for aryl and $1.5U_{\text{eq}}(\text{C})$ for methyl H atoms.

supplementary materials

Figures




Fig. 1. The molecular structure of the title compound with the atom numbering scheme. Displacement ellipsoids are drawn at the 50% probability level. H atoms are presented as a small spheres of arbitrary radius.




Fig. 2. A view of C–H···O and C–H···F interactions (dotted lines) in the crystal structure of the title compound. [Symmetry codes: (i) $-x + 1/2, y - 1/2, z - 1/2$; (ii) $-x + 2, -y + 1, z + 1/2$; (iii) $x + 1, y, z$; (v) $-x + 1/2, y + 1/2, z + 1/2$; (vi) $-x + 2, -y + 1, z - 1/2$; (vii) $x - 1, y, z$.]




Fig. 3. A view of C–H··· π and π – π interactions (dotted lines) in the crystal structure of the title compound. [Symmetry codes: (iv) $-x + 1, -y + 1, z - 1/2$; (viii) $x + 1/2, -y + 1/2, z$; (ix) $-x + 1, -y + 1, z + 1/2$.]

3-(4-Fluorophenylsulfonyl)-2-methylnaphtho[1,2-*b*]furan

Crystal data

| | |
|---------------------------------|---|
| $C_{19}H_{13}FO_3S$ | $F(000) = 704$ |
| $M_r = 340.35$ | $D_x = 1.452 \text{ Mg m}^{-3}$ |
| Orthorhombic, $Pna2_1$ | Mo $K\alpha$ radiation, $\lambda = 0.71073 \text{ \AA}$ |
| Hall symbol: P 2c -2n | Cell parameters from 3161 reflections |
| $a = 8.1456 (3) \text{ \AA}$ | $\theta = 2.3\text{--}27.5^\circ$ |
| $b = 18.4472 (5) \text{ \AA}$ | $\mu = 0.23 \text{ mm}^{-1}$ |
| $c = 10.3618 (4) \text{ \AA}$ | $T = 173 \text{ K}$ |
| $V = 1557.00 (9) \text{ \AA}^3$ | Block, colourless |
| $Z = 4$ | $0.30 \times 0.25 \times 0.12 \text{ mm}$ |

Data collection

| | |
|--|---|
| Bruker SMART APEXII CCD diffractometer | 2719 independent reflections |
| Radiation source: rotating anode graphite multilayer | 2502 reflections with $I > 2\sigma(I)$ |
| Detector resolution: 10.0 pixels mm^{-1} | $R_{\text{int}} = 0.026$ |
| | $\theta_{\text{max}} = 27.5^\circ, \theta_{\text{min}} = 2.2^\circ$ |

φ and ω scans $h = -10 \rightarrow 4$
 Absorption correction: multi-scan $k = -23 \rightarrow 22$
(SADABS; Bruker, 2009)
 $T_{\min} = 0.664, T_{\max} = 0.746$ $l = -10 \rightarrow 13$
 7603 measured reflections

Refinement

Refinement on F^2 Secondary atom site location: difference Fourier map
 Least-squares matrix: full Hydrogen site location: difference Fourier map
 $R[F^2 > 2\sigma(F^2)] = 0.030$ H-atom parameters constrained
 $wR(F^2) = 0.075$ $w = 1/[\sigma^2(F_o^2) + (0.0405P)^2 + 0.1905P]$
 $S = 1.05$ where $P = (F_o^2 + 2F_c^2)/3$
 2719 reflections $(\Delta/\sigma)_{\max} = 0.001$
 218 parameters $\Delta\rho_{\max} = 0.25 \text{ e \AA}^{-3}$
 1 restraint $\Delta\rho_{\min} = -0.23 \text{ e \AA}^{-3}$
 Primary atom site location: structure-invariant direct Absolute structure: Flack (1983), 830 Friedel pairs
 methods Flack parameter: 0.09 (7)

Special details

Geometry. All esds (except the esd in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell esds are taken into account individually in the estimation of esds in distances, angles and torsion angles; correlations between esds in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell esds is used for estimating esds involving l.s. planes.

Refinement. Refinement of F^2 against ALL reflections. The weighted R-factor wR and goodness of fit S are based on F^2 , conventional R-factors R are based on F, with F set to zero for negative F^2 . The threshold expression of $F^2 > 2\text{sigma}(F^2)$ is used only for calculating R-factors(gt) etc. and is not relevant to the choice of reflections for refinement. R-factors based on F^2 are statistically about twice as large as those based on F, and R- factors based on ALL data will be even larger.

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (\AA^2)

| | <i>x</i> | <i>y</i> | <i>z</i> | $U_{\text{iso}}^*/U_{\text{eq}}$ |
|----|--------------|--------------|--------------|----------------------------------|
| S1 | 0.52554 (5) | 0.41581 (2) | 0.52995 (6) | 0.02619 (12) |
| F1 | 1.07897 (19) | 0.56345 (10) | 0.29185 (19) | 0.0694 (5) |
| O1 | 0.46264 (16) | 0.22234 (6) | 0.38327 (16) | 0.0295 (3) |
| O2 | 0.57654 (19) | 0.39444 (8) | 0.65655 (16) | 0.0367 (4) |
| O3 | 0.38342 (15) | 0.46132 (7) | 0.51578 (19) | 0.0351 (4) |
| C1 | 0.4869 (2) | 0.33848 (9) | 0.4394 (2) | 0.0256 (4) |
| C2 | 0.3939 (2) | 0.33602 (9) | 0.3211 (2) | 0.0262 (4) |
| C3 | 0.3171 (2) | 0.38737 (10) | 0.2398 (2) | 0.0294 (5) |
| H3 | 0.3221 | 0.4377 | 0.2585 | 0.035* |
| C4 | 0.2358 (2) | 0.36252 (10) | 0.1336 (2) | 0.0321 (5) |
| H4 | 0.1835 | 0.3965 | 0.0784 | 0.039* |
| C5 | 0.2264 (2) | 0.28691 (11) | 0.1021 (2) | 0.0316 (5) |
| C6 | 0.1433 (3) | 0.26263 (12) | -0.0094 (2) | 0.0399 (6) |

supplementary materials

| | | | | |
|------|------------|--------------|-------------|------------|
| H6 | 0.0923 | 0.2967 | -0.0653 | 0.048* |
| C7 | 0.1359 (3) | 0.18989 (13) | -0.0374 (3) | 0.0475 (6) |
| H7 | 0.0807 | 0.1740 | -0.1130 | 0.057* |
| C8 | 0.2087 (3) | 0.13917 (11) | 0.0444 (3) | 0.0447 (6) |
| H8 | 0.2016 | 0.0891 | 0.0240 | 0.054* |
| C9 | 0.2901 (3) | 0.16034 (10) | 0.1534 (3) | 0.0371 (5) |
| H9 | 0.3390 | 0.1253 | 0.2084 | 0.044* |
| C10 | 0.3008 (2) | 0.23490 (9) | 0.1835 (2) | 0.0294 (5) |
| C11 | 0.3834 (2) | 0.26340 (9) | 0.2913 (2) | 0.0276 (4) |
| C12 | 0.5250 (2) | 0.26895 (10) | 0.4729 (2) | 0.0286 (4) |
| C13 | 0.6087 (3) | 0.23448 (11) | 0.5836 (3) | 0.0382 (5) |
| H13A | 0.6786 | 0.1949 | 0.5524 | 0.057* |
| H13B | 0.5266 | 0.2151 | 0.6435 | 0.057* |
| H13C | 0.6764 | 0.2706 | 0.6281 | 0.057* |
| C14 | 0.6915 (2) | 0.45978 (9) | 0.4545 (2) | 0.0251 (4) |
| C15 | 0.8504 (2) | 0.44194 (10) | 0.4931 (2) | 0.0304 (5) |
| H15 | 0.8679 | 0.4059 | 0.5572 | 0.036* |
| C16 | 0.9823 (3) | 0.47724 (12) | 0.4372 (3) | 0.0386 (5) |
| H16 | 1.0917 | 0.4664 | 0.4624 | 0.046* |
| C17 | 0.9511 (3) | 0.52791 (13) | 0.3451 (3) | 0.0423 (6) |
| C18 | 0.7955 (3) | 0.54585 (13) | 0.3041 (3) | 0.0455 (6) |
| H18 | 0.7794 | 0.5813 | 0.2388 | 0.055* |
| C19 | 0.6633 (3) | 0.51109 (11) | 0.3605 (2) | 0.0359 (5) |
| H19 | 0.5544 | 0.5224 | 0.3346 | 0.043* |

Atomic displacement parameters (\AA^2)

| | U^{11} | U^{22} | U^{33} | U^{12} | U^{13} | U^{23} |
|-----|-------------|--------------|-------------|--------------|--------------|--------------|
| S1 | 0.0226 (2) | 0.02804 (19) | 0.0279 (3) | 0.00002 (15) | 0.0020 (3) | -0.0023 (2) |
| F1 | 0.0527 (9) | 0.1072 (12) | 0.0483 (10) | -0.0428 (9) | 0.0082 (9) | 0.0102 (10) |
| O1 | 0.0281 (7) | 0.0237 (6) | 0.0368 (9) | 0.0001 (5) | 0.0003 (6) | 0.0019 (6) |
| O2 | 0.0381 (8) | 0.0429 (7) | 0.0291 (9) | -0.0014 (6) | 0.0014 (8) | 0.0032 (7) |
| O3 | 0.0239 (6) | 0.0338 (6) | 0.0477 (11) | 0.0044 (5) | 0.0033 (8) | -0.0078 (8) |
| C1 | 0.0223 (8) | 0.0251 (8) | 0.0293 (12) | -0.0009 (6) | 0.0016 (9) | -0.0008 (9) |
| C2 | 0.0229 (9) | 0.0256 (8) | 0.0301 (12) | -0.0012 (7) | 0.0028 (9) | -0.0025 (8) |
| C3 | 0.0295 (10) | 0.0241 (8) | 0.0344 (13) | -0.0008 (7) | 0.0016 (9) | -0.0006 (8) |
| C4 | 0.0329 (10) | 0.0325 (8) | 0.0310 (13) | 0.0016 (8) | 0.0001 (10) | 0.0035 (9) |
| C5 | 0.0257 (10) | 0.0358 (10) | 0.0332 (12) | -0.0049 (8) | 0.0040 (9) | -0.0025 (9) |
| C6 | 0.0381 (11) | 0.0440 (12) | 0.0375 (14) | -0.0043 (9) | 0.0003 (10) | -0.0036 (10) |
| C7 | 0.0465 (13) | 0.0545 (14) | 0.0416 (15) | -0.0139 (10) | 0.0016 (13) | -0.0153 (12) |
| C8 | 0.0481 (12) | 0.0354 (9) | 0.0506 (17) | -0.0112 (9) | 0.0099 (14) | -0.0125 (12) |
| C9 | 0.0361 (11) | 0.0291 (9) | 0.0459 (15) | -0.0062 (8) | 0.0074 (11) | -0.0035 (10) |
| C10 | 0.0263 (9) | 0.0275 (9) | 0.0346 (12) | -0.0032 (7) | 0.0077 (9) | -0.0040 (8) |
| C11 | 0.0239 (9) | 0.0252 (8) | 0.0338 (11) | -0.0008 (7) | 0.0039 (9) | 0.0013 (8) |
| C12 | 0.0227 (9) | 0.0297 (9) | 0.0335 (12) | -0.0007 (7) | 0.0036 (9) | 0.0019 (9) |
| C13 | 0.0352 (11) | 0.0347 (10) | 0.0448 (15) | 0.0048 (9) | -0.0024 (11) | 0.0075 (10) |
| C14 | 0.0219 (8) | 0.0271 (8) | 0.0263 (11) | -0.0021 (7) | 0.0002 (8) | -0.0029 (8) |
| C15 | 0.0263 (9) | 0.0306 (9) | 0.0343 (13) | 0.0027 (7) | -0.0029 (8) | -0.0025 (8) |

| | | | | | | |
|-----|-------------|-------------|-------------|--------------|--------------|--------------|
| C16 | 0.0252 (9) | 0.0505 (12) | 0.0400 (15) | -0.0028 (8) | -0.0008 (10) | -0.0093 (12) |
| C17 | 0.0389 (12) | 0.0574 (13) | 0.0307 (14) | -0.0206 (10) | 0.0064 (11) | -0.0057 (12) |
| C18 | 0.0508 (14) | 0.0525 (12) | 0.0333 (14) | -0.0185 (11) | -0.0082 (12) | 0.0124 (12) |
| C19 | 0.0355 (11) | 0.0384 (9) | 0.0339 (13) | -0.0074 (8) | -0.0086 (10) | 0.0057 (10) |

Geometric parameters (\AA , $^\circ$)

| | | | |
|------------|-------------|---------------|-------------|
| S1—O2 | 1.4313 (18) | C7—H7 | 0.9500 |
| S1—O3 | 1.4375 (13) | C8—C9 | 1.366 (4) |
| S1—C1 | 1.736 (2) | C8—H8 | 0.9500 |
| S1—C14 | 1.7596 (19) | C9—C10 | 1.413 (2) |
| F1—C17 | 1.348 (2) | C9—H9 | 0.9500 |
| O1—C12 | 1.364 (3) | C10—C11 | 1.406 (3) |
| O1—C11 | 1.378 (3) | C12—C13 | 1.478 (3) |
| C1—C12 | 1.365 (3) | C13—H13A | 0.9800 |
| C1—C2 | 1.442 (3) | C13—H13B | 0.9800 |
| C2—C11 | 1.377 (2) | C13—H13C | 0.9800 |
| C2—C3 | 1.414 (3) | C14—C19 | 1.378 (3) |
| C3—C4 | 1.364 (3) | C14—C15 | 1.394 (3) |
| C3—H3 | 0.9500 | C15—C16 | 1.384 (3) |
| C4—C5 | 1.434 (3) | C15—H15 | 0.9500 |
| C4—H4 | 0.9500 | C16—C17 | 1.360 (4) |
| C5—C6 | 1.412 (3) | C16—H16 | 0.9500 |
| C5—C10 | 1.414 (3) | C17—C18 | 1.377 (3) |
| C6—C7 | 1.374 (3) | C18—C19 | 1.382 (3) |
| C6—H6 | 0.9500 | C18—H18 | 0.9500 |
| C7—C8 | 1.395 (4) | C19—H19 | 0.9500 |
| O2—S1—O3 | 119.22 (11) | C11—C10—C9 | 124.7 (2) |
| O2—S1—C1 | 108.75 (9) | C11—C10—C5 | 115.16 (16) |
| O3—S1—C1 | 106.18 (9) | C9—C10—C5 | 120.2 (2) |
| O2—S1—C14 | 108.11 (10) | C2—C11—O1 | 110.52 (19) |
| O3—S1—C14 | 107.71 (9) | C2—C11—C10 | 124.85 (19) |
| C1—S1—C14 | 106.15 (10) | O1—C11—C10 | 124.62 (15) |
| C12—O1—C11 | 107.39 (14) | O1—C12—C1 | 109.55 (19) |
| C12—C1—C2 | 107.83 (17) | O1—C12—C13 | 115.41 (16) |
| C12—C1—S1 | 126.42 (18) | C1—C12—C13 | 135.0 (2) |
| C2—C1—S1 | 125.49 (14) | C12—C13—H13A | 109.5 |
| C11—C2—C3 | 119.37 (19) | C12—C13—H13B | 109.5 |
| C11—C2—C1 | 104.71 (18) | H13A—C13—H13B | 109.5 |
| C3—C2—C1 | 135.91 (17) | C12—C13—H13C | 109.5 |
| C4—C3—C2 | 118.08 (16) | H13A—C13—H13C | 109.5 |
| C4—C3—H3 | 121.0 | H13B—C13—H13C | 109.5 |
| C2—C3—H3 | 121.0 | C19—C14—C15 | 121.31 (19) |
| C3—C4—C5 | 122.4 (2) | C19—C14—S1 | 120.18 (15) |
| C3—C4—H4 | 118.8 | C15—C14—S1 | 118.50 (16) |
| C5—C4—H4 | 118.8 | C16—C15—C14 | 119.3 (2) |
| C6—C5—C10 | 118.56 (18) | C16—C15—H15 | 120.3 |
| C6—C5—C4 | 121.3 (2) | C14—C15—H15 | 120.3 |
| C10—C5—C4 | 120.1 (2) | C17—C16—C15 | 118.2 (2) |

supplementary materials

| | | | |
|---------------|--------------|-----------------|--------------|
| C7—C6—C5 | 120.2 (2) | C17—C16—H16 | 120.9 |
| C7—C6—H6 | 119.9 | C15—C16—H16 | 120.9 |
| C5—C6—H6 | 119.9 | F1—C17—C16 | 118.5 (2) |
| C6—C7—C8 | 120.6 (2) | F1—C17—C18 | 117.9 (2) |
| C6—C7—H7 | 119.7 | C16—C17—C18 | 123.6 (2) |
| C8—C7—H7 | 119.7 | C17—C18—C19 | 118.4 (2) |
| C9—C8—C7 | 121.1 (2) | C17—C18—H18 | 120.8 |
| C9—C8—H8 | 119.4 | C19—C18—H18 | 120.8 |
| C7—C8—H8 | 119.4 | C14—C19—C18 | 119.2 (2) |
| C8—C9—C10 | 119.4 (2) | C14—C19—H19 | 120.4 |
| C8—C9—H9 | 120.3 | C18—C19—H19 | 120.4 |
| C10—C9—H9 | 120.3 | | |
| O2—S1—C1—C12 | -10.6 (2) | C1—C2—C11—C10 | 179.16 (19) |
| O3—S1—C1—C12 | -140.05 (17) | C12—O1—C11—C2 | -0.1 (2) |
| C14—S1—C1—C12 | 105.50 (18) | C12—O1—C11—C10 | -179.04 (19) |
| O2—S1—C1—C2 | 162.76 (16) | C9—C10—C11—C2 | -179.9 (2) |
| O3—S1—C1—C2 | 33.3 (2) | C5—C10—C11—C2 | 0.6 (3) |
| C14—S1—C1—C2 | -81.12 (18) | C9—C10—C11—O1 | -1.1 (3) |
| C12—C1—C2—C11 | -0.3 (2) | C5—C10—C11—O1 | 179.34 (18) |
| S1—C1—C2—C11 | -174.67 (15) | C11—O1—C12—C1 | -0.1 (2) |
| C12—C1—C2—C3 | 178.4 (2) | C11—O1—C12—C13 | 177.31 (18) |
| S1—C1—C2—C3 | 4.0 (3) | C2—C1—C12—O1 | 0.2 (2) |
| C11—C2—C3—C4 | -0.4 (3) | S1—C1—C12—O1 | 174.55 (14) |
| C1—C2—C3—C4 | -178.9 (2) | C2—C1—C12—C13 | -176.4 (2) |
| C2—C3—C4—C5 | -0.2 (3) | S1—C1—C12—C13 | -2.1 (3) |
| C3—C4—C5—C6 | -179.1 (2) | O2—S1—C14—C19 | -152.63 (17) |
| C3—C4—C5—C10 | 1.0 (3) | O3—S1—C14—C19 | -22.6 (2) |
| C10—C5—C6—C7 | -0.1 (3) | C1—S1—C14—C19 | 90.83 (18) |
| C4—C5—C6—C7 | -180.0 (2) | O2—S1—C14—C15 | 26.78 (19) |
| C5—C6—C7—C8 | 0.6 (4) | O3—S1—C14—C15 | 156.85 (16) |
| C6—C7—C8—C9 | -0.5 (4) | C1—S1—C14—C15 | -89.76 (18) |
| C7—C8—C9—C10 | -0.1 (3) | C19—C14—C15—C16 | 1.0 (3) |
| C8—C9—C10—C11 | -178.9 (2) | S1—C14—C15—C16 | -178.42 (17) |
| C8—C9—C10—C5 | 0.7 (3) | C14—C15—C16—C17 | -0.6 (3) |
| C6—C5—C10—C11 | 179.01 (19) | C15—C16—C17—F1 | 178.8 (2) |
| C4—C5—C10—C11 | -1.1 (3) | C15—C16—C17—C18 | -0.2 (4) |
| C6—C5—C10—C9 | -0.6 (3) | F1—C17—C18—C19 | -178.3 (2) |
| C4—C5—C10—C9 | 179.30 (19) | C16—C17—C18—C19 | 0.7 (4) |
| C3—C2—C11—O1 | -178.73 (17) | C15—C14—C19—C18 | -0.5 (3) |
| C1—C2—C11—O1 | 0.2 (2) | S1—C14—C19—C18 | 178.88 (18) |
| C3—C2—C11—C10 | 0.2 (3) | C17—C18—C19—C14 | -0.3 (4) |

Hydrogen-bond geometry (\AA , $^\circ$)

Cg1 is the centroid of the C14—C19 4-fluorophenyl ring.

| $D—H\cdots A$ | $D—H$ | $H\cdots A$ | $D\cdots A$ | $D—H\cdots A$ |
|-----------------------------------|-------|-------------|-------------|---------------|
| C8—H8 \cdots O3 ⁱ | 0.95 | 2.46 | 3.379 (2) | 163 |
| C15—H15 \cdots F1 ⁱⁱ | 0.95 | 2.53 | 3.150 (3) | 123 |

supplementary materials

| | | | | |
|-----------------------------|------|------|-----------|-----|
| C16—H16···O3 ⁱⁱⁱ | 0.95 | 2.44 | 3.380 (2) | 170 |
| C4—H4···Cg1 ^{iv} | 0.95 | 2.75 | 3.625 (3) | 154 |

Symmetry codes: (i) $-x+1/2, y-1/2, z-1/2$; (ii) $-x+2, -y+1, z+1/2$; (iii) $x+1, y, z$; (iv) $-x+1, -y+1, z-1/2$.

supplementary materials

Fig. 1





Fig. 2



supplementary materials

Fig. 3

