

1,4-Dibenzylpiperazine

Meng Zhang, Yong-hong Zhou,* Li-hong Hu and Xiao-hui Yang

Institute of Chemical Industry of Forest Products, Chinese Academy of Forestry, Nanjing 210042, People's Republic of China, and Jiangsu Qiang Lin Bio-Energy Co. Ltd, Liyang 213364, People's Republic of China
Correspondence e-mail: zm205@sohu.com


Received 3 November 2010; accepted 24 November 2010

Key indicators: single-crystal X-ray study; $T = 293\text{ K}$; mean $\sigma(\text{C}-\text{C}) = 0.003\text{ \AA}$; R factor = 0.052; wR factor = 0.141; data-to-parameter ratio = 15.3.

In the title compound, $\text{C}_{18}\text{H}_{22}\text{N}_2$, which possesses non-crystallographic inversion symmetry, the central piperazine ring adopts a chair conformation. The phenyl rings are not exactly parallel and make a dihedral angle of $1.3(1)^\circ$. No significant intermolecular contacts are observed in the crystal.

Related literature

For the properties and applications of piperazine derivatives, see: Zhao *et al.* (2002); Sonurlikar *et al.* (1977); Bigoli *et al.* (2001). For the synthesis of related compounds, see: Zheng *et al.* (2005); Sarangarajan *et al.* (2005). For related structures, see: Yogavel *et al.* (2003); Gunasekaran *et al.* (1996); Thirumurugan *et al.* (1998).



Experimental

Crystal data

$\text{C}_{18}\text{H}_{22}\text{N}_2$	$V = 3070.3(11)\text{ \AA}^3$
$M_r = 266.38$	$Z = 8$
Orthorhombic, $Pbca$	Mo $K\alpha$ radiation
$a = 7.5130(15)\text{ \AA}$	$\mu = 0.07\text{ mm}^{-1}$
$b = 19.127(4)\text{ \AA}$	$T = 293\text{ K}$
$c = 21.366(4)\text{ \AA}$	$0.30 \times 0.20 \times 0.10\text{ mm}$

Data collection

Enraf–Nonius CAD-4 diffractometer	2781 independent reflections
Absorption correction: ψ scan (North <i>et al.</i> , 1968)	1650 reflections with $I > 2\sigma(I)$
$T_{\min} = 0.980$, $T_{\max} = 0.993$	$R_{\text{int}} = 0.045$
5468 measured reflections	3 standard reflections every 200 reflections
	intensity decay: 1%

Refinement

$R[F^2 > 2\sigma(F^2)] = 0.052$	182 parameters
$wR(F^2) = 0.141$	H-atom parameters constrained
$S = 1.01$	$\Delta\rho_{\max} = 0.16\text{ e \AA}^{-3}$
2781 reflections	$\Delta\rho_{\min} = -0.13\text{ e \AA}^{-3}$

Data collection: *CAD-4 EXPRESS* (Enraf–Nonius, 1989); cell refinement: *CAD-4 EXPRESS*; data reduction: *XCAD4* (Harms & Wocadlo, 1995); program(s) used to solve structure: *SHELXS97* (Sheldrick, 2008); program(s) used to refine structure: *SHELXL97* (Sheldrick, 2008); molecular graphics: *SHELXTL* (Sheldrick, 2008); software used to prepare material for publication: *SHELXTL*.

This work was supported by the 948 program of the State Forestry Administration (2009–4–55).

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: BH2322).

References

- Bigoli, F., Chen, C. T., Wu, W. C., Deplano, P., Mercuri, M. L., Pellinghelli, M. A., Pilia, L., Pintus, G., Serpe, A. & Trogu, E. F. (2001). *Chem. Commun.* **21**, 2246–2247.
Enraf–Nonius (1989). *CAD-4 EXPRESS*. Enraf–Nonius, Delft, The Netherlands.
Gunasekaran, K., Govindasamy, L., Shanmuga Sundara Raj, S., Velmurugan, D., Karunakaran, S. & Kandaswamy, M. (1996). *Acta Cryst. C52*, 1027–1028.
Harms, K. & Wocadlo, S. (1995). *XCAD4*. University of Marburg, Germany.
North, A. C. T., Phillips, D. C. & Mathews, F. S. (1968). *Acta Cryst. A24*, 351–359.
Sarangarajan, T. R., Panchanatheswaran, K., Low, J. N. & Glidewell, C. (2005). *Acta Cryst. C61*, o118–o121.
Sheldrick, G. M. (2008). *Acta Cryst. A64*, 112–122.
Sonurlikar, U. A., Shanker, B., Kirke, P. A. & Bhide, M. B. (1977). *Bull. Haffkine*, **5**, 94–96.
Thirumurugan, R., Shanmuga Sundara Raj, S., Shanmugam, G., Fun, H.-K., Marappan, M. & Kandaswamy, M. (1998). *Acta Cryst. C54*, 644–645.
Yogavel, M., Selvanayagam, S., Velmurugan, D., Shanmuga Sundara Raj, S., Fun, H.-K., Marappan, M. & Kandaswamy, M. (2003). *Acta Cryst. E59*, o83–o85.
Zhao, H., He, X., Thurkauf, A., Hoffman, D., Kieltyka, A., Brodbeck, R., Primus, R. & Wasley, J. W. F. (2002). *Biorg. Med. Chem. Lett.* **12**, 3111–3115.
Zheng, P.-W., Wang, W. & Duan, X.-M. (2005). *Acta Cryst. E61*, o2513–o2514.

supplementary materials

Acta Cryst. (2010). E66, o3336 [doi:10.1107/S1600536810049111]

1,4-Dibenzylpiperazine

M. Zhang, Y. Zhou, L. Hu and X. Yang

Comment

The structural study of piperazine derivatives is of interest, because some piperazine-containing derivatives constitute a novel class of mixed D2/D4 receptor antagonists (Zhao *et al.*, 2002), and disubstituted piperazine derivatives are antifilarial, antiamoebic and spermicidal agents (Sonurlikar *et al.*, 1977). In addition, piperazine derivatives are useful precursors of mixed-ligand dithiolenes of interest for non-linear optics (Bigoli *et al.*, 2001). Recently, many piperazine derivatives with various substituents have been synthesized (Zheng *et al.*, 2005; Sarangarajan *et al.*, 2005). Herein, we report the crystal structure of the title compound, (I).

The geometry and labeling scheme of the title compound are depicted in Fig. 1, and the packing structure is given in Fig. 2. The piperazine ring exhibits a chair conformation with the usual bond lengths and angles (Yogavel *et al.*, 2003), comparable with those of related reported structures (Gunasekaran *et al.*, 1996; Thirumurugan *et al.*, 1998).

Experimental

To a solution of anhydrous piperazine (5 mmol, 0.43 g) in CH₂Cl₂ (20 ml) was added 2.2 equivalents of triethylamine (1.5 ml), followed by benzyl bromide (10 mmol, 2.66 g) in CH₂Cl₂ (20 ml). After the mixture had been stirred for 10 min., the solvent was removed using a rotary evaporator. The solid residue was washed with water and recrystallized from ethanol-cyclohexane to give a colourless solid (76% yield). Crystals of (I) suitable for X-ray diffraction were obtained by slow evaporation of an ethanol solution.

Refinement

All H atoms were placed in geometrically idealized positions and constrained to ride on their parent atoms, with C—H distances in the range 0.93–0.97 Å, and $U_{\text{iso}}(\text{H}) = 1.2 U_{\text{eq}}$ of the carrier atom.

Figures



Fig. 1. A view of the molecular structure of the title compound, showing displacement ellipsoids at the 30% probability level. Hydrogen atoms are drawn as spheres of arbitrary radius.

supplementary materials




Fig. 2. The packing of the title compound, viewed along the a axis.

1,4-Dibenzylpiperazine

Crystal data

$C_{18}H_{22}N_2$	$D_x = 1.153 \text{ Mg m}^{-3}$
$M_r = 266.38$	Melting point: 372 K
Orthorhombic, $Pbca$	Mo $K\alpha$ radiation, $\lambda = 0.71073 \text{ \AA}$
Hall symbol: -P 2ac 2ab	Cell parameters from 25 reflections
$a = 7.5130 (15) \text{ \AA}$	$\theta = 9-13^\circ$
$b = 19.127 (4) \text{ \AA}$	$\mu = 0.07 \text{ mm}^{-1}$
$c = 21.366 (4) \text{ \AA}$	$T = 293 \text{ K}$
$V = 3070.3 (11) \text{ \AA}^3$	Strip, colorless
$Z = 8$	$0.30 \times 0.20 \times 0.10 \text{ mm}$
$F(000) = 1152$	

Data collection

Enraf–Nonius CAD-4 diffractometer	1650 reflections with $I > 2\sigma(I)$
Radiation source: fine-focus sealed tube graphite	$R_{\text{int}} = 0.045$
$\omega/2\theta$ scans	$\theta_{\max} = 25.3^\circ, \theta_{\min} = 1.9^\circ$
Absorption correction: ψ scan (North <i>et al.</i> , 1968)	$h = 0 \rightarrow 9$
$T_{\min} = 0.980, T_{\max} = 0.993$	$k = 0 \rightarrow 22$
5468 measured reflections	$l = -25 \rightarrow 25$
2781 independent reflections	3 standard reflections every 200 reflections intensity decay: 1%

Refinement

Refinement on F^2	Secondary atom site location: difference Fourier map
Least-squares matrix: full	Hydrogen site location: inferred from neighbouring sites
$R[F^2 > 2\sigma(F^2)] = 0.052$	H-atom parameters constrained
$wR(F^2) = 0.141$	$w = 1/[\sigma^2(F_o^2) + (0.065P)^2]$

$S = 1.00$ where $P = (F_o^2 + 2F_c^2)/3$
 2781 reflections $(\Delta/\sigma)_{\max} < 0.001$
 182 parameters $\Delta\rho_{\max} = 0.16 \text{ e } \text{\AA}^{-3}$
 $\Delta\rho_{\min} = -0.13 \text{ e } \text{\AA}^{-3}$
 0 restraints Extinction correction: *SHELXL97* (Sheldrick, 2008),
 $F_c^* = kFc[1 + 0.001xFc^2\lambda^3/\sin(2\theta)]^{-1/4}$
 0 constraints Extinction coefficient: 0.0097 (11)
 Primary atom site location: structure-invariant direct
 methods

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (\AA^2)

	<i>x</i>	<i>y</i>	<i>z</i>	$U_{\text{iso}}^*/U_{\text{eq}}$
N1	0.1744 (2)	0.09152 (8)	0.62991 (8)	0.0427 (5)
C1	0.6103 (3)	0.28645 (12)	0.57631 (12)	0.0604 (7)
H1A	0.6752	0.3217	0.5569	0.073*
N2	-0.1771 (2)	0.03706 (8)	0.61285 (8)	0.0432 (5)
C2	0.5050 (3)	0.24229 (12)	0.54133 (10)	0.0533 (6)
H2A	0.4989	0.2477	0.4981	0.064*
C3	0.4087 (3)	0.19002 (11)	0.57034 (10)	0.0449 (6)
H3A	0.3382	0.1604	0.5463	0.054*
C4	0.4150 (3)	0.18090 (10)	0.63433 (10)	0.0413 (5)
C5	0.5204 (3)	0.22622 (11)	0.66895 (11)	0.0542 (6)
H5A	0.5252	0.2218	0.7123	0.065*
C6	0.6185 (3)	0.27799 (12)	0.63953 (13)	0.0634 (7)
H6A	0.6908	0.3073	0.6632	0.076*
C7	0.3183 (3)	0.12199 (11)	0.66679 (10)	0.0504 (6)
H7A	0.4034	0.0855	0.6769	0.061*
H7B	0.2696	0.1393	0.7059	0.061*
C8	0.0184 (3)	0.13693 (10)	0.62714 (10)	0.0474 (6)
H8A	0.0520	0.1819	0.6098	0.057*
H8B	-0.0268	0.1446	0.6691	0.057*
C9	-0.1255 (3)	0.10464 (10)	0.58725 (10)	0.0480 (6)
H9A	-0.2282	0.1354	0.5860	0.058*
H9B	-0.0824	0.0987	0.5448	0.058*
C10	-0.0218 (3)	-0.00859 (10)	0.61446 (10)	0.0483 (6)
H10A	0.0230	-0.0154	0.5723	0.058*
H10B	-0.0551	-0.0539	0.6313	0.058*
C11	0.1210 (3)	0.02366 (10)	0.65474 (10)	0.0478 (6)
H11A	0.0768	0.0293	0.6971	0.057*
H11B	0.2235	-0.0072	0.6562	0.057*
C12	-0.3259 (3)	0.00581 (11)	0.57907 (10)	0.0497 (6)
H12A	-0.2825	-0.0132	0.5399	0.060*
H12B	-0.4116	0.0421	0.5692	0.060*
C13	-0.4189 (3)	-0.05156 (10)	0.61511 (9)	0.0396 (5)
C14	-0.5306 (3)	-0.09799 (11)	0.58415 (10)	0.0494 (6)
H14A	-0.5413	-0.0954	0.5408	0.059*
C15	-0.6260 (3)	-0.14791 (12)	0.61664 (12)	0.0569 (6)

supplementary materials

H15A	-0.7012	-0.1782	0.5952	0.068*
C16	-0.6102 (3)	-0.15299 (11)	0.68032 (12)	0.0555 (6)
H16A	-0.6745	-0.1866	0.7022	0.067*
C17	-0.4989 (3)	-0.10828 (11)	0.71169 (10)	0.0505 (6)
H17A	-0.4870	-0.1119	0.7549	0.061*
C18	-0.4041 (3)	-0.05765 (11)	0.67923 (10)	0.0440 (6)
H18A	-0.3295	-0.0274	0.7010	0.053*

Atomic displacement parameters (\AA^2)

	U^{11}	U^{22}	U^{33}	U^{12}	U^{13}	U^{23}
N1	0.0338 (10)	0.0385 (10)	0.0558 (11)	-0.0001 (8)	-0.0015 (8)	0.0068 (8)
C1	0.0516 (15)	0.0518 (15)	0.0779 (18)	-0.0086 (12)	0.0179 (14)	-0.0005 (14)
N2	0.0328 (9)	0.0414 (10)	0.0553 (11)	0.0003 (9)	-0.0012 (9)	0.0073 (9)
C2	0.0508 (14)	0.0543 (14)	0.0547 (14)	-0.0011 (13)	0.0103 (12)	0.0035 (11)
C3	0.0366 (12)	0.0465 (13)	0.0516 (14)	-0.0022 (10)	-0.0013 (10)	-0.0029 (11)
C4	0.0316 (11)	0.0426 (12)	0.0496 (13)	0.0028 (10)	-0.0007 (10)	-0.0019 (10)
C5	0.0519 (15)	0.0588 (15)	0.0520 (14)	-0.0035 (13)	-0.0045 (12)	-0.0074 (11)
C6	0.0495 (15)	0.0580 (16)	0.0826 (19)	-0.0142 (13)	-0.0008 (13)	-0.0161 (14)
C7	0.0457 (13)	0.0516 (13)	0.0541 (13)	-0.0050 (11)	-0.0061 (12)	0.0070 (11)
C8	0.0430 (13)	0.0377 (11)	0.0614 (14)	0.0003 (11)	0.0046 (11)	0.0055 (11)
C9	0.0363 (13)	0.0431 (13)	0.0645 (14)	0.0021 (10)	-0.0021 (11)	0.0114 (11)
C10	0.0402 (13)	0.0389 (12)	0.0658 (15)	0.0006 (10)	0.0027 (11)	0.0046 (11)
C11	0.0382 (12)	0.0422 (13)	0.0629 (14)	0.0030 (10)	-0.0014 (11)	0.0118 (11)
C12	0.0423 (13)	0.0561 (14)	0.0508 (13)	-0.0035 (11)	-0.0036 (11)	0.0066 (11)
C13	0.0308 (11)	0.0449 (12)	0.0432 (12)	0.0024 (10)	-0.0008 (10)	-0.0015 (10)
C14	0.0494 (14)	0.0521 (14)	0.0468 (12)	-0.0022 (12)	-0.0062 (11)	-0.0047 (11)
C15	0.0457 (14)	0.0484 (14)	0.0768 (17)	-0.0097 (12)	-0.0082 (12)	-0.0078 (13)
C16	0.0444 (14)	0.0506 (14)	0.0715 (17)	-0.0052 (12)	0.0107 (12)	0.0068 (12)
C17	0.0425 (14)	0.0584 (14)	0.0505 (13)	0.0004 (12)	0.0039 (11)	0.0055 (11)
C18	0.0345 (12)	0.0480 (13)	0.0495 (13)	-0.0042 (10)	0.0009 (10)	-0.0045 (10)

Geometric parameters (\AA , $^\circ$)


N1—C11	1.458 (2)	C8—H8B	0.9700
N1—C7	1.459 (3)	C9—H9A	0.9700
N1—C8	1.460 (3)	C9—H9B	0.9700
C1—C6	1.362 (3)	C10—C11	1.508 (3)
C1—C2	1.377 (3)	C10—H10A	0.9700
C1—H1A	0.9300	C10—H10B	0.9700
N2—C9	1.456 (2)	C11—H11A	0.9700
N2—C10	1.457 (3)	C11—H11B	0.9700
N2—C12	1.459 (3)	C12—C13	1.512 (3)
C2—C3	1.381 (3)	C12—H12A	0.9700
C2—H2A	0.9300	C12—H12B	0.9700
C3—C4	1.379 (3)	C13—C18	1.379 (3)
C3—H3A	0.9300	C13—C14	1.389 (3)
C4—C5	1.388 (3)	C14—C15	1.381 (3)
C4—C7	1.509 (3)	C14—H14A	0.9300

C5—C6	1.385 (3)	C15—C16	1.369 (3)
C5—H5A	0.9300	C15—H15A	0.9300
C6—H6A	0.9300	C16—C17	1.371 (3)
C7—H7A	0.9700	C16—H16A	0.9300
C7—H7B	0.9700	C17—C18	1.388 (3)
C8—C9	1.509 (3)	C17—H17A	0.9300
C8—H8A	0.9700	C18—H18A	0.9300
C11—N1—C7	111.26 (16)	N2—C9—H9B	109.7
C11—N1—C8	108.88 (16)	C8—C9—H9B	109.7
C7—N1—C8	112.29 (16)	H9A—C9—H9B	108.2
C6—C1—C2	119.4 (2)	N2—C10—C11	109.76 (17)
C6—C1—H1A	120.3	N2—C10—H10A	109.7
C2—C1—H1A	120.3	C11—C10—H10A	109.7
C9—N2—C10	109.14 (16)	N2—C10—H10B	109.7
C9—N2—C12	112.43 (16)	C11—C10—H10B	109.7
C10—N2—C12	112.31 (16)	H10A—C10—H10B	108.2
C1—C2—C3	120.1 (2)	N1—C11—C10	110.62 (17)
C1—C2—H2A	120.0	N1—C11—H11A	109.5
C3—C2—H2A	120.0	C10—C11—H11A	109.5
C4—C3—C2	121.2 (2)	N1—C11—H11B	109.5
C4—C3—H3A	119.4	C10—C11—H11B	109.5
C2—C3—H3A	119.4	H11A—C11—H11B	108.1
C3—C4—C5	118.0 (2)	N2—C12—C13	113.54 (17)
C3—C4—C7	122.24 (19)	N2—C12—H12A	108.9
C5—C4—C7	119.7 (2)	C13—C12—H12A	108.9
C6—C5—C4	120.6 (2)	N2—C12—H12B	108.9
C6—C5—H5A	119.7	C13—C12—H12B	108.9
C4—C5—H5A	119.7	H12A—C12—H12B	107.7
C1—C6—C5	120.7 (2)	C18—C13—C14	117.88 (19)
C1—C6—H6A	119.6	C18—C13—C12	122.00 (18)
C5—C6—H6A	119.6	C14—C13—C12	120.04 (19)
N1—C7—C4	113.99 (17)	C15—C14—C13	121.1 (2)
N1—C7—H7A	108.8	C15—C14—H14A	119.5
C4—C7—H7A	108.8	C13—C14—H14A	119.5
N1—C7—H7B	108.8	C16—C15—C14	120.2 (2)
C4—C7—H7B	108.8	C16—C15—H15A	119.9
H7A—C7—H7B	107.6	C14—C15—H15A	119.9
N1—C8—C9	110.77 (16)	C15—C16—C17	119.6 (2)
N1—C8—H8A	109.5	C15—C16—H16A	120.2
C9—C8—H8A	109.5	C17—C16—H16A	120.2
N1—C8—H8B	109.5	C16—C17—C18	120.3 (2)
C9—C8—H8B	109.5	C16—C17—H17A	119.9
H8A—C8—H8B	108.1	C18—C17—H17A	119.9
N2—C9—C8	109.98 (17)	C13—C18—C17	120.9 (2)
N2—C9—H9A	109.7	C13—C18—H18A	119.5
C8—C9—H9A	109.7	C17—C18—H18A	119.5
C6—C1—C2—C3	0.1 (3)	C9—N2—C10—C11	59.7 (2)
C1—C2—C3—C4	0.1 (3)	C12—N2—C10—C11	-174.95 (16)

supplementary materials

C2—C3—C4—C5	0.4 (3)	C7—N1—C11—C10	-177.43 (17)
C2—C3—C4—C7	-176.9 (2)	C8—N1—C11—C10	58.3 (2)
C3—C4—C5—C6	-1.2 (3)	N2—C10—C11—N1	-60.0 (2)
C7—C4—C5—C6	176.1 (2)	C9—N2—C12—C13	-162.08 (17)
C2—C1—C6—C5	-0.9 (4)	C10—N2—C12—C13	74.4 (2)
C4—C5—C6—C1	1.5 (4)	N2—C12—C13—C18	20.0 (3)
C11—N1—C7—C4	163.51 (17)	N2—C12—C13—C14	-163.34 (18)
C8—N1—C7—C4	-74.2 (2)	C18—C13—C14—C15	1.0 (3)
C3—C4—C7—N1	-19.8 (3)	C12—C13—C14—C15	-175.73 (19)
C5—C4—C7—N1	162.96 (18)	C13—C14—C15—C16	-0.8 (3)
C11—N1—C8—C9	-57.9 (2)	C14—C15—C16—C17	-0.1 (3)
C7—N1—C8—C9	178.46 (16)	C15—C16—C17—C18	0.6 (3)
C10—N2—C9—C8	-59.3 (2)	C14—C13—C18—C17	-0.5 (3)
C12—N2—C9—C8	175.41 (17)	C12—C13—C18—C17	176.23 (19)
N1—C8—C9—N2	59.1 (2)	C16—C17—C18—C13	-0.4 (3)

Fig. 1



supplementary materials

Fig. 2

