

N'-(3,5-Dibromo-2-hydroxybenzylidene)-2-methylbenzohydrazide

Chun-Bao Tang

Department of Chemistry, Jiaying University, Meizhou 514015, People's Republic of China

Correspondence e-mail: tangchunbao@yahoo.com.cn


Received 22 November 2010; accepted 23 November 2010

Key indicators: single-crystal X-ray study; $T = 298\text{ K}$; mean $\sigma(\text{C}-\text{C}) = 0.014\text{ \AA}$; R factor = 0.053; wR factor = 0.159; data-to-parameter ratio = 17.9.

The asymmetric unit of the title compound, $C_{15}H_{12}Br_2N_2O_2$, contains two independent molecules in which the dihedral angles between the benzene rings are 49.5 (7) and 66.4 (7) $^\circ$. Intramolecular O—H···N hydrogen bonds generate S(6) ring motifs in each molecule. In the crystal, molecules are linked through intermolecular N—H···O hydrogen bonds, forming chains along the b axis.

Related literature

For general background to hydrazones, see: Rasras *et al.* (2010); Pyta *et al.* (2010); Angelusiu *et al.* (2010). For related structures, see: Fun *et al.* (2008); Singh & Singh (2010); Ahmad *et al.* (2010); Tang (2010). For reference bond-length data, see: Allen *et al.* (1987) and for hydrogen-bond motifs, see: Bernstein *et al.* (1995).



Experimental

Crystal data

$C_{15}H_{12}Br_2N_2O_2$

$M_r = 412.09$

Monoclinic, $P2_1/c$

$a = 18.636 (3)\text{ \AA}$

$b = 9.606 (2)\text{ \AA}$

$c = 19.943 (3)\text{ \AA}$

$\beta = 113.726 (2)^\circ$

$V = 3268.4 (10)\text{ \AA}^3$

$Z = 8$

Mo $K\alpha$ radiation

$\mu = 4.97\text{ mm}^{-1}$

$T = 298\text{ K}$

$0.17 \times 0.13 \times 0.12\text{ mm}$

Data collection

Bruker SMART CCD area-detector diffractometer

Absorption correction: multi-scan (*SADABS*; Sheldrick, 1996)

$T_{\min} = 0.486$, $T_{\max} = 0.587$

17117 measured reflections

6973 independent reflections

2208 reflections with $I > 2\sigma(I)$

$R_{\text{int}} = 0.116$

Refinement

$R[F^2 > 2\sigma(F^2)] = 0.053$

$wR(F^2) = 0.159$

$S = 0.89$

6973 reflections

389 parameters

2 restraints

H atoms treated by a mixture of independent and constrained refinement

$\Delta\rho_{\max} = 0.32\text{ e \AA}^{-3}$

$\Delta\rho_{\min} = -0.49\text{ e \AA}^{-3}$

Table 1
Hydrogen-bond geometry (\AA , $^\circ$).

$D-\text{H}\cdots A$	$D-\text{H}$	$\text{H}\cdots A$	$D\cdots A$	$D-\text{H}\cdots A$
N2—H2···O4 ⁱ	0.90 (1)	1.85 (2)	2.743 (8)	172 (7)
N4—H4···O2	0.90 (1)	1.92 (2)	2.815 (8)	174 (7)
O3—H3···N3	0.82	1.90	2.624 (8)	146
O1—H1···N1	0.82	1.87	2.584 (8)	145

Symmetry code: (i) $x, y - 1, z$.

Data collection: *SMART* (Bruker, 2002); cell refinement: *SAINT* (Bruker, 2002); data reduction: *SAINT*; program(s) used to solve structure: *SHELXS97* (Sheldrick, 2008); program(s) used to refine structure: *SHELXL97* (Sheldrick, 2008); molecular graphics: *SHELXTL* (Sheldrick, 2008); software used to prepare material for publication: *SHELXL97*.

Financial support from the Jiaying University research fund is gratefully acknowledged.

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: SJ5065).

References

- Ahmad, T., Zia-ur-Rehman, M., Siddiqui, H. L., Mahmud, S. & Parvez, M. (2010). *Acta Cryst. E66*, o976.
- Allen, F. H., Kennard, O., Watson, D. G., Brammer, L., Orpen, A. G. & Taylor, R. (1987). *J. Chem. Soc. Perkin Trans. 2*, pp. S1–19.
- Angelusiu, M. V., Barbuceanu, S. F., Draghici, C. & Almajan, G. L. (2010). *Eur. J. Med. Chem. 45*, 2055–2062.
- Bernstein, J., Davis, R. E., Shimoni, L. & Chang, N.-L. (1995). *Angew. Chem. Int. Ed. Engl. 34*, 1555–1573.
- Bruker (2002). *SMART* and *SAINT*. Bruker AXS Inc., Madison, Wisconsin, USA.
- Fun, H.-K., Sujith, K. V., Patil, P. S., Kalluraya, B. & Chantrapromma, S. (2008). *Acta Cryst. E64*, o1961–o1962.
- Pyta, K., Przybylski, P., Huczynski, A., Hosser, A., Wozniak, K., Schilf, W., Kamienski, B., Grech, E. & Brzezinski, B. (2010). *J. Mol. Struct. 970*, 147–154.
- Rasras, A. J. M., Al-Tel, T. H., Al-Aboudi, A. F. & Al-Qawasmeh, R. A. (2010). *Eur. J. Med. Chem. 45*, 2307–2313.
- Sheldrick, G. M. (1996). *SADABS*. University of Göttingen, Germany.
- Sheldrick, G. M. (2008). *Acta Cryst. A64*, 112–122.
- Singh, V. P. & Singh, S. (2010). *Acta Cryst. E66*, o1172.
- Tang, C.-B. (2010). *Acta Cryst. E66*, o2482.

supplementary materials

Acta Cryst. (2010). E66, o3361 [doi:10.1107/S1600536810048889]

N'-(3,5-Dibromo-2-hydroxybenzylidene)-2-methylbenzohydrazide

C.-B. Tang

Comment

Hydrazone compounds have received much attention in biological and structural chemistry in the last few years (Rasras *et al.*, 2010; Pyta *et al.*, 2010; Angelusiu *et al.*, 2010; Fun *et al.*, 2008; Singh & Singh, 2010; Ahmad *et al.*, 2010). In the present paper, the author reports the crystal structure of the new title hydrazone compound (Fig. 1).

The asymmetric unit of the title compound contains two independent molecules. The dihedral angles between the two benzene rings in the two molecules are 49.5 (7) and 66.4 (7) $^{\circ}$, respectively. The torsion angles C1—C7—N1—N2, C7—N1—N2—C8, N1—N2—C8—C9, C15—C21—N3—N4, C21—N3—N4—C22, and N3—N4—C22—C23 are 3.9 (6), 4.3 (6), 1.8 (6), 2.4 (6), 7.3 (6), and 2.4 (6) $^{\circ}$, respectively. Bond lengths in the molecules are normal (Allen *et al.*, 1987) and comparable to those in the similar compound the author reported recently (Tang, 2010). Intramolecular O1—H1 \cdots N1 and O3—H3 \cdots N3 hydrogen bonds generate S(6) ring motifs in each molecule (Bernstein *et al.*, 1995). In the crystal structure, molecules are linked through intermolecular N—H \cdots O hydrogen bonds (Table 1), forming chains along the *b* axis (Fig. 2).

Experimental

3,5-Dibromo-2-hydroxybenzaldehyde (0.1 mmol, 28.0 mg) and 2-methylbenzohydrazide (0.1 mmol, 15.0 mg) were dissolved in methanol (20 ml). The mixture was stirred at reflux for 10 min to give a clear colourless solution. Colourless block-shaped crystals of the compound were formed by slow evaporation of the solvent over several days.

Refinement

The amino H atoms were located in a difference Fourier map and refined isotropically, with the N—H distances restrained to 0.90 (1) Å [$U_{\text{iso}}(\text{H}) = 0.08 \text{ \AA}^2$]. Other H atoms were constrained to ideal geometries and refined as riding, with C_{sp}^2 —H = 0.93 Å, C(methyl)—H = 0.96 Å, and O—H = 0.82 Å; $U_{\text{iso}}(\text{H}) = 1.2U_{\text{eq}}(\text{C})$ and $1.5U_{\text{eq}}(\text{O and C}_\text{methyl})$.

Figures




Fig. 1. The molecular structure of the compound, showing the atom-numbering scheme. Displacement ellipsoids are drawn at the 30% probability level. Hydrogen atoms are shown as spheres of arbitrary radius.

supplementary materials




Fig. 2. Molecular packing of the title compound, with hydrogen bonds shown as dashed lines.

***N'*-(3,5-Dibromo-2-hydroxybenzylidene)-2-methylbenzohydrazide**

Crystal data

C ₁₅ H ₁₂ Br ₂ N ₂ O ₂	F(000) = 1616
M _r = 412.09	D _x = 1.675 Mg m ⁻³
Monoclinic, P2 ₁ /c	Mo K α radiation, λ = 0.71073 Å
Hall symbol: -P 2ybc	Cell parameters from 1348 reflections
a = 18.636 (3) Å	θ = 2.5–24.1°
b = 9.606 (2) Å	μ = 4.97 mm ⁻¹
c = 19.943 (3) Å	T = 298 K
β = 113.726 (2)°	Block, colourless
V = 3268.4 (10) Å ³	0.17 × 0.13 × 0.12 mm
Z = 8	

Data collection

Bruker SMART CCD area-detector diffractometer	6973 independent reflections
Radiation source: fine-focus sealed tube graphite	2208 reflections with $I > 2\sigma(I)$
ω scans	$R_{\text{int}} = 0.116$
Absorption correction: multi-scan (SADABS; Sheldrick, 1996)	$\theta_{\text{max}} = 27.0^\circ$, $\theta_{\text{min}} = 2.2^\circ$
$T_{\text{min}} = 0.486$, $T_{\text{max}} = 0.587$	$h = -23 \rightarrow 19$
17117 measured reflections	$k = -12 \rightarrow 11$
	$l = -25 \rightarrow 25$

Refinement

Refinement on F^2	Primary atom site location: structure-invariant direct methods
Least-squares matrix: full	Secondary atom site location: difference Fourier map
$R[F^2 > 2\sigma(F^2)] = 0.053$	Hydrogen site location: inferred from neighbouring sites
$wR(F^2) = 0.159$	H atoms treated by a mixture of independent and constrained refinement
$S = 0.89$	$w = 1/[\sigma^2(F_o^2)]$
6973 reflections	where $P = (F_o^2 + 2F_c^2)/3$
389 parameters	$(\Delta/\sigma)_{\text{max}} < 0.001$
2 restraints	$\Delta\rho_{\text{max}} = 0.32 \text{ e \AA}^{-3}$
	$\Delta\rho_{\text{min}} = -0.49 \text{ e \AA}^{-3}$

Special details

Geometry. All esds (except the esd in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell esds are taken into account individually in the estimation of esds in distances, angles and torsion angles; correlations between esds in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell esds is used for estimating esds involving l.s. planes.

Refinement. Refinement of F^2 against ALL reflections. The weighted R-factor wR and goodness of fit S are based on F^2 , conventional R-factors R are based on F, with F set to zero for negative F^2 . The threshold expression of $F^2 > 2\text{sigma}(F^2)$ is used only for calculating R-factors(gt) etc. and is not relevant to the choice of reflections for refinement. R-factors based on F^2 are statistically about twice as large as those based on F, and R- factors based on ALL data will be even larger.

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (\AA^2)

	x	y	z	$U_{\text{iso}}^*/U_{\text{eq}}$
Br1	0.69344 (6)	0.40876 (10)	0.38431 (6)	0.0971 (4)
Br2	0.72350 (5)	-0.17458 (10)	0.37439 (6)	0.0928 (4)
Br3	-0.06953 (6)	0.85420 (10)	-0.01006 (5)	0.0876 (4)
Br4	-0.12081 (6)	0.29585 (11)	0.04659 (6)	0.0984 (4)
N1	0.4069 (4)	0.1700 (7)	0.3119 (3)	0.0615 (18)
N2	0.3301 (4)	0.1455 (7)	0.3007 (4)	0.0614 (19)
N3	0.1898 (4)	0.6601 (6)	0.2214 (3)	0.0576 (18)
N4	0.2608 (4)	0.6440 (7)	0.2795 (4)	0.0605 (18)
O1	0.5235 (3)	0.3360 (6)	0.3345 (3)	0.0757 (17)
H1	0.4775	0.3154	0.3240	0.114*
O2	0.3146 (3)	0.3703 (6)	0.3204 (3)	0.0781 (18)
O3	0.0844 (3)	0.8048 (6)	0.1160 (3)	0.0726 (16)
H3	0.1266	0.7907	0.1504	0.109*
O4	0.2768 (3)	0.8762 (6)	0.2838 (3)	0.0822 (19)
C1	0.5314 (4)	0.0854 (9)	0.3282 (4)	0.051 (2)
C2	0.5659 (5)	0.2190 (9)	0.3415 (4)	0.057 (2)
C3	0.6469 (5)	0.2318 (8)	0.3630 (4)	0.060 (2)
C4	0.6930 (5)	0.1155 (10)	0.3713 (4)	0.067 (2)
H4A	0.7467	0.1249	0.3851	0.081*
C5	0.6590 (5)	-0.0157 (9)	0.3591 (4)	0.065 (2)
C6	0.5792 (5)	-0.0284 (9)	0.3376 (4)	0.058 (2)
H6	0.5569	-0.1168	0.3292	0.069*
C7	0.4486 (4)	0.0642 (9)	0.3118 (4)	0.057 (2)
H7	0.4265	-0.0242	0.3015	0.069*
C8	0.2854 (5)	0.2545 (10)	0.3053 (4)	0.055 (2)
C9	0.2028 (5)	0.2208 (9)	0.2904 (5)	0.061 (2)
C10	0.1668 (6)	0.2838 (11)	0.3318 (5)	0.088 (3)
C11	0.0869 (7)	0.2519 (13)	0.3119 (7)	0.109 (4)
H11	0.0607	0.2906	0.3384	0.131*
C12	0.0470 (7)	0.1646 (14)	0.2540 (7)	0.111 (4)
H12	-0.0059	0.1473	0.2417	0.133*
C13	0.0832 (6)	0.1029 (9)	0.2144 (6)	0.089 (3)
H13	0.0561	0.0433	0.1757	0.107*

supplementary materials

C14	0.1627 (5)	0.1327 (9)	0.2342 (5)	0.073 (3)
H14	0.1888	0.0911	0.2083	0.087*
C15	0.0682 (4)	0.5631 (8)	0.1399 (4)	0.048 (2)
C16	0.0418 (5)	0.6869 (9)	0.1003 (4)	0.054 (2)
C17	-0.0330 (5)	0.6879 (9)	0.0439 (4)	0.059 (2)
C18	-0.0814 (5)	0.5726 (9)	0.0281 (4)	0.066 (2)
H18	-0.1315	0.5755	-0.0090	0.079*
C19	-0.0538 (4)	0.4536 (8)	0.0683 (4)	0.057 (2)
C20	0.0207 (4)	0.4455 (9)	0.1218 (4)	0.056 (2)
H20	0.0394	0.3616	0.1458	0.067*
C21	0.1458 (4)	0.5550 (8)	0.1993 (4)	0.051 (2)
H21	0.1636	0.4695	0.2217	0.061*
C22	0.3020 (5)	0.7621 (9)	0.3066 (4)	0.051 (2)
C23	0.3807 (5)	0.7368 (8)	0.3682 (4)	0.051 (2)
C24	0.4012 (5)	0.8118 (9)	0.4335 (5)	0.065 (2)
C25	0.4759 (7)	0.7929 (11)	0.4871 (5)	0.103 (3)
H25	0.4908	0.8405	0.5312	0.124*
C26	0.5285 (6)	0.7049 (11)	0.4762 (6)	0.102 (3)
H26	0.5786	0.6952	0.5129	0.122*
C27	0.5087 (5)	0.6310 (10)	0.4121 (6)	0.102 (3)
H27	0.5443	0.5710	0.4052	0.122*
C28	0.4339 (5)	0.6489 (9)	0.3583 (5)	0.075 (3)
H28	0.4193	0.6003	0.3145	0.090*
C29	0.2102 (6)	0.3762 (12)	0.3981 (6)	0.138 (5)
H29A	0.2560	0.3286	0.4313	0.208*
H29B	0.1763	0.3970	0.4226	0.208*
H29C	0.2254	0.4613	0.3822	0.208*
C30	0.3438 (6)	0.9038 (10)	0.4487 (5)	0.111 (4)
H30A	0.3303	0.9819	0.4159	0.167*
H30B	0.3673	0.9365	0.4983	0.167*
H30C	0.2974	0.8516	0.4415	0.167*
H4	0.281 (4)	0.559 (3)	0.295 (4)	0.080*
H2	0.309 (4)	0.060 (3)	0.291 (4)	0.080*

Atomic displacement parameters (\AA^2)

	U^{11}	U^{22}	U^{33}	U^{12}	U^{13}	U^{23}
Br1	0.0765 (7)	0.0809 (7)	0.1295 (9)	-0.0250 (6)	0.0369 (7)	-0.0015 (7)
Br2	0.0685 (6)	0.0922 (8)	0.1017 (8)	0.0176 (6)	0.0175 (6)	-0.0253 (6)
Br3	0.0767 (7)	0.0865 (8)	0.0826 (7)	0.0029 (5)	0.0143 (6)	0.0252 (6)
Br4	0.0707 (7)	0.0878 (8)	0.1113 (8)	-0.0274 (6)	0.0101 (6)	-0.0070 (6)
N1	0.051 (5)	0.047 (4)	0.067 (5)	-0.004 (4)	0.004 (4)	0.006 (4)
N2	0.046 (5)	0.045 (5)	0.083 (5)	-0.004 (4)	0.015 (4)	0.007 (4)
N3	0.051 (4)	0.051 (4)	0.059 (5)	-0.013 (4)	0.009 (4)	-0.015 (4)
N4	0.040 (4)	0.056 (5)	0.065 (5)	0.005 (4)	0.000 (4)	0.001 (4)
O1	0.049 (3)	0.062 (4)	0.104 (5)	-0.003 (3)	0.019 (4)	0.015 (3)
O2	0.060 (4)	0.057 (4)	0.107 (5)	0.006 (3)	0.022 (4)	-0.006 (4)
O3	0.060 (4)	0.066 (4)	0.073 (4)	-0.019 (3)	0.006 (3)	0.006 (3)

O4	0.069 (4)	0.047 (4)	0.095 (5)	-0.004 (3)	-0.004 (4)	0.012 (4)
C1	0.050 (5)	0.054 (6)	0.041 (5)	-0.007 (5)	0.009 (4)	-0.006 (4)
C2	0.053 (6)	0.064 (7)	0.041 (5)	-0.002 (5)	0.005 (4)	0.005 (5)
C3	0.060 (6)	0.061 (6)	0.049 (5)	-0.013 (5)	0.010 (5)	-0.001 (4)
C4	0.053 (5)	0.091 (8)	0.055 (6)	0.006 (6)	0.018 (5)	-0.007 (6)
C5	0.062 (6)	0.073 (7)	0.056 (6)	0.010 (5)	0.019 (5)	-0.021 (5)
C6	0.050 (5)	0.063 (6)	0.050 (5)	-0.003 (5)	0.009 (4)	-0.010 (5)
C7	0.042 (5)	0.057 (6)	0.061 (5)	-0.006 (5)	0.009 (4)	0.001 (5)
C8	0.046 (6)	0.053 (6)	0.047 (5)	-0.010 (5)	0.000 (4)	0.006 (5)
C9	0.048 (6)	0.054 (6)	0.075 (7)	-0.011 (5)	0.019 (5)	0.001 (5)
C10	0.087 (8)	0.112 (9)	0.071 (7)	-0.003 (7)	0.038 (7)	0.017 (6)
C11	0.085 (9)	0.168 (12)	0.096 (9)	0.009 (8)	0.058 (8)	0.026 (9)
C12	0.074 (8)	0.164 (13)	0.101 (10)	-0.006 (8)	0.042 (8)	0.051 (9)
C13	0.073 (8)	0.080 (7)	0.105 (8)	-0.008 (6)	0.026 (7)	0.028 (6)
C14	0.048 (6)	0.066 (7)	0.101 (8)	0.005 (5)	0.027 (6)	0.017 (6)
C15	0.055 (5)	0.040 (5)	0.046 (5)	0.006 (4)	0.019 (4)	-0.001 (4)
C16	0.051 (5)	0.060 (6)	0.054 (6)	-0.010 (5)	0.024 (5)	-0.017 (5)
C17	0.046 (5)	0.074 (6)	0.049 (5)	0.009 (5)	0.011 (5)	0.018 (5)
C18	0.052 (5)	0.082 (7)	0.060 (6)	-0.019 (6)	0.018 (5)	-0.007 (5)
C19	0.050 (5)	0.050 (6)	0.057 (6)	-0.004 (4)	0.007 (5)	-0.002 (5)
C20	0.044 (5)	0.064 (6)	0.046 (5)	0.000 (5)	0.004 (4)	-0.010 (4)
C21	0.046 (5)	0.061 (6)	0.034 (5)	0.007 (4)	0.003 (4)	-0.009 (4)
C22	0.049 (5)	0.056 (6)	0.046 (5)	-0.018 (5)	0.016 (5)	-0.009 (5)
C23	0.052 (5)	0.049 (5)	0.049 (6)	-0.008 (4)	0.017 (5)	-0.005 (4)
C24	0.067 (6)	0.073 (6)	0.049 (6)	-0.002 (5)	0.017 (5)	-0.004 (5)
C25	0.104 (9)	0.116 (9)	0.055 (7)	0.005 (8)	-0.004 (7)	-0.018 (6)
C26	0.074 (8)	0.101 (9)	0.093 (9)	-0.013 (7)	-0.007 (7)	-0.013 (7)
C27	0.058 (7)	0.092 (8)	0.114 (9)	0.006 (6)	-0.008 (6)	-0.028 (7)
C28	0.058 (6)	0.078 (7)	0.070 (6)	-0.004 (5)	0.005 (6)	-0.003 (5)
C29	0.124 (10)	0.209 (14)	0.103 (9)	-0.013 (9)	0.067 (8)	-0.035 (9)
C30	0.121 (8)	0.144 (10)	0.078 (7)	0.008 (8)	0.049 (7)	-0.012 (7)

Geometric parameters (\AA , $^\circ$)

Br1—C3	1.878 (8)	C11—H11	0.9300
Br2—C5	1.890 (8)	C12—C13	1.364 (13)
Br3—C17	1.894 (7)	C12—H12	0.9300
Br4—C19	1.900 (8)	C13—C14	1.402 (11)
N1—C7	1.280 (8)	C13—H13	0.9300
N1—N2	1.377 (8)	C14—H14	0.9300
N2—C8	1.364 (10)	C15—C20	1.391 (9)
N2—H2	0.898 (10)	C15—C16	1.402 (10)
N3—C21	1.263 (8)	C15—C21	1.457 (9)
N3—N4	1.372 (8)	C16—C17	1.396 (10)
N4—C22	1.355 (9)	C17—C18	1.382 (10)
N4—H4	0.898 (10)	C18—C19	1.371 (10)
O1—C2	1.348 (8)	C18—H18	0.9300
O1—H1	0.8200	C19—C20	1.370 (9)
O2—C8	1.221 (8)	C20—H20	0.9300

supplementary materials

O3—C16	1.345 (8)	C21—H21	0.9300
O3—H3	0.8200	C22—C23	1.507 (10)
O4—C22	1.207 (8)	C23—C28	1.375 (10)
C1—C6	1.375 (9)	C23—C24	1.400 (10)
C1—C2	1.412 (10)	C24—C25	1.384 (11)
C1—C7	1.457 (9)	C24—C30	1.508 (11)
C2—C3	1.399 (10)	C25—C26	1.378 (12)
C3—C4	1.377 (10)	C25—H25	0.9300
C4—C5	1.387 (10)	C26—C27	1.376 (12)
C4—H4A	0.9300	C26—H26	0.9300
C5—C6	1.378 (9)	C27—C28	1.386 (11)
C6—H6	0.9300	C27—H27	0.9300
C7—H7	0.9300	C28—H28	0.9300
C8—C9	1.481 (10)	C29—H29A	0.9600
C9—C14	1.363 (10)	C29—H29B	0.9600
C9—C10	1.396 (11)	C29—H29C	0.9600
C10—C11	1.411 (13)	C30—H30A	0.9600
C10—C29	1.526 (12)	C30—H30B	0.9600
C11—C12	1.377 (14)	C30—H30C	0.9600
C7—N1—N2	117.1 (6)	C20—C15—C21	119.2 (7)
C8—N2—N1	118.7 (6)	C16—C15—C21	120.8 (7)
C8—N2—H2	119 (5)	O3—C16—C17	119.1 (8)
N1—N2—H2	122 (5)	O3—C16—C15	122.9 (7)
C21—N3—N4	118.3 (7)	C17—C16—C15	118.0 (8)
C22—N4—N3	116.2 (6)	C18—C17—C16	121.8 (8)
C22—N4—H4	122 (5)	C18—C17—Br3	119.6 (6)
N3—N4—H4	122 (5)	C16—C17—Br3	118.6 (7)
C2—O1—H1	109.5	C19—C18—C17	118.5 (7)
C16—O3—H3	109.5	C19—C18—H18	120.8
C6—C1—C2	118.4 (7)	C17—C18—H18	120.8
C6—C1—C7	119.3 (8)	C20—C19—C18	121.8 (7)
C2—C1—C7	122.1 (8)	C20—C19—Br4	119.9 (7)
O1—C2—C3	118.3 (8)	C18—C19—Br4	118.3 (6)
O1—C2—C1	122.3 (7)	C19—C20—C15	119.7 (7)
C3—C2—C1	119.4 (8)	C19—C20—H20	120.2
C4—C3—C2	120.6 (8)	C15—C20—H20	120.2
C4—C3—Br1	119.9 (7)	N3—C21—C15	122.2 (7)
C2—C3—Br1	119.5 (7)	N3—C21—H21	118.9
C3—C4—C5	119.8 (8)	C15—C21—H21	118.9
C3—C4—H4A	120.1	O4—C22—N4	122.5 (8)
C5—C4—H4A	120.1	O4—C22—C23	123.9 (7)
C6—C5—C4	119.6 (8)	N4—C22—C23	113.6 (8)
C6—C5—Br2	121.0 (7)	C28—C23—C24	120.5 (8)
C4—C5—Br2	119.3 (7)	C28—C23—C22	120.2 (7)
C1—C6—C5	122.1 (8)	C24—C23—C22	119.1 (8)
C1—C6—H6	119.0	C25—C24—C23	117.6 (9)
C5—C6—H6	119.0	C25—C24—C30	119.7 (9)
N1—C7—C1	118.5 (7)	C23—C24—C30	122.6 (8)
N1—C7—H7	120.8	C26—C25—C24	121.2 (9)

C1—C7—H7	120.8	C26—C25—H25	119.4
O2—C8—N2	119.5 (7)	C24—C25—H25	119.4
O2—C8—C9	124.9 (9)	C27—C26—C25	121.4 (10)
N2—C8—C9	115.6 (8)	C27—C26—H26	119.3
C14—C9—C10	121.4 (8)	C25—C26—H26	119.3
C14—C9—C8	118.9 (9)	C26—C27—C28	117.7 (9)
C10—C9—C8	119.7 (9)	C26—C27—H27	121.1
C9—C10—C11	116.3 (10)	C28—C27—H27	121.1
C9—C10—C29	123.5 (10)	C23—C28—C27	121.6 (9)
C11—C10—C29	120.1 (11)	C23—C28—H28	119.2
C12—C11—C10	121.4 (11)	C27—C28—H28	119.2
C12—C11—H11	119.3	C10—C29—H29A	109.5
C10—C11—H11	119.3	C10—C29—H29B	109.5
C13—C12—C11	121.6 (11)	H29A—C29—H29B	109.5
C13—C12—H12	119.2	C10—C29—H29C	109.5
C11—C12—H12	119.2	H29A—C29—H29C	109.5
C12—C13—C14	117.5 (10)	H29B—C29—H29C	109.5
C12—C13—H13	121.2	C24—C30—H30A	109.5
C14—C13—H13	121.2	C24—C30—H30B	109.5
C9—C14—C13	121.7 (9)	H30A—C30—H30B	109.5
C9—C14—H14	119.1	C24—C30—H30C	109.5
C13—C14—H14	119.1	H30A—C30—H30C	109.5
C20—C15—C16	120.0 (7)	H30B—C30—H30C	109.5

Hydrogen-bond geometry (Å, °)

<i>D—H···A</i>	<i>D—H</i>	<i>H···A</i>	<i>D···A</i>	<i>D—H···A</i>
N2—H2···O4 ⁱ	0.90 (1)	1.85 (2)	2.743 (8)	172 (7)
N4—H4···O2	0.90 (1)	1.92 (2)	2.815 (8)	174 (7)
O3—H3···N3	0.82	1.90	2.624 (8)	146
O1—H1···N1	0.82	1.87	2.584 (8)	145

Symmetry codes: (i) $x, y-1, z$.

supplementary materials

Fig. 1




Fig. 2

