

4-[(3-Oxo-1,3-dihydro-2-benzofuran-1-yl)amino]benzoic acid

Chao-Jun Du

School of Biology and Chemical Engineering, Nanyang Institute of Technology, Nanyang 473004, People's Republic of China
Correspondence e-mail: chjdu@yahoo.com.cn


Received 13 November 2010; accepted 22 November 2010

Key indicators: single-crystal X-ray study; $T = 296$ K; mean $\sigma(\text{C}-\text{C}) = 0.002$ Å; R factor = 0.035; wR factor = 0.101; data-to-parameter ratio = 12.2.

In the title compound, $\text{C}_{15}\text{H}_{11}\text{NO}_4$, the dihedral angle formed by the benzene ring and the essentially planar 2-benzofuran ring system is $55.93(3)^\circ$. In the crystal, intermolecular $\text{O}-\text{H}\cdots\text{O}$ hydrogen bonds link pairs of molecules, generating centrosymmetric $R_2^2(8)$ ring motifs. These dimeric units are connected via $\text{N}-\text{H}\cdots\text{O}$ hydrogen bonds, forming $C(6)$ chains along $[100]$.

Related literature

For the structure of 2-(3-oxo-1,3-dihydroisobenzofuran-1-yl-amino)benzoic acid, see: Odabaşoğlu & Büyükgüngör (2008). For the structure of 3-(3-oxo-1,3-dihydroisobenzofuran-1-yl-amino)benzoic acid, see: Li *et al.* (2009). For hydrogen-bond motifs, see: Bernstein *et al.* (1995).



Experimental

Crystal data

$\text{C}_{15}\text{H}_{11}\text{NO}_4$
 $M_r = 269.25$
Triclinic, $P\bar{1}$
 $a = 5.9727(18)$ Å
 $b = 6.987(2)$ Å
 $c = 15.451(5)$ Å
 $\alpha = 78.135(3)^\circ$
 $\beta = 87.217(3)^\circ$
 $\gamma = 77.804(3)^\circ$
 $V = 616.8(3)$ Å³
 $Z = 2$
Mo $K\alpha$ radiation
 $\mu = 0.11$ mm⁻¹
 $T = 296$ K
 $0.21 \times 0.16 \times 0.11$ mm

Data collection

Bruker APEXII CCD diffractometer
Absorption correction: multi-scan (SADABS; Sheldrick, 1996)
 $T_{\min} = 0.98$, $T_{\max} = 0.988$
5940 measured reflections
2213 independent reflections
1971 reflections with $I > 2\sigma(I)$
 $R_{\text{int}} = 0.015$

Refinement

$R[F^2 > 2\sigma(F^2)] = 0.035$
 $wR(F^2) = 0.101$
 $S = 1.05$
2213 reflections
182 parameters
H-atom parameters constrained
 $\Delta\rho_{\max} = 0.21$ e Å⁻³
 $\Delta\rho_{\min} = -0.26$ e Å⁻³

Table 1

Hydrogen-bond geometry (Å, °).

$D-\text{H}\cdots A$	$D-\text{H}$	$\text{H}\cdots A$	$D\cdots A$	$D-\text{H}\cdots A$
$\text{N1}-\text{H1}\cdots\text{O4}^i$	0.88	2.11	2.9802 (17)	168
$\text{O1}-\text{H1A}\cdots\text{O2}^{ii}$	0.86	1.79	2.6438 (16)	175

Symmetry codes: (i) $x - 1, y, z$; (ii) $-x + 1, -y + 3, -z$.

Data collection: APEX2 (Bruker, 2004); cell refinement: SAINT (Bruker, 2004); data reduction: SAINT; program(s) used to solve structure: SHELXS97 (Sheldrick, 2008); program(s) used to refine structure: SHELXL97 (Sheldrick, 2008); molecular graphics: SHELXTL (Sheldrick, 2008); software used to prepare material for publication: SHELXTL.

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: LH5161).

References

- Bernstein, J., Davis, R. E., Shimoni, L. & Chang, N.-L. (1995). *Angew. Chem. Int. Ed. Engl.* **34**, 1555–1573.
Bruker (2004). APEX2 and SAINT. Bruker AXS Inc., Madison, Wisconsin, USA.
Li, W., Yin, H., Wen, L., Li, K. & Fan, W. (2009). *Acta Cryst.* **E65**, o2579.
Odabaşoğlu, M. & Büyükgüngör, O. (2008). *Acta Cryst.* **E64**, o752–o753.
Sheldrick, G. M. (1996). SADABS. University of Göttingen, Germany.
Sheldrick, G. M. (2008). *Acta Cryst.* **A64**, 112–122.

supplementary materials

Acta Cryst. (2010). E66, o3364 [doi:10.1107/S1600536810048695]

4-[(3-Oxo-1,3-dihydro-2-benzofuran-1-yl)amino]benzoic acid

C.-J. Du

Comment

The molecular structure of the title compound is shown in Fig. 1. The dihedral angle formed by the benzene ring and the essentially planar isobenzofuran ring system is $55.93(3)^\circ$. In the crystal, intermolecular O—H \cdots O hydrogen bonds link pairs of molecules, generating centrosymmetric $R_2^2(8)$ ring motifs. These dimeric units are connected via N—H \cdots O hydrogen bonds forming C(6) chains along [100] (see Fig. 2). The crystal structures of 2-(3-oxo-1,3-dihydroisobenzofuran-1-ylamino)benzoic acid (Odabaşođlu & Büyükğüngör, 2008) and 3-(3-oxo-1,3-dihydroisobenzofuran-1-ylamino)benzoic (Li *et al.* (2009) have been published previously.

Experimental

4-aminobenzoic acid (5.00 mmol) and 2-formylbenzoic acid (5.00 mmol) were added to an ethanol (35 ml) and DMF (15 ml) mix. The mixture was stirred at 353 K for 5 h. The resulting clear solution was evaporated under vacuum. The product was crystallized from a solution of DMF/methanol(1:1) yielding the title compound. Anal. yield: *ca* 98.6%. Single crystals suitable for X-ray analysis were obtained within one week by slow evaporation of a DMF/methanol (1:3) solution of the title compound.

Refinement

All H atoms were placed in idealized positions (C—H = 0.93 or 0.98 Å, N—H = 0.88 Å and O—H = 0.86 Å), and constrained to ride on the atom to which they are bonded, and were included in the refinement in the riding-model approximation. $U_{\text{iso}}(\text{H})$ values were set equal to $1.5U_{\text{eq}}(\text{parent atom})$ for carboxyl and the secondary amine H atom and to $1.2U_{\text{eq}}(\text{parent atom})$ for all other H atoms.

Figures




Fig. 1. The molecular structure of the title compound with 50% probability displacement ellipsoids.




Fig. 2. Part of the crystal structure with hydrogen bonds shown as dashed lines.

4-[(3-Oxo-1,3-dihydro-2-benzofuran-1-yl)amino]benzoic acid

Crystal data

$C_{15}H_{11}NO_4$	$Z = 2$
$M_r = 269.25$	$F(000) = 280$
Triclinic, PT	$D_x = 1.450 \text{ Mg m}^{-3}$
Hall symbol: $-P 1$	Mo $K\alpha$ radiation, $\lambda = 0.71073 \text{ \AA}$
$a = 5.9727 (18) \text{ \AA}$	Cell parameters from 3521 reflections
$b = 6.987 (2) \text{ \AA}$	$\theta = 2.7\text{--}27.8^\circ$
$c = 15.451 (5) \text{ \AA}$	$\mu = 0.11 \text{ mm}^{-1}$
$\alpha = 78.135 (3)^\circ$	$T = 296 \text{ K}$
$\beta = 87.217 (3)^\circ$	Block, colorless
$\gamma = 77.804 (3)^\circ$	$0.21 \times 0.16 \times 0.11 \text{ mm}$
$V = 616.8 (3) \text{ \AA}^3$	

Data collection

Bruker APEXII CCD diffractometer	2213 independent reflections
Radiation source: fine-focus sealed tube graphite	1971 reflections with $I > 2\sigma(I)$
φ and ω scans	$R_{\text{int}} = 0.015$
Absorption correction: multi-scan (<i>SADABS</i> ; Sheldrick, 1996)	$\theta_{\text{max}} = 25.2^\circ$, $\theta_{\text{min}} = 2.7^\circ$
$T_{\text{min}} = 0.98$, $T_{\text{max}} = 0.988$	$h = -7 \rightarrow 7$
5940 measured reflections	$k = -8 \rightarrow 8$
	$l = -18 \rightarrow 18$

Refinement

Refinement on F^2	Primary atom site location: structure-invariant direct methods
Least-squares matrix: full	Secondary atom site location: difference Fourier map
$R[F^2 > 2\sigma(F^2)] = 0.035$	Hydrogen site location: inferred from neighbouring sites
$wR(F^2) = 0.101$	H-atom parameters constrained
$S = 1.05$	$w = 1/[\sigma^2(F_o^2) + (0.0516P)^2 + 0.1431P]$
2213 reflections	where $P = (F_o^2 + 2F_c^2)/3$
182 parameters	$(\Delta/\sigma)_{\text{max}} < 0.001$
0 restraints	$\Delta\rho_{\text{max}} = 0.21 \text{ e \AA}^{-3}$
	$\Delta\rho_{\text{min}} = -0.26 \text{ e \AA}^{-3}$

Special details

Geometry. All e.s.d.'s (except the e.s.d. in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell e.s.d.'s are taken into account individually in the estimation of e.s.d.'s in distances, angles and torsion angles; correlations

between e.s.d.'s in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell e.s.d.'s is used for estimating e.s.d.'s involving l.s. planes.

Refinement. Refinement of F^2 against ALL reflections. The weighted R -factor wR and goodness of fit S are based on F^2 , conventional R -factors R are based on F , with F set to zero for negative F^2 . The threshold expression of $F^2 > \sigma(F^2)$ is used only for calculating R -factors(gt) *etc.* and is not relevant to the choice of reflections for refinement. R -factors based on F^2 are statistically about twice as large as those based on F , and R -factors based on ALL data will be even larger.

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (\AA^2)

	x	y	z	$U_{\text{iso}}^*/U_{\text{eq}}$
C1	0.5514 (2)	1.2253 (2)	0.06142 (9)	0.0419 (3)
C2	0.5971 (2)	1.0142 (2)	0.10724 (9)	0.0380 (3)
C3	0.4306 (2)	0.9320 (2)	0.15965 (10)	0.0413 (3)
H3	0.2840	1.0090	0.1624	0.050*
C4	0.4797 (2)	0.7390 (2)	0.20729 (10)	0.0409 (3)
H4	0.3668	0.6872	0.2422	0.049*
C5	0.6981 (2)	0.62056 (19)	0.20341 (9)	0.0363 (3)
C6	0.8628 (2)	0.6997 (2)	0.14808 (9)	0.0440 (4)
H6	1.0072	0.6210	0.1428	0.053*
C7	0.8119 (3)	0.8932 (2)	0.10154 (9)	0.0443 (4)
H7	0.9235	0.9443	0.0655	0.053*
C8	0.9645 (2)	0.3035 (2)	0.25671 (9)	0.0403 (3)
H8	1.0056	0.2839	0.1965	0.048*
C9	0.9832 (2)	0.10315 (19)	0.31764 (9)	0.0389 (3)
C10	0.8538 (3)	-0.0406 (2)	0.32249 (10)	0.0498 (4)
H10	0.7294	-0.0204	0.2854	0.060*
C11	0.9156 (3)	-0.2158 (2)	0.38446 (11)	0.0581 (4)
H11	0.8300	-0.3144	0.3894	0.070*
C12	1.1024 (3)	-0.2478 (2)	0.43950 (11)	0.0573 (4)
H12	1.1423	-0.3685	0.4795	0.069*
C13	1.2293 (3)	-0.1033 (2)	0.43565 (10)	0.0495 (4)
H13	1.3534	-0.1229	0.4729	0.059*
C14	1.1648 (2)	0.0728 (2)	0.37402 (9)	0.0397 (3)
C15	1.2613 (2)	0.2526 (2)	0.35724 (10)	0.0433 (3)
N1	0.7473 (2)	0.43002 (16)	0.25574 (8)	0.0417 (3)
H1	0.6533	0.4023	0.3003	0.063*
H1A	0.3357	1.4504	0.0381	0.063*
O1	0.35440 (19)	1.33180 (15)	0.06907 (8)	0.0583 (3)
O2	0.7119 (2)	1.29313 (16)	0.01765 (8)	0.0654 (4)
O3	1.14390 (17)	0.38706 (15)	0.29107 (7)	0.0478 (3)
O4	1.41808 (19)	0.28973 (18)	0.39302 (8)	0.0595 (3)

Atomic displacement parameters (\AA^2)

	U^{11}	U^{22}	U^{33}	U^{12}	U^{13}	U^{23}
C1	0.0449 (8)	0.0338 (7)	0.0402 (7)	-0.0024 (6)	0.0006 (6)	0.0019 (6)
C2	0.0421 (7)	0.0327 (7)	0.0344 (7)	-0.0034 (6)	-0.0009 (6)	0.0000 (5)

supplementary materials

C3	0.0338 (7)	0.0347 (7)	0.0503 (8)	-0.0015 (6)	-0.0016 (6)	-0.0019 (6)
C4	0.0346 (7)	0.0359 (7)	0.0488 (8)	-0.0088 (6)	0.0014 (6)	0.0006 (6)
C5	0.0402 (7)	0.0292 (7)	0.0368 (7)	-0.0060 (5)	-0.0035 (5)	-0.0006 (5)
C6	0.0398 (8)	0.0375 (8)	0.0440 (8)	0.0031 (6)	0.0058 (6)	0.0033 (6)
C7	0.0433 (8)	0.0399 (8)	0.0399 (7)	-0.0030 (6)	0.0079 (6)	0.0068 (6)
C8	0.0446 (8)	0.0325 (7)	0.0387 (7)	-0.0034 (6)	-0.0005 (6)	-0.0004 (6)
C9	0.0472 (8)	0.0300 (7)	0.0355 (7)	-0.0018 (6)	0.0008 (6)	-0.0040 (5)
C10	0.0656 (10)	0.0345 (7)	0.0491 (8)	-0.0105 (7)	-0.0088 (7)	-0.0058 (6)
C11	0.0833 (12)	0.0326 (8)	0.0593 (10)	-0.0185 (8)	-0.0057 (9)	-0.0037 (7)
C12	0.0818 (12)	0.0324 (8)	0.0495 (9)	-0.0058 (8)	-0.0033 (8)	0.0052 (6)
C13	0.0547 (9)	0.0407 (8)	0.0443 (8)	-0.0007 (7)	-0.0043 (7)	0.0027 (6)
C14	0.0408 (7)	0.0351 (7)	0.0377 (7)	-0.0013 (6)	0.0033 (6)	-0.0025 (6)
C15	0.0363 (7)	0.0413 (8)	0.0451 (8)	-0.0034 (6)	0.0035 (6)	0.0025 (6)
N1	0.0402 (6)	0.0314 (6)	0.0459 (7)	-0.0041 (5)	0.0018 (5)	0.0055 (5)
O1	0.0537 (7)	0.0341 (6)	0.0706 (8)	0.0054 (5)	0.0083 (6)	0.0104 (5)
O2	0.0602 (7)	0.0409 (6)	0.0777 (8)	-0.0019 (5)	0.0204 (6)	0.0142 (6)
O3	0.0423 (6)	0.0378 (5)	0.0561 (6)	-0.0092 (4)	-0.0037 (5)	0.0091 (4)
O4	0.0455 (6)	0.0632 (7)	0.0657 (7)	-0.0194 (5)	-0.0085 (5)	0.0068 (6)

Geometric parameters (Å, °)

C1—O2	1.2640 (18)	C8—H8	0.9800
C1—O1	1.2670 (17)	C9—C14	1.378 (2)
C1—C2	1.4736 (19)	C9—C10	1.379 (2)
C2—C7	1.389 (2)	C10—C11	1.383 (2)
C2—C3	1.395 (2)	C10—H10	0.9300
C3—C4	1.3755 (19)	C11—C12	1.388 (3)
C3—H3	0.9300	C11—H11	0.9300
C4—C5	1.395 (2)	C12—C13	1.375 (2)
C4—H4	0.9300	C12—H12	0.9300
C5—N1	1.3884 (17)	C13—C14	1.387 (2)
C5—C6	1.399 (2)	C13—H13	0.9300
C6—C7	1.3729 (19)	C14—C15	1.462 (2)
C6—H6	0.9300	C15—O4	1.2086 (18)
C7—H7	0.9300	C15—O3	1.3476 (17)
C8—N1	1.4054 (18)	N1—H1	0.8837
C8—O3	1.4892 (18)	O1—H1A	0.8551
C8—C9	1.5037 (19)		
O2—C1—O1	122.93 (13)	C14—C9—C10	120.58 (13)
O2—C1—C2	118.54 (13)	C14—C9—C8	109.02 (12)
O1—C1—C2	118.52 (12)	C10—C9—C8	130.40 (13)
C7—C2—C3	118.18 (12)	C9—C10—C11	117.69 (15)
C7—C2—C1	120.29 (13)	C9—C10—H10	121.2
C3—C2—C1	121.49 (12)	C11—C10—H10	121.2
C4—C3—C2	121.09 (13)	C10—C11—C12	121.48 (15)
C4—C3—H3	119.5	C10—C11—H11	119.3
C2—C3—H3	119.5	C12—C11—H11	119.3
C3—C4—C5	120.32 (12)	C13—C12—C11	120.85 (14)
C3—C4—H4	119.8	C13—C12—H12	119.6

C5—C4—H4	119.8	C11—C12—H12	119.6
N1—C5—C4	119.29 (12)	C12—C13—C14	117.30 (15)
N1—C5—C6	121.97 (12)	C12—C13—H13	121.3
C4—C5—C6	118.73 (12)	C14—C13—H13	121.3
C7—C6—C5	120.26 (13)	C9—C14—C13	122.06 (14)
C7—C6—H6	119.9	C9—C14—C15	108.59 (12)
C5—C6—H6	119.9	C13—C14—C15	129.34 (14)
C6—C7—C2	121.32 (13)	O4—C15—O3	121.22 (13)
C6—C7—H7	119.3	O4—C15—C14	129.98 (13)
C2—C7—H7	119.3	O3—C15—C14	108.80 (12)
N1—C8—O3	111.92 (11)	C5—N1—C8	122.57 (12)
N1—C8—C9	114.49 (12)	C5—N1—H1	117.3
O3—C8—C9	102.61 (11)	C8—N1—H1	117.7
N1—C8—H8	109.2	C1—O1—H1A	114.1
O3—C8—H8	109.2	C15—O3—C8	110.77 (11)
C9—C8—H8	109.2		
O2—C1—C2—C7	-0.4 (2)	C10—C11—C12—C13	1.7 (3)
O1—C1—C2—C7	-179.08 (14)	C11—C12—C13—C14	-0.9 (2)
O2—C1—C2—C3	177.30 (14)	C10—C9—C14—C13	1.9 (2)
O1—C1—C2—C3	-1.3 (2)	C8—C9—C14—C13	-177.47 (13)
C7—C2—C3—C4	2.5 (2)	C10—C9—C14—C15	-176.90 (13)
C1—C2—C3—C4	-175.24 (13)	C8—C9—C14—C15	3.73 (15)
C2—C3—C4—C5	-0.5 (2)	C12—C13—C14—C9	-0.8 (2)
C3—C4—C5—N1	176.53 (13)	C12—C13—C14—C15	177.69 (14)
C3—C4—C5—C6	-2.1 (2)	C9—C14—C15—O4	178.44 (15)
N1—C5—C6—C7	-175.90 (14)	C13—C14—C15—O4	-0.2 (3)
C4—C5—C6—C7	2.7 (2)	C9—C14—C15—O3	-1.20 (16)
C5—C6—C7—C2	-0.7 (2)	C13—C14—C15—O3	-179.88 (14)
C3—C2—C7—C6	-2.0 (2)	C4—C5—N1—C8	-178.74 (12)
C1—C2—C7—C6	175.86 (13)	C6—C5—N1—C8	-0.2 (2)
N1—C8—C9—C14	-126.09 (13)	O3—C8—N1—C5	63.86 (17)
O3—C8—C9—C14	-4.61 (14)	C9—C8—N1—C5	-179.92 (12)
N1—C8—C9—C10	54.6 (2)	O4—C15—O3—C8	178.43 (13)
O3—C8—C9—C10	176.11 (14)	C14—C15—O3—C8	-1.89 (15)
C14—C9—C10—C11	-1.1 (2)	N1—C8—O3—C15	127.16 (12)
C8—C9—C10—C11	178.09 (14)	C9—C8—O3—C15	3.93 (14)
C9—C10—C11—C12	-0.6 (3)		

Hydrogen-bond geometry (Å, °)

<i>D</i> —H \cdots <i>A</i>	<i>D</i> —H	H \cdots <i>A</i>	<i>D</i> \cdots <i>A</i>	<i>D</i> —H \cdots <i>A</i>
N1—H1 \cdots O4 ⁱ	0.88	2.11	2.9802 (17)	168
O1—H1A \cdots O2 ⁱⁱ	0.86	1.79	2.6438 (16)	175

Symmetry codes: (i) $x-1, y, z$; (ii) $-x+1, -y+3, -z$.

Fig. 1




Fig. 2

