

Bis(μ -4-amino-3,5-dimethyl-4H-1,2,4-triazole)bis[diiodidozinc(II)]

Rongxian Zhang,^{a*} Qiuyun Chen,^a Xiaofei Yang^b and Xiangyang Wu^c

^aCollege of Chemistry and Chemical Engineering, Jiangsu University, Zhenjiang 212013, People's Republic of China, ^bCollege of Material Science and Engineering, Jiangsu University, Zhenjiang 212013, People's Republic of China, and ^cSchool of Chemistry and Chemical Engineering, Jiangsu University, Zhenjiang 212013, People's Republic of China

Correspondence e-mail: rong@ujjs.edu.cn


Received 8 October 2010; accepted 21 November 2010

Key indicators: single-crystal X-ray study; $T = 293$ K; mean $\sigma(\text{C}-\text{C}) = 0.010$ Å; R factor = 0.040; wR factor = 0.098; data-to-parameter ratio = 18.9.

In the title compound, $[\text{Zn}_2\text{I}_4(\text{C}_4\text{H}_8\text{N}_4)_2]$, the Zn^{II} atom is coordinated in a distorted tetrahedral geometry by two N atoms from the triazole rings of two 4-amino-3,5-dimethyl-4H-1,2,4-triazole (admt) ligands and two iodide ligands. Doubly bridging admt ligands connect two Zn^{II} atoms, forming a centrosymmetric dimer. Weak $\text{N}-\text{H}\cdots\text{I}$ and $\text{C}-\text{H}\cdots\text{I}$ hydrogen bonds play an important role in the intermolecular packing.

Related literature

For background to transition metal complexes of 1,2,4-triazole derivatives, see: Liu *et al.* (1999, 2003); Zhao *et al.* (2002); Yi *et al.* (2004); Lavrenova *et al.* (1992); Haasnoot (2000); Zhang *et al.* (2007).



Experimental

Crystal data

$[\text{Zn}_2\text{I}_4(\text{C}_4\text{H}_8\text{N}_4)_2]$
 $M_r = 862.63$

Monoclinic, $P2_1/c$
 $a = 7.4674$ (19) Å

$b = 13.442$ (3) Å
 $c = 11.412$ (3) Å
 $\beta = 102.598$ (6)°
 $V = 1117.9$ (5) Å³
 $Z = 2$

Mo $K\alpha$ radiation
 $\mu = 7.68$ mm⁻¹
 $T = 293$ K
 $0.30 \times 0.20 \times 0.20$ mm

Data collection

Rigaku Mercury CCD diffractometer
 Absorption correction: multi-scan (REQAB; Jacobson, 1998)
 $T_{\text{min}} = 0.207$, $T_{\text{max}} = 0.309$

10214 measured reflections
 2038 independent reflections
 1760 reflections with $I > 2\sigma(I)$
 $R_{\text{int}} = 0.030$

Refinement

$R[F^2 > 2\sigma(F^2)] = 0.040$
 $wR(F^2) = 0.098$
 $S = 1.04$
 2038 reflections
 108 parameters
 2 restraints

H atoms treated by a mixture of independent and constrained refinement

$\Delta\rho_{\text{max}} = 1.35$ e Å⁻³
 $\Delta\rho_{\text{min}} = -1.11$ e Å⁻³

Table 1

Hydrogen-bond geometry (Å, °).

| $D-\text{H}\cdots A$ | $D-\text{H}$ | $\text{H}\cdots A$ | $D\cdots A$ | $D-\text{H}\cdots A$ |
|----------------------------------------------------|--------------|--------------------|-------------|----------------------|
| $\text{N4}-\text{H4A}\cdots\text{I2}^{\text{i}}$ | 0.86 (2) | 2.98 (5) | 3.706 (7) | 144 (7) |
| $\text{N4}-\text{H4A}\cdots\text{I1}^{\text{ii}}$ | 0.86 (2) | 3.23 (8) | 3.720 (7) | 119 (7) |
| $\text{N4}-\text{H4B}\cdots\text{I1}^{\text{iii}}$ | 0.86 (2) | 3.27 (4) | 4.090 (7) | 161 (7) |
| $\text{C3}-\text{H3A}\cdots\text{I1}^{\text{iv}}$ | 0.96 | 3.24 | 3.930 (8) | 130 |
| $\text{C3}-\text{H3B}\cdots\text{I1}^{\text{iii}}$ | 0.96 | 3.43 | 3.888 (8) | 112 |

Symmetry codes: (i) $-x + 2, y + \frac{1}{2}, -z + \frac{1}{2}$; (ii) $x + 1, y, z$; (iii) $-x + 1, y + \frac{1}{2}, -z + \frac{1}{2}$; (iv) $-x + 1, -y + 1, -z$.

Data collection: *CrystalClear* (Rigaku, 2000); cell refinement: *CrystalClear*; data reduction: *CrystalClear*; program(s) used to solve structure: *SHELXS97* (Sheldrick, 2008); program(s) used to refine structure: *SHELXL97* (Sheldrick, 2008); molecular graphics: *SHELXTL* (Sheldrick, 2008); software used to prepare material for publication: *SHELXTL*.

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: ZL2313).

References

- Haasnoot, J. G. (2000). *Coord. Chem. Rev.* **200–202**, 131–185.
 Jacobson, R. (1998). *REQAB*. Private communication to the Rigaku corporation, Tokyo, Japan.
 Lavrenova, L. G., Baidina, I. A., Ikorskii, V. N., Sheludjkova, L. A. & Larionov, S. V. (1992). *Zh. Neorg. Khim.* **37**, 630–634.
 Liu, J. C., Guo, G. C., Huang, J. S. & You, X. Z. (2003). *Inorg. Chem.* **42**, 235–243.
 Liu, J. C., Xu, Y., Duan, C. Y., Wang, S. L., Liao, F. L., Zhuang, J. Z. & You, X. Z. (1999). *Inorg. Chim. Acta*, **295**, 229–233.
 Rigaku (2000). *CrystalClear*. Rigaku Corporation, Tokyo, Japan.
 Sheldrick, G. M. (2008). *Acta Cryst.* **A64**, 112–122.
 Yi, L., Ding, B., Zhao, B., Cheng, P., Liao, D. Z., Yan, S. P. & Jiang, Z. H. (2004). *Inorg. Chem.* **43**, 33–43.
 Zhang, Y.-M., Zhang, Y.-P., Li, B.-L. & Zhang, Y. (2007). *Acta Cryst.* **C63**, m120–m122.
 Zhao, Q. H., Li, H. F., Chen, Z. D. & Fang, R. B. (2002). *Inorg. Chim. Acta*, **36**, 142–146.

supplementary materials

Acta Cryst. (2011). E67, m26 [doi:10.1107/S160053681004852X]

Bis(μ -4-amino-3,5-dimethyl-4H-1,2,4-triazole)bis[diiodidozinc(II)]

R. Zhang, Q. Chen, X. Yang and X. Wu

Comment

1,2,4-Triazole and its derivatives are very interesting ligands because they combine the coordination geometry of both pyrazole and imidazole with regard to the arrangement of their three heteroatoms. A large number of mononuclear, oligonuclear and polynuclear transition metal complexes of 1,2,4-triazole derivatives have been synthesized and characterized due to their magnetic properties and novel topologies (Haasnoot, 2000). For 4-amino-3,5-dimethyl-1,2,4-triazole (admt), several Mn^{II} (Liu *et al.*, 1999), Co^{II} , Ni^{II} (Zhao *et al.*, 2002), Cu^{II} (Liu *et al.*, 2003) and Cd^{II} compounds (Yi *et al.*, 2004) were synthesized. However, to best of our knowledge, only one Zn^{II} -admt compound, $[\text{Zn}_2(\text{admt})_2\text{Cl}_4]$, was synthesized (Lavrenova *et al.*, 1992). Here, we report the preparation and crystal structure of a dimeric Zn^{II} complex of $[\text{Zn}(\text{admt})\text{I}_2]_2$.

The structure of the title compound is made up of neutral dimeric metallacycle. The title compound has the same molecular structure as its chloro derivative $[\text{Zn}_2(\text{admt})_2\text{Cl}_4]$, but the two compounds have different packing patterns and are not isostructural (Lavrenova *et al.*, 1992). In each dimeric metallacycle, as shown in Fig. 1, two zinc^{II} centers are connected by two admt ligands, resulting in a discrete $\text{Zn}_2(\text{admt})_2$ 6-membered metallacycle which represents the smallest closed cyclic structure with a 1:1 metal-to-ligand ratio. Two triazole rings are coplanar. Each zinc^{II} center is four-coordinated with two nitrogen donors of two admt ligands (Zn1—N1 2.013 (5) Å; Zn1—N2ⁱ (symmetry code i: $-x+1, -y+1, -z$) 2.046 (5) Å) and two I⁻ anions ligands (Zn1—I1 2.560 (1) Å; Zn1—I2 2.549 (1) Å), forming a distorted tetrahedral geometry. The Zn—N (triazole) bond lengths in the title compound are consistent with values in other Zn-triazole complexes (Zhang *et al.*, 2007; Lavrenova *et al.*, 1992). The N—Zn—N, N—Zn—I and I—Zn—I bond angles in the title compound are in the range of 106.8 (2) to 113.75 (3)°, near to the ideal tetrahedral value of ca. 109.5°.

The ligand admt is a 4-substituted 1,2,4-triazole and exhibits in the title compound the N1,N2-bidentate bridging coordination mode. Two admt ligands bridge two Zn(II) atoms to form a dimer with a Zn...Zn distance of 3.803 (2) Å. For a 4-substituted 1,2,4-triazole, by blocking the N4 donor position through substitution, only the N1 monodentate and N1,N2-bidentate coordination modes are possible. The N1,N2-bidentate coordination mode in the dimer has been observed.

There are weak hydrogen bonding interactions between the hydrogen atom of the amino NH_2 group and the I⁻ anion of adjacent dimers (N4...I2ⁱⁱ = 3.706 (7) Å; N4...I1ⁱⁱⁱ = 3.720 (7) Å; N4...I1^{iv} = 4.090 (7) Å) (symmetry codes: ii = $-x+2, y+1/2, -z+1/2$; iii = $x+1, y, z$; iv = $-x+1, y+1/2, -z+1/2$). There are also weak inter-dimer hydrogen bonding interactions between methyl hydrogen atoms and I⁻ anions (C3...I1ⁱ = 3.930 (8) Å; C3...I1^{iv} = 3.888 (8) Å). These hydrogen bonding interactions do direct the packing of the crystal structure of the title compound (Fig. 2). No obvious π - π stacking interactions between the triazole rings is observed.

Experimental

A 15 ml aqueous solution of 4-amino-3,5-dimethyl-1,2,4-triazole (admt) (1.0 mmol) was added to a 10 ml aqueous solution of $\text{Zn}(\text{NO}_3)_2 \cdot 6\text{H}_2\text{O}$ (1.0 mmol) and KI (2.0 mmol) with stirring. The resultant solution was stored at room temperature and colourless crystals were obtained after about two weeks. Anal. Calcd. for $\text{C}_8\text{H}_{16}\text{I}_4\text{N}_8\text{Zn}_2$: C, 11.14; H, 1.87; N, 12.99%. Found: C, 11.09; H, 1.83; N, 12.93%.

Refinement

The H atoms of the amino group were obtained from difference Fourier maps and were refined with N—H distances of 0.86 Å and $U_{\text{iso}}(\text{H}) = 1.2U_{\text{eq}}(\text{N})$. All other H atoms were placed in idealized positions and refined as riding with C—H distances of 0.96 Å and $U_{\text{iso}}(\text{H}) = 1.5U_{\text{eq}}(\text{C})$.

Figures




Fig. 1. The dimeric structure of the title compound, showing the atom-numbering scheme. Displacement ellipsoids are drawn at the 30% probability level. [Symmetry codes: (i) $-x + 1$, $-y + 1$, $-z$.]




Fig. 2. Cell packing plot of the title compound. The dashed lines represent N—H...I hydrogen bond interactions.

Bis(μ -4-amino-3,5-dimethyl-4H-1,2,4-triazole)bis[diiodidozinc(II)]

Crystal data

$[\text{Zn}_2\text{I}_4(\text{C}_4\text{H}_8\text{N}_4)_2]$

$M_r = 862.63$

Monoclinic, $P2_1/c$

Hall symbol: $-P\ 2ybc$

$a = 7.4674(19)$ Å

$b = 13.442(3)$ Å

$F(000) = 784$

$D_x = 2.563$ Mg m $^{-3}$

Mo $K\alpha$ radiation, $\lambda = 0.71070$ Å

Cell parameters from 3727 reflections

$\theta = 3.0$ – 25.4°

$\mu = 7.68$ mm $^{-1}$

$c = 11.412 (3) \text{ \AA}$
 $\beta = 102.598 (6)^\circ$
 $V = 1117.9 (5) \text{ \AA}^3$
 $Z = 2$

$T = 293 \text{ K}$
 Block, colorless
 $0.30 \times 0.20 \times 0.20 \text{ mm}$

Data collection

Rigaku Mercury CCD diffractometer
 Radiation source: fine-focus sealed tube graphite
 Detector resolution: $7.31 \text{ pixels mm}^{-1}$
 ω scans
 Absorption correction: multi-scan (Jacobson, 1998)
 $T_{\min} = 0.207, T_{\max} = 0.309$
 10214 measured reflections

2038 independent reflections
 1760 reflections with $I > 2\sigma(I)$
 $R_{\text{int}} = 0.030$
 $\theta_{\max} = 25.4^\circ, \theta_{\min} = 3.0^\circ$
 $h = -8 \rightarrow 8$
 $k = -16 \rightarrow 14$
 $l = -13 \rightarrow 13$

Refinement

Refinement on F^2
 Least-squares matrix: full
 $R[F^2 > 2\sigma(F^2)] = 0.040$
 $wR(F^2) = 0.098$
 $S = 1.04$
 2038 reflections
 108 parameters
 2 restraints

Primary atom site location: structure-invariant direct methods
 Secondary atom site location: difference Fourier map
 Hydrogen site location: inferred from neighbouring sites
 H atoms treated by a mixture of independent and constrained refinement
 $w = 1/[\sigma^2(F_o^2) + (0.0483P)^2 + 3.598P]$
 where $P = (F_o^2 + 2F_c^2)/3$
 $(\Delta/\sigma)_{\max} < 0.001$
 $\Delta\rho_{\max} = 1.35 \text{ e \AA}^{-3}$
 $\Delta\rho_{\min} = -1.11 \text{ e \AA}^{-3}$

Special details

Geometry. All e.s.d.'s (except the e.s.d. in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell e.s.d.'s are taken into account individually in the estimation of e.s.d.'s in distances, angles and torsion angles; correlations between e.s.d.'s in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell e.s.d.'s is used for estimating e.s.d.'s involving l.s. planes.

Refinement. Refinement of F^2 against ALL reflections. The weighted R -factor wR and goodness of fit S are based on F^2 , conventional R -factors R are based on F , with F set to zero for negative F^2 . The threshold expression of $F^2 > \sigma(F^2)$ is used only for calculating R -factors(gt) *etc.* and is not relevant to the choice of reflections for refinement. R -factors based on F^2 are statistically about twice as large as those based on F , and R -factors based on ALL data will be even larger.

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (\AA^2)

| | | | |
|-----|-----|-----|----------------------------------|
| x | y | z | $U_{\text{iso}}^*/U_{\text{eq}}$ |
|-----|-----|-----|----------------------------------|

supplementary materials

| | | | | |
|-----|-------------|-------------|-------------|-------------|
| Zn1 | 0.42494 (9) | 0.40526 (5) | 0.10329 (6) | 0.0386 (2) |
| I1 | 0.16668 (7) | 0.43239 (4) | 0.21514 (5) | 0.0657 (2) |
| I2 | 0.56923 (7) | 0.23271 (4) | 0.13413 (6) | 0.0743 (2) |
| N1 | 0.6190 (7) | 0.5117 (4) | 0.1541 (4) | 0.0405 (12) |
| N2 | 0.6793 (7) | 0.5755 (4) | 0.0763 (5) | 0.0417 (12) |
| N3 | 0.8344 (7) | 0.6026 (4) | 0.2567 (4) | 0.0387 (12) |
| N4 | 0.9608 (9) | 0.6397 (5) | 0.3571 (6) | 0.0565 (15) |
| C1 | 0.8137 (9) | 0.6303 (5) | 0.1401 (6) | 0.0439 (15) |
| C2 | 0.7150 (8) | 0.5286 (5) | 0.2633 (5) | 0.0391 (14) |
| C3 | 0.9265 (11) | 0.7049 (6) | 0.0964 (7) | 0.060 (2) |
| H3A | 0.9910 | 0.6741 | 0.0418 | 0.090* |
| H3B | 1.0133 | 0.7325 | 0.1631 | 0.090* |
| H3C | 0.8492 | 0.7569 | 0.0556 | 0.090* |
| C4 | 0.7035 (11) | 0.4766 (6) | 0.3741 (6) | 0.0603 (19) |
| H4C | 0.5954 | 0.4358 | 0.3600 | 0.090* |
| H4D | 0.6975 | 0.5244 | 0.4356 | 0.090* |
| H4E | 0.8101 | 0.4355 | 0.3993 | 0.090* |
| H4A | 1.069 (5) | 0.632 (6) | 0.345 (7) | 0.072* |
| H4B | 0.963 (12) | 0.7033 (17) | 0.353 (8) | 0.072* |

Atomic displacement parameters (\AA^2)

| | U^{11} | U^{22} | U^{33} | U^{12} | U^{13} | U^{23} |
|-----|------------|------------|------------|-------------|------------|-------------|
| Zn1 | 0.0331 (4) | 0.0396 (4) | 0.0418 (4) | -0.0002 (3) | 0.0052 (3) | 0.0039 (3) |
| I1 | 0.0538 (3) | 0.0680 (4) | 0.0837 (4) | 0.0007 (2) | 0.0337 (3) | -0.0101 (3) |
| I2 | 0.0540 (3) | 0.0465 (3) | 0.1169 (5) | 0.0152 (2) | 0.0064 (3) | 0.0084 (3) |
| N1 | 0.037 (3) | 0.043 (3) | 0.040 (3) | -0.004 (2) | 0.006 (2) | 0.007 (2) |
| N2 | 0.039 (3) | 0.044 (3) | 0.039 (3) | -0.004 (2) | 0.003 (2) | 0.001 (2) |
| N3 | 0.034 (3) | 0.039 (3) | 0.038 (3) | 0.000 (2) | -0.001 (2) | -0.005 (2) |
| N4 | 0.051 (4) | 0.063 (4) | 0.048 (3) | -0.012 (3) | -0.005 (3) | -0.015 (3) |
| C1 | 0.046 (4) | 0.039 (3) | 0.042 (4) | -0.004 (3) | 0.001 (3) | -0.006 (3) |
| C2 | 0.035 (3) | 0.040 (3) | 0.040 (3) | 0.002 (3) | 0.003 (3) | -0.002 (3) |
| C3 | 0.065 (5) | 0.058 (4) | 0.054 (4) | -0.028 (4) | 0.008 (4) | -0.005 (3) |
| C4 | 0.060 (5) | 0.072 (5) | 0.046 (4) | -0.008 (4) | 0.005 (3) | 0.008 (4) |

Geometric parameters (\AA , $^\circ$)

| | | | |
|---------------------|-------------|--------|-----------|
| Zn1—N1 | 2.029 (5) | N4—H4A | 0.86 (2) |
| Zn1—N2 ⁱ | 2.044 (5) | N4—H4B | 0.86 (2) |
| Zn1—I2 | 2.5493 (10) | C1—C3 | 1.465 (9) |
| Zn1—I1 | 2.5603 (10) | C2—C4 | 1.464 (9) |
| N1—C2 | 1.314 (8) | C3—H3A | 0.9600 |
| N1—N2 | 1.378 (7) | C3—H3B | 0.9600 |
| N2—C1 | 1.327 (8) | C3—H3C | 0.9600 |
| N2—Zn1 ⁱ | 2.044 (5) | C4—H4C | 0.9600 |
| N3—C2 | 1.349 (8) | C4—H4D | 0.9600 |
| N3—C1 | 1.358 (8) | C4—H4E | 0.9600 |
| N3—N4 | 1.408 (7) | | |

| | | | |
|----------------------------|-------------|----------------------------|------------|
| N1—Zn1—N2 ⁱ | 106.8 (2) | N2—C1—N3 | 107.2 (6) |
| N1—Zn1—I2 | 110.38 (15) | N2—C1—C3 | 128.0 (6) |
| N2 ⁱ —Zn1—I2 | 108.08 (15) | N3—C1—C3 | 124.8 (6) |
| N1—Zn1—I1 | 109.00 (15) | N1—C2—N3 | 107.7 (5) |
| N2 ⁱ —Zn1—I1 | 108.58 (16) | N1—C2—C4 | 127.9 (6) |
| I2—Zn1—I1 | 113.73 (3) | N3—C2—C4 | 124.4 (6) |
| C2—N1—N2 | 108.4 (5) | C1—C3—H3A | 109.5 |
| C2—N1—Zn1 | 126.9 (4) | C1—C3—H3B | 109.5 |
| N2—N1—Zn1 | 124.6 (4) | H3A—C3—H3B | 109.5 |
| C1—N2—N1 | 107.9 (5) | C1—C3—H3C | 109.5 |
| C1—N2—Zn1 ⁱ | 123.8 (4) | H3A—C3—H3C | 109.5 |
| N1—N2—Zn1 ⁱ | 128.2 (4) | H3B—C3—H3C | 109.5 |
| C2—N3—C1 | 108.8 (5) | C2—C4—H4C | 109.5 |
| C2—N3—N4 | 123.4 (5) | C2—C4—H4D | 109.5 |
| C1—N3—N4 | 127.8 (6) | H4C—C4—H4D | 109.5 |
| N3—N4—H4A | 108 (6) | C2—C4—H4E | 109.5 |
| N3—N4—H4B | 109 (6) | H4C—C4—H4E | 109.5 |
| H4A—N4—H4B | 94 (8) | H4D—C4—H4E | 109.5 |
| N2 ⁱ —Zn1—N1—C2 | 177.2 (5) | Zn1 ⁱ —N2—C1—C3 | -7.6 (10) |
| I2—Zn1—N1—C2 | -65.6 (5) | C2—N3—C1—N2 | 1.6 (7) |
| I1—Zn1—N1—C2 | 60.0 (5) | N4—N3—C1—N2 | 179.7 (6) |
| N2 ⁱ —Zn1—N1—N2 | -6.8 (6) | C2—N3—C1—C3 | -176.4 (7) |
| I2—Zn1—N1—N2 | 110.5 (4) | N4—N3—C1—C3 | 1.7 (11) |
| I1—Zn1—N1—N2 | -124.0 (4) | N2—N1—C2—N3 | 0.5 (7) |
| C2—N1—N2—C1 | 0.5 (7) | Zn1—N1—C2—N3 | 177.1 (4) |
| Zn1—N1—N2—C1 | -176.2 (4) | N2—N1—C2—C4 | -177.8 (7) |
| C2—N1—N2—Zn1 ⁱ | -175.0 (4) | Zn1—N1—C2—C4 | -1.2 (10) |
| Zn1—N1—N2—Zn1 ⁱ | 8.3 (7) | C1—N3—C2—N1 | -1.3 (7) |
| N1—N2—C1—N3 | -1.3 (7) | N4—N3—C2—N1 | -179.5 (6) |
| Zn1 ⁱ —N2—C1—N3 | 174.5 (4) | C1—N3—C2—C4 | 177.1 (6) |
| N1—N2—C1—C3 | 176.7 (7) | N4—N3—C2—C4 | -1.1 (10) |

Symmetry codes: (i) $-x+1, -y+1, -z$.

Hydrogen-bond geometry (Å, °)

| <i>D</i> —H... <i>A</i> | <i>D</i> —H | H... <i>A</i> | <i>D</i> ... <i>A</i> | <i>D</i> —H... <i>A</i> |
|----------------------------|-------------|---------------|-----------------------|-------------------------|
| N4—H4A...I2 ⁱⁱ | 0.86 (2) | 2.98 (5) | 3.706 (7) | 144 (7) |
| N4—H4A...I1 ⁱⁱⁱ | 0.86 (2) | 3.23 (8) | 3.720 (7) | 119 (7) |
| N4—H4B...I1 ^{iv} | 0.86 (2) | 3.27 (4) | 4.090 (7) | 161 (7) |
| C3—H3A...I1 ⁱ | 0.96 | 3.24 | 3.930 (8) | 130 |
| C3—H3B...I1 ^{iv} | 0.96 | 3.43 | 3.888 (8) | 112 |

Symmetry codes: (ii) $-x+2, y+1/2, -z+1/2$; (iii) $x+1, y, z$; (iv) $-x+1, y+1/2, -z+1/2$; (i) $-x+1, -y+1, -z$.

Fig. 1




Fig. 2

