

## Dibromido(2,2'-dimethyl-4,4'-bi-1,3-thiazole- $\kappa^2N,N'$ )mercury(II)

**Anita Abedi**

Department of Chemistry, Islamic Azad University, North Tehran Branch, Tehran, Iran

Correspondence e-mail: anita\_abedi@yahoo.com


Received 3 December 2010; accepted 8 December 2010

 Key indicators: single-crystal X-ray study;  $T = 100$  K; mean  $\sigma(\text{C}-\text{C}) = 0.007$  Å;  $R$  factor = 0.026;  $wR$  factor = 0.062; data-to-parameter ratio = 25.1.

The asymmetric unit of the title compound,  $[\text{HgBr}_2(\text{C}_8\text{H}_8\text{N}_2\text{S}_2)]$ , contains two crystallographically independent molecules. The  $\text{Hg}^{\text{II}}$  atoms are four-coordinated in a distorted tetrahedral geometry by two N atoms from a 2,2'-dimethyl-4,4'-bi-1,3-thiazole ligand and two Br atoms. In the crystal structure, intermolecular  $\text{C}-\text{H} \cdots \text{Br}$  hydrogen bonds and  $\pi-\pi$  contacts between the thiazole rings [centroid-centroid distances = 3.670 (3) and 3.614 (2) Å] stabilize the structure.

### Related literature

For metal complexes with 2,2'-dimethyl-4,4'-bithiazole ligands, see: Abedi & Yahyazade Bali (2010); Al-Hashemi *et al.* (2009, 2010); Khavasi *et al.* (2008); Notash *et al.* (2008, 2009). For related structures, see: Kalateh *et al.* (2008); Safari *et al.* (2009).



### Experimental

#### Crystal data

$[\text{HgBr}_2(\text{C}_8\text{H}_8\text{N}_2\text{S}_2)]$   
 $M_r = 556.69$   
 Triclinic,  $P\bar{1}$

$a = 10.2799$  (6) Å  
 $b = 11.1595$  (7) Å  
 $c = 11.6821$  (7) Å

$\alpha = 88.4456$  (11)°  
 $\beta = 85.3290$  (11)°  
 $\gamma = 77.1162$  (11)°  
 $V = 1302.02$  (14) Å<sup>3</sup>  
 $Z = 4$

Mo  $K\alpha$  radiation  
 $\mu = 18.25$  mm<sup>-1</sup>  
 $T = 100$  K  
 $0.14 \times 0.12 \times 0.08$  mm

#### Data collection

Bruker APEXII CCD diffractometer  
 Absorption correction: multi-scan (SADABS; Bruker, 2001)  
 $T_{\min} = 0.070$ ,  $T_{\max} = 0.240$

20881 measured reflections  
 6912 independent reflections  
 5865 reflections with  $I > 2\sigma(I)$   
 $R_{\text{int}} = 0.041$

#### Refinement

$R[F^2 > 2\sigma(F^2)] = 0.026$   
 $wR(F^2) = 0.062$   
 $S = 0.94$   
 6912 reflections

275 parameters  
 H-atom parameters constrained  
 $\Delta\rho_{\max} = 1.28$  e Å<sup>-3</sup>  
 $\Delta\rho_{\min} = -1.68$  e Å<sup>-3</sup>

**Table 1**

Selected bond lengths (Å).

Hg1—N1	2.379 (4)	Hg2—N3	2.357 (4)
Hg1—N2	2.383 (4)	Hg2—N4	2.410 (4)
Hg1—Br1	2.4970 (5)	Hg2—Br3	2.4999 (5)
Hg1—Br2	2.5206 (5)	Hg2—Br4	2.4957 (5)

**Table 2**

Hydrogen-bond geometry (Å, °).

$D-H \cdots A$	$D-H$	$H \cdots A$	$D \cdots A$	$D-H \cdots A$
C8—H8B $\cdots$ Br4 <sup>i</sup>	0.98	2.92	3.826 (5)	155
C10—H10A $\cdots$ Br4 <sup>i</sup>	0.95	2.92	3.760 (5)	148
C16—H16B $\cdots$ Br3 <sup>ii</sup>	0.98	2.87	3.772 (5)	154
C16—H16C $\cdots$ Br2 <sup>iii</sup>	0.98	2.88	3.837 (5)	165

Symmetry codes: (i)  $-x + 1, -y + 2, -z + 1$ ; (ii)  $-x + 1, -y + 1, -z + 1$ ; (iii)  $x - 1, y, z$ .

Data collection: APEX2 (Bruker, 2007); cell refinement: SAINT (Bruker, 2007); data reduction: SAINT; program(s) used to solve structure: SHELXS97 (Sheldrick, 2008); program(s) used to refine structure: SHELXL97 (Sheldrick, 2008); molecular graphics: SHELXTL (Sheldrick, 2008) and Mercury (Macrae *et al.*, 2006); software used to prepare material for publication: SHELXTL.

The author is grateful to the Islamic Azad University, North Tehran Branch, for financial support.

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: HY2386).

### References

- Abedi, A. & Yahyazade Bali, E. (2010). *Acta Cryst.* **E66**, m1023.  
 Al-Hashemi, R., Safari, N., Abedi, A., Notash, B., Amani, V. & Khavasi, H. R. (2009). *J. Coord. Chem.* **62**, 2909–2918.  
 Al-Hashemi, R., Safari, N., Amani, S., Amani, V., Abedi, A., Khavasi, H. R. & Ng, S. W. (2010). *J. Coord. Chem.* **63**, 3207–3217.  
 Bruker (2001). SADABS. Bruker AXS Inc., Madison, Wisconsin, USA.  
 Bruker (2007). APEX2 and SAINT. Bruker AXS Inc., Madison, Wisconsin, USA.  
 Kalateh, K., Norouzi, A., Ebadi, A., Ahmadi, R. & Amani, V. (2008). *Acta Cryst.* **E64**, m1583–m1584.

- Khavasi, H. R., Abedi, A., Amani, V., Notash, B. & Safari, N. (2008). *Polyhedron*, **27**, 1848–1854.
- Macrae, C. F., Edgington, P. R., McCabe, P., Pidcock, E., Shields, G. P., Taylor, R., Towler, M. & van de Streek, J. (2006). *J. Appl. Cryst.* **39**, 453–457.
- Notash, B., Safari, N., Abedi, A., Amani, V. & Khavasi, H. R. (2009). *J. Coord. Chem.* **62**, 1638–1649.
- Notash, B., Safari, N., Khavasi, H. R., Amani, V. & Abedi, A. (2008). *J. Organomet. Chem.* **693**, 3553–3557.
- Safari, N., Amani, V., Abedi, A., Notash, B. & Ng, S. W. (2009). *Acta Cryst. E* **65**, m372.
- Sheldrick, G. M. (2008). *Acta Cryst. A* **64**, 112–122.

**supplementary materials**

*Acta Cryst.* (2011). E67, m76-m77 [ doi:10.1107/S1600536810051494 ]

## Dibromido(2,2'-dimethyl-4,4'-bi-1,3-thiazole- $\kappa^2$ N,N')mercury(II)

A. Abedi

### Comment

Recently, we reported the synthesis and crystal structure of [HgI<sub>2</sub>(dm4bt)] (dm4bt is 2,2'-dimethyl-4,4'-bithiazole) (Abedi & Yahyazade Bali, 2010). Dm4bt is a good bidentate ligand, and numerous complexes with dm4bt have been prepared, such as that of zinc (Khavasi *et al.*, 2008), thallium (Notash *et al.*, 2008), cadmium (Notash *et al.*, 2009) and copper (Al-Hashemi *et al.*, 2009, 2010). For further investigation of dm4bt, we synthesis the title complex and report herein its crystal structure.

The asymmetric unit of the title compound (Fig. 1) contains two crystallographically independent molecules. The Hg<sup>II</sup> atom is four-coordinated in a distorted tetrahedral configuration by two N atoms from a dm4bt ligand and two Br atoms. The Hg—N and Hg—Br bond lengths (Table 1) and angles are within normal range found in [Hg(SCN)<sub>2</sub>(dm4bt)] (Safari *et al.*, 2009) and {HgBr<sub>2</sub>[NH(py)<sub>2</sub>]} [NH(py)<sub>2</sub> is di-2-pyridylamine] (Kalatch *et al.*, 2008).

In the crystal structure, intermolecular C—H $\cdots$ Br hydrogen bonds (Table 2) and  $\pi$ – $\pi$  contacts (Fig. 2) between the thiazole rings, Cg1 $\cdots$ Cg2<sup>i</sup> and Cg3 $\cdots$ Cg4<sup>ii</sup> [symmetry codes: (i) 2-x, 1-y, -z; (ii) 1-x, 2-y, 1-z. Cg1, Cg2, Cg3 and Cg4 are the centroids of the rings S1/C1/N1/C3/C2, S2/C4/N2/C6/C5, S3/C9/N3/C11/C10 and S4/C12/N4/C14/C13, respectively] stabilize the structure, with centroid–centroid distances of 3.670 (3) and 3.614 (2) Å.

### Experimental

For the preparation of the title compound, a solution of dm4bt (0.26 g, 1.3 mmol) in methanol (15 ml) was added to a solution of HgBr<sub>2</sub> (0.47 g, 1.3 mmol) in methanol (15 ml) at room temperature. Crystals suitable for X-ray diffraction experiment were obtained by methanol diffusion into a colorless solution in DMSO after one week (yield: 0.52 g, 71.8%).

### Refinement

H atoms were positioned geometrically and refined as riding atoms, with C—H = 0.95 (CH) and 0.98 (CH<sub>3</sub>) Å and with  $U_{\text{iso}}(\text{H}) = 1.2(1.5 \text{ for methyl})U_{\text{eq}}(\text{C})$ . The highest residual electron density was found at 0.85 Å from Hg2 atom and the deepest hole at 0.70 Å from Hg1 atom.

### Figures




Fig. 1. The molecular structure of the title compound. Displacement ellipsoids are drawn at the 50% probability level.




Fig. 2. Crystal packing diagram for the title compound. Dashed lines denote hydrogen bonds.

## Dibromido(2,2'-dimethyl-4,4'-bi-1,3-thiazole- $\kappa^2N,N'$ )mercury(II)

### Crystal data

[HgBr<sub>2</sub>(C<sub>8</sub>H<sub>8</sub>N<sub>2</sub>S<sub>2</sub>)]

$M_r = 556.69$

Triclinic, *P* $\bar{1}$

Hall symbol: -P 1

$a = 10.2799$  (6) Å

$b = 11.1595$  (7) Å

$c = 11.6821$  (7) Å

$\alpha = 88.4456$  (11)°

$\beta = 85.3290$  (11)°

$\gamma = 77.1162$  (11)°

$V = 1302.02$  (14) Å<sup>3</sup>

$Z = 4$

$F(000) = 1008$

$D_x = 2.840$  Mg m<sup>-3</sup>

Mo  $K\alpha$  radiation,  $\lambda = 0.71073$  Å

Cell parameters from 2931 reflections

$\theta = 3.0$ – $29.0$ °

$\mu = 18.25$  mm<sup>-1</sup>

$T = 100$  K

Prism, colorless

$0.14 \times 0.12 \times 0.08$  mm

### Data collection

Bruker APEXII CCD  
diffractometer

Radiation source: fine-focus sealed tube  
graphite

$\varphi$  and  $\omega$  scans

Absorption correction: multi-scan  
(*SADABS*; Bruker, 2001)

$T_{\min} = 0.070$ ,  $T_{\max} = 0.240$

20881 measured reflections

6912 independent reflections

5865 reflections with  $I > 2\sigma(I)$

$R_{\text{int}} = 0.041$

$\theta_{\max} = 29.0$ °,  $\theta_{\min} = 1.8$ °

$h = -14 \rightarrow 14$

$k = -15 \rightarrow 15$

$l = -15 \rightarrow 15$

### Refinement

Refinement on  $F^2$

Least-squares matrix: full

$R[F^2 > 2\sigma(F^2)] = 0.026$

$wR(F^2) = 0.062$

$S = 0.94$

6912 reflections

275 parameters

Primary atom site location: structure-invariant direct methods

Secondary atom site location: difference Fourier map

Hydrogen site location: inferred from neighbouring sites

H-atom parameters constrained

$w = 1/[\sigma^2(F_o^2) + (0.0299P)^2 + 2.0838P]$

where  $P = (F_o^2 + 2F_c^2)/3$

$(\Delta/\sigma)_{\max} = 0.002$

$\Delta\rho_{\max} = 1.28$  e Å<sup>-3</sup>

0 restraints

$$\Delta\rho_{\min} = -1.68 \text{ e } \text{\AA}^{-3}$$

*Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters ( $\text{\AA}^2$ )*

	<i>x</i>	<i>y</i>	<i>z</i>	$U_{\text{iso}}^*/U_{\text{eq}}$
Hg1	1.080846 (18)	0.703396 (17)	0.207265 (14)	0.01438 (5)
Br1	1.27509 (5)	0.73077 (5)	0.07429 (4)	0.01974 (10)
Br2	1.01869 (5)	0.79183 (4)	0.40562 (4)	0.01472 (9)
S1	0.96732 (13)	0.30682 (11)	0.30191 (10)	0.0162 (2)
S2	0.74266 (12)	0.71656 (11)	-0.06846 (10)	0.0167 (2)
N1	1.0123 (4)	0.5145 (4)	0.2400 (3)	0.0126 (8)
N2	0.9123 (4)	0.6940 (3)	0.0815 (3)	0.0118 (7)
C1	1.0489 (4)	0.4251 (4)	0.3135 (4)	0.0117 (8)
C2	0.8814 (5)	0.3826 (4)	0.1925 (4)	0.0160 (9)
H2A	0.8189	0.3523	0.1522	0.019*
C3	0.9157 (5)	0.4932 (4)	0.1714 (4)	0.0129 (9)
C4	0.8605 (5)	0.7689 (4)	0.0020 (4)	0.0148 (9)
C5	0.7668 (5)	0.5861 (4)	0.0144 (4)	0.0150 (9)
H5A	0.7206	0.5217	0.0097	0.018*
C6	0.8618 (5)	0.5883 (4)	0.0881 (4)	0.0131 (9)
C7	1.1512 (5)	0.4226 (5)	0.3961 (4)	0.0186 (10)
H7A	1.1390	0.5043	0.4296	0.028*
H7B	1.2407	0.3995	0.3561	0.028*
H7C	1.1418	0.3625	0.4571	0.028*
C8	0.8973 (5)	0.8888 (4)	-0.0288 (4)	0.0189 (10)
H8A	0.9945	0.8786	-0.0294	0.028*
H8B	0.8531	0.9507	0.0280	0.028*
H8C	0.8684	0.9155	-0.1050	0.028*
Hg2	0.411665 (18)	0.785100 (16)	0.717555 (15)	0.01467 (5)
Br3	0.50713 (5)	0.58257 (4)	0.80847 (4)	0.01660 (10)
Br4	0.19137 (5)	0.93379 (4)	0.74712 (4)	0.01734 (10)
S3	0.72536 (12)	1.05628 (11)	0.68030 (10)	0.0150 (2)
S4	0.59092 (12)	0.68616 (11)	0.31954 (10)	0.0161 (2)
N3	0.5664 (4)	0.9109 (4)	0.6782 (3)	0.0142 (8)
N4	0.5044 (4)	0.7580 (3)	0.5213 (3)	0.0110 (7)
C9	0.6015 (5)	0.9912 (4)	0.7442 (4)	0.0136 (9)
C10	0.7285 (5)	0.9747 (4)	0.5581 (4)	0.0154 (9)
H10A	0.7849	0.9794	0.4904	0.018*
C11	0.6378 (4)	0.9014 (4)	0.5721 (4)	0.0114 (8)
C12	0.4840 (4)	0.6864 (4)	0.4407 (4)	0.0138 (9)
C13	0.6630 (5)	0.7909 (4)	0.3806 (4)	0.0137 (9)
H13A	0.7328	0.8249	0.3443	0.016*
C14	0.6059 (4)	0.8182 (4)	0.4879 (4)	0.0100 (8)
C15	0.5358 (5)	1.0239 (5)	0.8604 (4)	0.0176 (10)
H15A	0.4717	0.9721	0.8807	0.026*
H15B	0.4885	1.1105	0.8608	0.026*
H15C	0.6036	1.0107	0.9164	0.026*
C16	0.3766 (5)	0.6164 (5)	0.4523 (4)	0.0181 (10)

## supplementary materials

---

H16A	0.3632	0.5912	0.5326	0.027*
H16B	0.4023	0.5434	0.4038	0.027*
H16C	0.2932	0.6685	0.4281	0.027*

### Atomic displacement parameters ( $\text{\AA}^2$ )

	$U^{11}$	$U^{22}$	$U^{33}$	$U^{12}$	$U^{13}$	$U^{23}$
Hg1	0.01399 (9)	0.01808 (10)	0.01340 (8)	-0.00844 (7)	-0.00148 (6)	0.00061 (6)
Br1	0.0193 (2)	0.0236 (3)	0.0190 (2)	-0.0124 (2)	0.00513 (19)	-0.00411 (18)
Br2	0.0117 (2)	0.0184 (2)	0.0138 (2)	-0.00327 (17)	0.00036 (16)	-0.00020 (17)
S1	0.0190 (6)	0.0165 (6)	0.0161 (5)	-0.0099 (5)	-0.0037 (4)	0.0045 (4)
S2	0.0160 (6)	0.0177 (6)	0.0179 (5)	-0.0051 (5)	-0.0079 (5)	0.0019 (4)
N1	0.0086 (18)	0.015 (2)	0.0150 (18)	-0.0052 (15)	-0.0010 (15)	0.0000 (15)
N2	0.0097 (18)	0.0125 (19)	0.0134 (18)	-0.0036 (15)	0.0000 (14)	0.0017 (14)
C1	0.007 (2)	0.012 (2)	0.015 (2)	-0.0021 (16)	0.0031 (16)	-0.0011 (16)
C2	0.017 (2)	0.020 (2)	0.014 (2)	-0.011 (2)	-0.0026 (18)	0.0032 (18)
C3	0.013 (2)	0.015 (2)	0.011 (2)	-0.0035 (18)	-0.0011 (17)	0.0001 (16)
C4	0.017 (2)	0.018 (2)	0.011 (2)	-0.0052 (19)	-0.0008 (17)	-0.0031 (17)
C5	0.014 (2)	0.016 (2)	0.015 (2)	-0.0045 (18)	-0.0055 (18)	0.0006 (17)
C6	0.009 (2)	0.016 (2)	0.015 (2)	-0.0045 (18)	0.0010 (17)	-0.0052 (17)
C7	0.018 (2)	0.022 (3)	0.019 (2)	-0.008 (2)	-0.0075 (19)	0.0053 (19)
C8	0.020 (3)	0.015 (2)	0.023 (2)	-0.005 (2)	-0.004 (2)	-0.0003 (19)
Hg2	0.01185 (9)	0.01587 (9)	0.01664 (9)	-0.00476 (7)	0.00171 (7)	-0.00051 (6)
Br3	0.0161 (2)	0.0170 (2)	0.0169 (2)	-0.00399 (18)	-0.00171 (17)	0.00080 (17)
Br4	0.0137 (2)	0.0168 (2)	0.0201 (2)	-0.00250 (18)	0.00373 (18)	0.00131 (17)
S3	0.0160 (6)	0.0169 (6)	0.0139 (5)	-0.0084 (5)	0.0024 (4)	-0.0025 (4)
S4	0.0136 (5)	0.0197 (6)	0.0151 (5)	-0.0042 (5)	0.0012 (4)	-0.0057 (4)
N3	0.0134 (19)	0.015 (2)	0.0149 (18)	-0.0048 (16)	-0.0012 (15)	-0.0004 (15)
N4	0.0065 (17)	0.0114 (18)	0.0140 (17)	0.0002 (14)	0.0005 (14)	-0.0018 (14)
C9	0.012 (2)	0.015 (2)	0.013 (2)	-0.0031 (17)	0.0027 (17)	0.0006 (17)
C10	0.015 (2)	0.017 (2)	0.014 (2)	-0.0043 (18)	0.0002 (18)	-0.0004 (17)
C11	0.007 (2)	0.015 (2)	0.012 (2)	-0.0022 (17)	-0.0010 (16)	0.0004 (16)
C12	0.0029 (19)	0.020 (2)	0.017 (2)	0.0006 (17)	0.0003 (16)	-0.0006 (18)
C13	0.011 (2)	0.013 (2)	0.017 (2)	-0.0019 (17)	-0.0004 (17)	-0.0004 (17)
C14	0.0051 (19)	0.009 (2)	0.015 (2)	-0.0003 (16)	-0.0022 (16)	0.0033 (16)
C15	0.022 (3)	0.023 (3)	0.009 (2)	-0.009 (2)	0.0028 (18)	-0.0022 (18)
C16	0.012 (2)	0.023 (3)	0.021 (2)	-0.0072 (19)	-0.0017 (19)	-0.006 (2)

### Geometric parameters ( $\text{\AA}$ , $^\circ$ )

Hg1—N1	2.379 (4)	Hg2—N3	2.357 (4)
Hg1—N2	2.383 (4)	Hg2—N4	2.410 (4)
Hg1—Br1	2.4970 (5)	Hg2—Br3	2.4999 (5)
Hg1—Br2	2.5206 (5)	Hg2—Br4	2.4957 (5)
S1—C2	1.709 (5)	S3—C10	1.708 (5)
S1—C1	1.727 (4)	S3—C9	1.713 (5)
S2—C5	1.708 (5)	S4—C13	1.715 (5)
S2—C4	1.729 (5)	S4—C12	1.720 (5)
N1—C1	1.309 (6)	N3—C9	1.324 (6)

N1—C3	1.389 (6)	N3—C11	1.383 (6)
N2—C4	1.298 (6)	N4—C12	1.310 (6)
N2—C6	1.388 (6)	N4—C14	1.387 (5)
C1—C7	1.480 (6)	C9—C15	1.483 (6)
C2—C3	1.368 (6)	C10—C11	1.368 (6)
C2—H2A	0.9500	C10—H10A	0.9500
C3—C6	1.467 (7)	C11—C14	1.475 (6)
C4—C8	1.496 (7)	C12—C16	1.483 (6)
C5—C6	1.358 (6)	C13—C14	1.353 (6)
C5—H5A	0.9500	C13—H13A	0.9500
C7—H7A	0.9800	C15—H15A	0.9800
C7—H7B	0.9800	C15—H15B	0.9800
C7—H7C	0.9800	C15—H15C	0.9800
C8—H8A	0.9800	C16—H16A	0.9800
C8—H8B	0.9800	C16—H16B	0.9800
C8—H8C	0.9800	C16—H16C	0.9800
N1—Hg1—N2	71.23 (13)	N3—Hg2—N4	70.25 (13)
N1—Hg1—Br1	124.79 (9)	N3—Hg2—Br4	104.07 (10)
N2—Hg1—Br1	103.63 (9)	N4—Hg2—Br4	115.64 (9)
N1—Hg1—Br2	98.36 (9)	N3—Hg2—Br3	114.30 (10)
N2—Hg1—Br2	120.67 (9)	N4—Hg2—Br3	102.77 (9)
Br1—Hg1—Br2	126.492 (16)	Br4—Hg2—Br3	132.848 (17)
C2—S1—C1	90.8 (2)	C10—S3—C9	90.6 (2)
C5—S2—C4	90.4 (2)	C13—S4—C12	90.4 (2)
C1—N1—C3	112.7 (4)	C9—N3—C11	111.5 (4)
C1—N1—Hg1	131.2 (3)	C9—N3—Hg2	130.2 (3)
C3—N1—Hg1	116.2 (3)	C11—N3—Hg2	118.3 (3)
C4—N2—C6	111.8 (4)	C12—N4—C14	112.3 (4)
C4—N2—Hg1	131.6 (3)	C12—N4—Hg2	130.9 (3)
C6—N2—Hg1	116.5 (3)	C14—N4—Hg2	116.6 (3)
N1—C1—C7	124.3 (4)	N3—C9—C15	122.7 (4)
N1—C1—S1	112.7 (3)	N3—C9—S3	113.5 (3)
C7—C1—S1	123.0 (4)	C15—C9—S3	123.8 (4)
C3—C2—S1	110.1 (3)	C11—C10—S3	110.1 (3)
C3—C2—H2A	125.0	C11—C10—H10A	125.0
S1—C2—H2A	125.0	S3—C10—H10A	125.0
C2—C3—N1	113.8 (4)	C10—C11—N3	114.4 (4)
C2—C3—C6	127.8 (4)	C10—C11—C14	128.1 (4)
N1—C3—C6	118.4 (4)	N3—C11—C14	117.5 (4)
N2—C4—C8	124.7 (4)	N4—C12—C16	123.1 (4)
N2—C4—S2	113.3 (3)	N4—C12—S4	112.9 (3)
C8—C4—S2	122.0 (4)	C16—C12—S4	124.0 (4)
C6—C5—S2	109.8 (4)	C14—C13—S4	110.1 (3)
C6—C5—H5A	125.1	C14—C13—H13A	124.9
S2—C5—H5A	125.1	S4—C13—H13A	124.9
C5—C6—N2	114.7 (4)	C13—C14—N4	114.2 (4)
C5—C6—C3	127.7 (4)	C13—C14—C11	128.8 (4)
N2—C6—C3	117.7 (4)	N4—C14—C11	117.0 (4)
C1—C7—H7A	109.5	C9—C15—H15A	109.5



## supplementary materials

---

C1—C7—H7B	109.5	C9—C15—H15B	109.5
H7A—C7—H7B	109.5	H15A—C15—H15B	109.5
C1—C7—H7C	109.5	C9—C15—H15C	109.5
H7A—C7—H7C	109.5	H15A—C15—H15C	109.5
H7B—C7—H7C	109.5	H15B—C15—H15C	109.5
C4—C8—H8A	109.5	C12—C16—H16A	109.5
C4—C8—H8B	109.5	C12—C16—H16B	109.5
H8A—C8—H8B	109.5	H16A—C16—H16B	109.5
C4—C8—H8C	109.5	C12—C16—H16C	109.5
H8A—C8—H8C	109.5	H16A—C16—H16C	109.5
H8B—C8—H8C	109.5	H16B—C16—H16C	109.5
N2—Hg1—N1—C1	-178.0 (4)	N4—Hg2—N3—C9	177.3 (5)
Br1—Hg1—N1—C1	88.2 (4)	Br4—Hg2—N3—C9	64.7 (4)
Br2—Hg1—N1—C1	-58.3 (4)	Br3—Hg2—N3—C9	-87.2 (4)
N2—Hg1—N1—C3	1.3 (3)	N4—Hg2—N3—C11	-4.7 (3)
Br1—Hg1—N1—C3	-92.5 (3)	Br4—Hg2—N3—C11	-117.3 (3)
Br2—Hg1—N1—C3	121.0 (3)	Br3—Hg2—N3—C11	90.8 (3)
N1—Hg1—N2—C4	-177.5 (4)	N3—Hg2—N4—C12	178.4 (4)
Br1—Hg1—N2—C4	-55.0 (4)	Br4—Hg2—N4—C12	-85.0 (4)
Br2—Hg1—N2—C4	93.8 (4)	Br3—Hg2—N4—C12	66.9 (4)
N1—Hg1—N2—C6	-0.9 (3)	N3—Hg2—N4—C14	5.4 (3)
Br1—Hg1—N2—C6	121.7 (3)	Br4—Hg2—N4—C14	101.9 (3)
Br2—Hg1—N2—C6	-89.5 (3)	Br3—Hg2—N4—C14	-106.2 (3)
C3—N1—C1—C7	179.2 (4)	C11—N3—C9—C15	176.8 (4)
Hg1—N1—C1—C7	-1.5 (7)	Hg2—N3—C9—C15	-5.0 (7)
C3—N1—C1—S1	0.6 (5)	C11—N3—C9—S3	-1.2 (5)
Hg1—N1—C1—S1	179.9 (2)	Hg2—N3—C9—S3	177.0 (2)
C2—S1—C1—N1	0.3 (4)	C10—S3—C9—N3	1.2 (4)
C2—S1—C1—C7	-178.3 (4)	C10—S3—C9—C15	-176.8 (5)
C1—S1—C2—C3	-1.2 (4)	C9—S3—C10—C11	-0.9 (4)
S1—C2—C3—N1	1.7 (5)	S3—C10—C11—N3	0.4 (5)
S1—C2—C3—C6	-177.4 (4)	S3—C10—C11—C14	178.7 (4)
C1—N1—C3—C2	-1.5 (6)	C9—N3—C11—C10	0.5 (6)
Hg1—N1—C3—C2	179.0 (3)	Hg2—N3—C11—C10	-177.9 (3)
C1—N1—C3—C6	177.7 (4)	C9—N3—C11—C14	-178.0 (4)
Hg1—N1—C3—C6	-1.7 (5)	Hg2—N3—C11—C14	3.6 (5)
C6—N2—C4—C8	-178.1 (4)	C14—N4—C12—C16	-176.9 (4)
Hg1—N2—C4—C8	-1.3 (7)	Hg2—N4—C12—C16	9.8 (7)
C6—N2—C4—S2	1.5 (5)	C14—N4—C12—S4	1.2 (5)
Hg1—N2—C4—S2	178.2 (2)	Hg2—N4—C12—S4	-172.1 (2)
C5—S2—C4—N2	-0.4 (4)	C13—S4—C12—N4	-1.5 (4)
C5—S2—C4—C8	179.2 (4)	C13—S4—C12—C16	176.6 (4)
C4—S2—C5—C6	-0.8 (4)	C12—S4—C13—C14	1.3 (4)
S2—C5—C6—N2	1.9 (5)	S4—C13—C14—N4	-0.9 (5)
S2—C5—C6—C3	-177.9 (4)	S4—C13—C14—C11	178.8 (4)
C4—N2—C6—C5	-2.2 (6)	C12—N4—C14—C13	-0.2 (6)
Hg1—N2—C6—C5	-179.5 (3)	Hg2—N4—C14—C13	174.1 (3)
C4—N2—C6—C3	177.6 (4)	C12—N4—C14—C11	-179.9 (4)
Hg1—N2—C6—C3	0.3 (5)	Hg2—N4—C14—C11	-5.6 (5)

C2—C3—C6—C5	-0.1 (8)	C10—C11—C14—C13	3.5 (8)
N1—C3—C6—C5	-179.3 (4)	N3—C11—C14—C13	-178.2 (5)
C2—C3—C6—N2	-179.9 (4)	C10—C11—C14—N4	-176.8 (5)
N1—C3—C6—N2	1.0 (6)	N3—C11—C14—N4	1.5 (6)

*Hydrogen-bond geometry* ( $\text{\AA}$ ,  $^\circ$ )

$D-H\cdots A$	$D-H$	$H\cdots A$	$D\cdots A$	$D-H\cdots A$
C8—H8B $\cdots$ Br4 <sup>i</sup>	0.98	2.92	3.826 (5)	155
C10—H10A $\cdots$ Br4 <sup>i</sup>	0.95	2.92	3.760 (5)	148
C16—H16B $\cdots$ Br3 <sup>ii</sup>	0.98	2.87	3.772 (5)	154
C16—H16C $\cdots$ Br2 <sup>iii</sup>	0.98	2.88	3.837 (5)	165

Symmetry codes: (i)  $-x+1, -y+2, -z+1$ ; (ii)  $-x+1, -y+1, -z+1$ ; (iii)  $x-1, y, z$ .

Fig. 1




Fig. 2

