

Acta Crystallographica Section E

Structure Reports

Online

ISSN 1600-5368

Bis{tris[2-(2-oxidobenzylideneazaniumyl)-ethyl]amine- $\kappa^3 O, O', O''$ }calcium bis(perchlorate) acetonitrile disolvate

Muhammet Kose* and Vickie McKee

Chemistry Department, Loughborough University, Leicestershire, LE11 3TU, England
Correspondence e-mail: m.kose@lboro.ac.uk


Received 13 December 2010; accepted 23 December 2010

Key indicators: single-crystal X-ray study; $T = 150$ K; mean $\sigma(C-C) = 0.002$ Å;
R factor = 0.039; wR factor = 0.110; data-to-parameter ratio = 17.4.

The title complex, $[Ca(C_{27}H_{30}N_4O_3)_2](ClO_4)_2 \cdot 2CH_3CN$, is composed of centrosymmetric $(CaL_2)^{2+}$ cations [$L = \text{tris}(2\text{-hydroxybenzoylaminoethyl})\text{amine} = H_3\text{saltren}$], uncoordinated perchlorate anions and acetonitrile solvent molecules. The calcium ion is six-coordinated and is bonded to all phenoxy O atoms from both zwitterionic saltren molecules. There are strong intramolecular $N-H \cdots O$ hydrogen bonds. The cations are linked into chains *via* weak intermolecular $C-H \cdots O$ hydrogen bonds and $C-H \cdots \pi$ and $\pi-\pi$ stacking interactions [centroid-centroid distances = 3.306 (3) and 3.415 (3) Å].

Related literature

For crystal structure of the free ligand, see: Gündüz *et al.* (1985). For structures of transition metal complexes of $H_3\text{saltren}$, see: Steinhäuser *et al.* (2004); Elerman *et al.* (1994, 1995).



Experimental

Crystal data

 $[Ca(C_{27}H_{30}N_4O_3)_2](ClO_4)_2 \cdot 2C_2H_3N$ $M_r = 1238.19$
Orthorhombic, $Pbca$ $a = 11.3469$ (7) Å $b = 19.5307$ (12) Å $c = 27.3178$ (16) Å $V = 6054.0$ (6) Å³ $Z = 4$ Mo $K\alpha$ radiation $\mu = 0.27$ mm⁻¹ $T = 150$ K $0.12 \times 0.10 \times 0.09$ mm

Data collection

 Bruker APEXII CCD
diffractometer
Absorption correction: multi-scan
(*SADABS*; Bruker, 2008)
 $T_{\min} = 0.662$, $T_{\max} = 0.746$

 55467 measured reflections
6915 independent reflections
5810 reflections with $I > 2.0\sigma(I)$
 $R_{\text{int}} = 0.030$

Refinement

 $R[F^2 > 2\sigma(F^2)] = 0.039$ $wR(F^2) = 0.110$ $S = 1.05$

6915 reflections

398 parameters

 H atoms treated by a mixture of
independent and constrained
refinement
 $\Delta\rho_{\text{max}} = 0.60$ e Å⁻³ $\Delta\rho_{\text{min}} = -0.39$ e Å⁻³
Table 1

Hydrogen-bond geometry (Å, °).

Cg1 is the centroid of the C1–C6 ring.

$D-H \cdots A$	$D-H$	$H \cdots A$	$D \cdots A$	$D-H \cdots A$
$N1-H1A \cdots O1$	0.85 (2)	1.98 (2)	2.6597 (17)	136 (2)
$N3-H3A \cdots O2$	0.86 (2)	1.91 (2)	2.6256 (18)	140 (2)
$N4-H4A \cdots O3$	0.83 (2)	1.98 (2)	2.6449 (17)	137 (2)
$C3-H3 \cdots O7^i$	0.95	2.41	3.316 (2)	160
$C20-H20A \cdots O5^{ii}$	0.99	2.50	3.380 (2)	148
$C10-H10A \cdots O5^{iii}$	0.99	2.50	3.271 (2)	134
$C21-H21 \cdots O5^{iv}$	0.95	2.58	3.259 (2)	129
$C29-H29B \cdots O6^{iv}$	0.98	2.58	3.555 (3)	173
$C26-H26 \cdots Cg1^v$	0.95	2.68	3.489 (2)	143

 Symmetry codes: (i) $-x + 1, -y + 1, -z + 1$; (ii) $x - 1, y - 1, z$; (iii) $x - \frac{1}{2}, y - 1, -z + \frac{3}{2}$; (iv) $-x + \frac{3}{2}, y - \frac{1}{2}, z$; (v) $-x + 1, -y, -z + 1$.

Data collection: *APEX2* (Bruker, 2008); cell refinement: *SAINT* (Bruker, 2008); data reduction: *SAINT*; program(s) used to solve structure: *SHELXS97* (Sheldrick, 2008); program(s) used to refine structure: *SHELXL97* (Sheldrick, 2008); molecular graphics: *SHELXTL* (Sheldrick, 2008); software used to prepare material for publication: *SHELXTL*.

We are grateful to the Turkish Government for the award of a postgraduate scholarship (to MK).

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: PV2371).

References

- Bruker (2008). *APEX2*, *SAINT* and *SADABS*. Bruker AXS Inc., Madison, Wisconsin, USA.
- Elerman, Y., Kabak, M., Svoboda, I., Fuess, H. & Atakol, O. (1995). *J. Chem. Crystallogr.* **25**, 227–230.
- Elerman, Y., Kabak, M., Svoboda, I. & Geselle, M. (1994). *Acta Cryst.* **C50**, 1694–1696.
- Gündüz, N., Gündüz, T., Hursthouse, M. B., Parkes, H. G., Shaw (née Gözen), L. S., Shaw, R. A. & Tüzün, M. (1985). *J. Chem. Soc. Perkin Trans. 2*, pp. 899–902.
- Sheldrick, G. M. (2008). *Acta Cryst.* **A64**, 112–122.
- Steinhäuser, S., Bachman, F., Hazenkamp, M., Heinz, U., Dannacher, J. & Hegetschweiler, K. (2004). *Z. Kristallogr. New Cryst. Struct.* **219**, 325–326.

supplementary materials

Acta Cryst. (2011). E67, m149 [doi:10.1107/S1600536810053961]

Bis{tris[2-(2-oxidobenzylideneazaniumyl)ethyl]amine- κ^3O,O',O'' }calcium bis(perchlorate) acetonitrile disolvate

M. Kose and V. McKee

Comment

The crystal structures and coordination chemistry of transition metal complexes of H₃saltren, tris(2-hydroxybenzoylaminoethyl)amine (*L*), a tripodal ligand, have been described in detail (Steinhauser *et al.*, 2004; Elerman *et al.*, 1994); Elerman *et al.*, 1995). Here, we report the first example of an earth alkaline metal (Ca⁺²) complex of H₃saltren.

The crystal structure of the title complex contains centrosymmetric (CaL₂)⁺² cations, uncoordinated perchlorate anions and acetonitrile molecules (Fig. 1). Each calcium ion is six-coordinated, bonded to all phenoxy O atoms from two saltren molecules. There are strong intramolecular hydrogen bonds of the type N—H \cdots O between protonated imine and deprotonated phenol-oxygen atoms (Table 1). The cations are linked into chains *via* intermolecular C—H \cdots O and C—H \cdots N type weak hydrogen bonds and C—H \cdots π interactions (Tab. 1 and Fig. 2). There is evidence of π - π stacking in the crystal structure; C16 is stacked with C16* and C15* of an adjacent molecule (* = 2 - x, -y, 1 - z) with separation of 3.306 (3) and 3.415 (3) Å, respectively (Fig. 2). Protonated imine distances (C=N distances) (mean 1.300 (2) Å) are longer than the corresponding C=N distances reported in the neutral ligand (mean 1.267 Å) (Gündüz *et al.*, 1985). The longer C=N distances were also determined by IR spectroscopy; while C=N stretch is observed at 1633 cm⁻¹ in the neutral ligand, by protonation of imine, C=N stretch shifts to 1654 cm⁻¹.

Experimental

H₃Saltren (0.508 g, 1.11 mmol) was dissolved in methanol (50 ml) followed by addition of Ca(ClO₄)₂·3H₂O (0.185 g, 0.63 mmol) in methanol (20 ml). The reaction mixture was stirred at room temperature for two hours. Yellow precipitate was collected and washed with methanol (5 ml) and diethylether (20 ml) (yield 0.506 g, 0.44 mmol, 69.6%). X-ray quality crystals were obtained from acetonitrile solution of the title complex by slow evaporation.

Refinement

H atoms bonded to C were inserted at calculated positions with C—H distances of 0.95, 0.98 and 0.99 Å for aryl, methyl and methylene H atoms, respectively, and refined using a riding model with $U_{\text{iso}}(\text{H}) = 1.2$ or $1.5U_{\text{eq}}(\text{C})$. The H-atoms bonded to N-atoms were taken from the difference Fourier maps and were refined isotropically.

Figures




Fig. 1. Perspective view of centrosymmetric calcium complex $[\text{Ca}(\text{C}_{27}\text{H}_{30}\text{N}_4\text{O}_3)_2](\text{ClO}_4)_2(\text{CH}_3\text{CN})_2$; displacement ellipsoids are drawn at the 50% probability level and hydrogen atoms are omitted for clarity (Symmetry code (i) $-x + 1, -y, -z + 1$).




Fig. 2. Packing diagram viewed perpendicular to b , showing intramolecular hydrogen bonds and π - π (phenyl-phenyl) interactions.

Bis{tris[2-(2-oxidobenzylideneazaniumyl)ethyl]amine- $\kappa^3\text{O},\text{O}',\text{O}''$ }calcium bis(perchlorate) acetonitrile disolvate

Crystal data

$[\text{Ca}(\text{C}_{27}\text{H}_{30}\text{N}_4\text{O}_3)_2](\text{ClO}_4)_2 \cdot 2\text{C}_2\text{H}_3\text{N}$

$M_r = 1238.19$

Orthorhombic, $Pbca$

Hall symbol: $-P\ 2ac\ 2ab$

$a = 11.3469\ (7)\ \text{\AA}$

$b = 19.5307\ (12)\ \text{\AA}$

$c = 27.3178\ (16)\ \text{\AA}$

$V = 6054.0\ (6)\ \text{\AA}^3$

$Z = 4$

$F(000) = 2600$

$D_x = 1.358\ \text{Mg m}^{-3}$

Mo $K\alpha$ radiation, $\lambda = 0.71073\ \text{\AA}$

Cell parameters from 9950 reflections

$\theta = 2.2\text{--}27.4^\circ$

$\mu = 0.27\ \text{mm}^{-1}$

$T = 150\ \text{K}$

Block, yellow

$0.12 \times 0.10 \times 0.09\ \text{mm}$

Data collection

Bruker APEXII CCD
diffractometer

Radiation source: fine-focus sealed tube
graphite

φ and ω scans

Absorption correction: multi-scan
(*SADABS*; Bruker, 2008)

$T_{\min} = 0.662$, $T_{\max} = 0.746$

55467 measured reflections

6915 independent reflections

5810 reflections with $I > 2.0\sigma(I)$

$R_{\text{int}} = 0.030$

$\theta_{\max} = 27.5^\circ$, $\theta_{\min} = 1.5^\circ$

$h = -14 \rightarrow 14$

$k = -25 \rightarrow 25$

$l = -34 \rightarrow 35$

Refinement

Refinement on F^2

Least-squares matrix: full

Primary atom site location: structure-invariant direct methods

Secondary atom site location: difference Fourier map

$$R[F^2 > 2\sigma(F^2)] = 0.039$$

$$wR(F^2) = 0.110$$

$$S = 1.05$$

6915 reflections

398 parameters

0 restraints

Hydrogen site location: inferred from neighbouring sites

H atoms treated by a mixture of independent and constrained refinement

$$w = 1/[\sigma^2(F_o^2) + (0.0538P)^2 + 3.4246P]$$

$$\text{where } P = (F_o^2 + 2F_c^2)/3$$

$$(\Delta/\sigma)_{\max} = 0.002$$

$$\Delta\rho_{\max} = 0.60 \text{ e } \text{\AA}^{-3}$$

$$\Delta\rho_{\min} = -0.39 \text{ e } \text{\AA}^{-3}$$

Special details

Geometry. All e.s.d.'s (except the e.s.d. in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell e.s.d.'s are taken into account individually in the estimation of e.s.d.'s in distances, angles and torsion angles; correlations between e.s.d.'s in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell e.s.d.'s is used for estimating e.s.d.'s involving l.s. planes.

Refinement. Refinement of F^2 against ALL reflections. The weighted R -factor wR and goodness of fit S are based on F^2 , conventional R -factors R are based on F , with F set to zero for negative F^2 . The threshold expression of $F^2 > \sigma(F^2)$ is used only for calculating R -factors(gt) *etc.* and is not relevant to the choice of reflections for refinement. R -factors based on F^2 are statistically about twice as large as those based on F , and R -factors based on ALL data will be even larger.

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (\AA^2)

	<i>x</i>	<i>y</i>	<i>z</i>	$U_{\text{iso}}^*/U_{\text{eq}}$
Ca1	0.5000	0.0000	0.5000	0.01901 (10)
O1	0.38283 (10)	0.08611 (5)	0.53706 (4)	0.0260 (2)
C1	0.32707 (13)	0.13534 (8)	0.51499 (5)	0.0224 (3)
C2	0.25547 (15)	0.12409 (8)	0.47327 (6)	0.0291 (3)
H2	0.2438	0.0786	0.4619	0.035*
C3	0.20240 (16)	0.17746 (10)	0.44891 (7)	0.0349 (4)
H3	0.1557	0.1680	0.4208	0.042*
C4	0.21547 (17)	0.24547 (9)	0.46452 (7)	0.0395 (4)
H4	0.1795	0.2818	0.4469	0.047*
C5	0.28043 (16)	0.25839 (9)	0.50526 (7)	0.0340 (4)
H5	0.2879	0.3042	0.5166	0.041*
C6	0.33725 (13)	0.20497 (8)	0.53117 (6)	0.0239 (3)
C7	0.40674 (14)	0.22286 (8)	0.57213 (5)	0.0247 (3)
H7	0.4066	0.2695	0.5821	0.030*
N1	0.47075 (12)	0.18036 (7)	0.59710 (5)	0.0243 (3)
C8	0.53399 (15)	0.19940 (8)	0.64171 (6)	0.0277 (3)
H8A	0.5444	0.2497	0.6429	0.033*
H8B	0.6130	0.1779	0.6418	0.033*
C9	0.46415 (15)	0.17550 (8)	0.68610 (6)	0.0264 (3)
H9A	0.5005	0.1939	0.7163	0.032*
H9B	0.3825	0.1931	0.6840	0.032*
N2	0.46228 (11)	0.10039 (6)	0.68827 (5)	0.0228 (3)
C10	0.56302 (15)	0.07362 (9)	0.71617 (6)	0.0290 (3)

supplementary materials

H10A	0.5388	0.0659	0.7505	0.035*
H10B	0.6271	0.1080	0.7162	0.035*
C11	0.60888 (15)	0.00683 (9)	0.69464 (6)	0.0307 (4)
H11A	0.6682	-0.0133	0.7170	0.037*
H11B	0.5431	-0.0261	0.6913	0.037*
N3	0.66223 (12)	0.01909 (7)	0.64666 (5)	0.0283 (3)
C12	0.77320 (14)	0.03357 (9)	0.64084 (6)	0.0303 (3)
H12	0.8224	0.0329	0.6690	0.036*
C13	0.82570 (14)	0.05027 (8)	0.59528 (6)	0.0282 (3)
C14	0.94893 (15)	0.06200 (9)	0.59374 (7)	0.0362 (4)
H14	0.9933	0.0605	0.6232	0.043*
C15	1.00481 (16)	0.07539 (9)	0.55058 (8)	0.0385 (4)
H15	1.0872	0.0841	0.5499	0.046*
C16	0.93896 (16)	0.07617 (9)	0.50724 (7)	0.0353 (4)
H16	0.9780	0.0844	0.4770	0.042*
C17	0.81902 (16)	0.06529 (8)	0.50726 (6)	0.0313 (4)
H17	0.7772	0.0660	0.4771	0.038*
C18	0.75700 (14)	0.05306 (7)	0.55146 (6)	0.0253 (3)
O2	0.64376 (10)	0.04299 (6)	0.55239 (4)	0.0287 (2)
C19	0.34878 (15)	0.07321 (8)	0.70527 (6)	0.0280 (3)
H19A	0.3107	0.1072	0.7270	0.034*
H19B	0.3626	0.0311	0.7246	0.034*
C20	0.26672 (14)	0.05700 (8)	0.66278 (6)	0.0287 (3)
H20A	0.1863	0.0484	0.6754	0.034*
H20B	0.2630	0.0968	0.6404	0.034*
N4	0.30786 (12)	-0.00316 (6)	0.63575 (5)	0.0241 (3)
C21	0.29327 (13)	-0.06510 (8)	0.65229 (6)	0.0252 (3)
H21	0.2469	-0.0706	0.6810	0.030*
C22	0.34110 (13)	-0.12487 (8)	0.63089 (6)	0.0233 (3)
C23	0.31923 (15)	-0.18869 (8)	0.65401 (6)	0.0308 (4)
H23	0.2708	-0.1903	0.6824	0.037*
C24	0.36664 (16)	-0.24768 (9)	0.63612 (7)	0.0361 (4)
H24	0.3526	-0.2901	0.6521	0.043*
C25	0.43665 (16)	-0.24480 (9)	0.59373 (7)	0.0376 (4)
H25	0.4690	-0.2859	0.5809	0.045*
C26	0.45932 (16)	-0.18409 (9)	0.57042 (7)	0.0341 (4)
H26	0.5066	-0.1842	0.5417	0.041*
C27	0.41389 (13)	-0.12088 (8)	0.58811 (5)	0.0231 (3)
O3	0.43711 (10)	-0.06298 (6)	0.56747 (4)	0.0285 (3)
C28	0.8355 (2)	0.24381 (14)	0.68889 (8)	0.0537 (6)
C29	0.85639 (19)	0.31612 (11)	0.68130 (9)	0.0505 (5)
H29A	0.9341	0.3283	0.6943	0.076*
H29B	0.7955	0.3427	0.6982	0.076*
H29C	0.8538	0.3263	0.6462	0.076*
N5	0.8201 (3)	0.18674 (14)	0.69423 (9)	0.0870 (9)
H4A	0.3474 (18)	-0.0002 (10)	0.6103 (8)	0.034 (5)*
H3A	0.6222 (19)	0.0209 (11)	0.6199 (8)	0.040 (6)*
H1A	0.4716 (19)	0.1395 (12)	0.5863 (8)	0.046 (6)*
C11	0.96706 (4)	0.89879 (2)	0.698669 (15)	0.03209 (11)

O4	1.04850 (18)	0.84563 (10)	0.70855 (9)	0.0828 (6)
O5	1.02963 (14)	0.96257 (8)	0.69527 (6)	0.0547 (4)
O6	0.88221 (16)	0.90508 (10)	0.73665 (7)	0.0743 (6)
O7	0.90949 (18)	0.88552 (10)	0.65378 (6)	0.0699 (5)

Atomic displacement parameters (\AA^2)

	U^{11}	U^{22}	U^{33}	U^{12}	U^{13}	U^{23}
Ca1	0.0219 (2)	0.01807 (19)	0.01709 (19)	0.00034 (15)	0.00046 (15)	0.00076 (14)
O1	0.0295 (6)	0.0200 (5)	0.0285 (6)	0.0045 (4)	-0.0004 (4)	0.0005 (4)
C1	0.0202 (7)	0.0226 (7)	0.0243 (7)	0.0021 (6)	0.0052 (6)	0.0012 (6)
C2	0.0294 (8)	0.0270 (8)	0.0309 (8)	0.0006 (6)	-0.0015 (7)	-0.0034 (6)
C3	0.0317 (9)	0.0410 (10)	0.0321 (9)	0.0026 (7)	-0.0085 (7)	0.0006 (7)
C4	0.0426 (10)	0.0326 (9)	0.0432 (10)	0.0087 (8)	-0.0118 (8)	0.0084 (8)
C5	0.0383 (9)	0.0225 (8)	0.0413 (10)	0.0049 (7)	-0.0063 (7)	0.0016 (7)
C6	0.0253 (7)	0.0219 (7)	0.0245 (7)	0.0019 (6)	0.0014 (6)	0.0013 (6)
C7	0.0284 (8)	0.0207 (7)	0.0251 (7)	-0.0012 (6)	0.0041 (6)	0.0001 (6)
N1	0.0290 (7)	0.0219 (6)	0.0219 (6)	-0.0040 (5)	0.0004 (5)	0.0009 (5)
C8	0.0337 (8)	0.0272 (8)	0.0222 (7)	-0.0086 (6)	-0.0027 (6)	0.0023 (6)
C9	0.0342 (8)	0.0238 (7)	0.0211 (7)	-0.0018 (6)	0.0006 (6)	-0.0010 (6)
N2	0.0250 (6)	0.0220 (6)	0.0214 (6)	-0.0021 (5)	0.0007 (5)	0.0024 (5)
C10	0.0324 (9)	0.0321 (8)	0.0224 (7)	0.0001 (7)	-0.0039 (6)	0.0022 (6)
C11	0.0312 (8)	0.0319 (8)	0.0290 (8)	0.0031 (7)	-0.0017 (7)	0.0065 (6)
N3	0.0261 (7)	0.0308 (7)	0.0279 (7)	0.0026 (6)	-0.0029 (6)	0.0001 (6)
C12	0.0266 (8)	0.0315 (8)	0.0326 (8)	0.0039 (6)	-0.0062 (6)	-0.0042 (7)
C13	0.0246 (8)	0.0255 (7)	0.0344 (8)	0.0007 (6)	-0.0018 (6)	-0.0045 (6)
C14	0.0257 (8)	0.0388 (10)	0.0442 (10)	-0.0014 (7)	-0.0047 (7)	-0.0077 (8)
C15	0.0254 (8)	0.0340 (9)	0.0561 (12)	-0.0043 (7)	0.0050 (8)	-0.0066 (8)
C16	0.0367 (9)	0.0250 (8)	0.0442 (10)	-0.0025 (7)	0.0113 (8)	-0.0018 (7)
C17	0.0355 (9)	0.0232 (8)	0.0352 (9)	-0.0019 (6)	0.0016 (7)	0.0001 (6)
C18	0.0266 (7)	0.0164 (7)	0.0330 (8)	-0.0008 (6)	-0.0011 (6)	-0.0029 (6)
O2	0.0248 (6)	0.0308 (6)	0.0304 (6)	-0.0023 (4)	-0.0034 (4)	-0.0008 (5)
C19	0.0309 (8)	0.0266 (8)	0.0265 (8)	-0.0046 (6)	0.0083 (6)	-0.0034 (6)
C20	0.0237 (8)	0.0246 (8)	0.0377 (9)	0.0013 (6)	0.0047 (6)	-0.0053 (6)
N4	0.0224 (6)	0.0230 (6)	0.0269 (7)	-0.0008 (5)	0.0023 (5)	-0.0025 (5)
C21	0.0216 (7)	0.0280 (8)	0.0262 (7)	-0.0026 (6)	0.0026 (6)	-0.0006 (6)
C22	0.0226 (7)	0.0227 (7)	0.0245 (7)	-0.0018 (6)	0.0009 (6)	0.0010 (6)
C23	0.0315 (8)	0.0283 (8)	0.0328 (9)	-0.0072 (6)	0.0058 (7)	0.0050 (7)
C24	0.0385 (9)	0.0211 (8)	0.0486 (10)	-0.0063 (7)	0.0035 (8)	0.0059 (7)
C25	0.0402 (10)	0.0217 (8)	0.0509 (11)	-0.0003 (7)	0.0077 (8)	-0.0063 (7)
C26	0.0387 (9)	0.0281 (8)	0.0356 (9)	-0.0017 (7)	0.0125 (7)	-0.0049 (7)
C27	0.0241 (7)	0.0236 (7)	0.0217 (7)	-0.0019 (6)	-0.0004 (6)	0.0009 (6)
O3	0.0356 (6)	0.0242 (6)	0.0257 (6)	-0.0002 (5)	0.0054 (5)	0.0050 (4)
C28	0.0484 (12)	0.0695 (15)	0.0431 (11)	-0.0228 (11)	-0.0127 (9)	0.0133 (10)
C29	0.0410 (11)	0.0524 (13)	0.0582 (13)	0.0026 (9)	0.0053 (10)	0.0008 (10)
N5	0.105 (2)	0.0817 (17)	0.0743 (16)	-0.0495 (15)	-0.0320 (14)	0.0278 (13)
Cl1	0.0298 (2)	0.0318 (2)	0.0346 (2)	-0.00051 (15)	-0.00320 (16)	-0.00806 (16)
O4	0.0778 (13)	0.0586 (11)	0.1119 (17)	0.0288 (10)	-0.0151 (12)	0.0028 (11)

supplementary materials

O5	0.0497 (9)	0.0468 (9)	0.0675 (10)	-0.0183 (7)	0.0038 (7)	-0.0143 (7)
O6	0.0635 (11)	0.0894 (13)	0.0702 (12)	-0.0249 (10)	0.0329 (9)	-0.0277 (10)
O7	0.0905 (13)	0.0724 (11)	0.0469 (9)	-0.0234 (10)	-0.0246 (9)	-0.0098 (8)

Geometric parameters (Å, °)

Ca1—O2 ⁱ	2.3269 (11)	C13—C18	1.430 (2)
Ca1—O2	2.3269 (11)	C14—C15	1.364 (3)
Ca1—O3 ⁱ	2.3279 (11)	C14—H14	0.9500
Ca1—O3	2.3279 (11)	C15—C16	1.400 (3)
Ca1—O1 ⁱ	2.3709 (10)	C15—H15	0.9500
Ca1—O1	2.3709 (10)	C16—C17	1.377 (3)
O1—C1	1.2993 (18)	C16—H16	0.9500
C1—C2	1.417 (2)	C17—C18	1.418 (2)
C1—C6	1.435 (2)	C17—H17	0.9500
C2—C3	1.375 (2)	C18—O2	1.3001 (19)
C2—H2	0.9500	C19—C20	1.522 (2)
C3—C4	1.403 (3)	C19—H19A	0.9900
C3—H3	0.9500	C19—H19B	0.9900
C4—C5	1.358 (3)	C20—N4	1.4642 (19)
C4—H4	0.9500	C20—H20A	0.9900
C5—C6	1.416 (2)	C20—H20B	0.9900
C5—H5	0.9500	N4—C21	1.302 (2)
C6—C7	1.413 (2)	N4—H4A	0.83 (2)
C7—N1	1.297 (2)	C21—C22	1.414 (2)
C7—H7	0.9500	C21—H21	0.9500
N1—C8	1.462 (2)	C22—C23	1.419 (2)
N1—H1A	0.85 (2)	C22—C27	1.433 (2)
C8—C9	1.522 (2)	C23—C24	1.362 (2)
C8—H8A	0.9900	C23—H23	0.9500
C8—H8B	0.9900	C24—C25	1.405 (3)
C9—N2	1.4684 (19)	C24—H24	0.9500
C9—H9A	0.9900	C25—C26	1.370 (2)
C9—H9B	0.9900	C25—H25	0.9500
N2—C19	1.468 (2)	C26—C27	1.422 (2)
N2—C10	1.470 (2)	C26—H26	0.9500
C10—C11	1.523 (2)	C27—O3	1.2908 (18)
C10—H10A	0.9900	C28—N5	1.138 (3)
C10—H10B	0.9900	C28—C29	1.447 (3)
C11—N3	1.463 (2)	C29—H29A	0.9800
C11—H11A	0.9900	C29—H29B	0.9800
C11—H11B	0.9900	C29—H29C	0.9800
N3—C12	1.300 (2)	C11—O7	1.4135 (16)
N3—H3A	0.86 (2)	C11—O4	1.4158 (18)
C12—C13	1.418 (2)	C11—O6	1.4209 (16)
C12—H12	0.9500	C11—O5	1.4368 (15)
C13—C14	1.418 (2)		
O2 ⁱ —Ca1—O2	180.00 (4)	N3—C12—H12	117.8

O2 ⁱ —Ca1—O3 ⁱ	85.33 (4)	C13—C12—H12	117.8
O2—Ca1—O3 ⁱ	94.67 (4)	C14—C13—C12	118.54 (16)
O2 ⁱ —Ca1—O3	94.67 (4)	C14—C13—C18	120.45 (16)
O2—Ca1—O3	85.33 (4)	C12—C13—C18	120.98 (14)
O3 ⁱ —Ca1—O3	180.0	C15—C14—C13	121.02 (17)
O2 ⁱ —Ca1—O1 ⁱ	82.79 (4)	C15—C14—H14	119.5
O2—Ca1—O1 ⁱ	97.21 (4)	C13—C14—H14	119.5
O3 ⁱ —Ca1—O1 ⁱ	82.23 (4)	C14—C15—C16	119.01 (16)
O3—Ca1—O1 ⁱ	97.77 (4)	C14—C15—H15	120.5
O2 ⁱ —Ca1—O1	97.21 (4)	C16—C15—H15	120.5
O2—Ca1—O1	82.79 (4)	C17—C16—C15	121.68 (17)
O3 ⁱ —Ca1—O1	97.77 (4)	C17—C16—H16	119.2
O3—Ca1—O1	82.23 (4)	C15—C16—H16	119.2
O1 ⁱ —Ca1—O1	180.00 (5)	C16—C17—C18	121.11 (17)
C1—O1—Ca1	126.86 (9)	C16—C17—H17	119.4
O1—C1—C2	122.54 (14)	C18—C17—H17	119.4
O1—C1—C6	121.28 (14)	O2—C18—C17	122.19 (15)
C2—C1—C6	116.17 (13)	O2—C18—C13	121.11 (14)
C3—C2—C1	121.52 (15)	C17—C18—C13	116.68 (15)
C3—C2—H2	119.2	C18—O2—Ca1	137.34 (10)
C1—C2—H2	119.2	N2—C19—C20	111.75 (13)
C2—C3—C4	121.61 (16)	N2—C19—H19A	109.3
C2—C3—H3	119.2	C20—C19—H19A	109.3
C4—C3—H3	119.2	N2—C19—H19B	109.3
C5—C4—C3	118.83 (16)	C20—C19—H19B	109.3
C5—C4—H4	120.6	H19A—C19—H19B	107.9
C3—C4—H4	120.6	N4—C20—C19	110.89 (13)
C4—C5—C6	121.31 (16)	N4—C20—H20A	109.5
C4—C5—H5	119.3	C19—C20—H20A	109.5
C6—C5—H5	119.3	N4—C20—H20B	109.5
C7—C6—C5	117.89 (14)	C19—C20—H20B	109.5
C7—C6—C1	121.57 (14)	H20A—C20—H20B	108.0
C5—C6—C1	120.51 (14)	C21—N4—C20	122.02 (14)
N1—C7—C6	124.81 (14)	C21—N4—H4A	115.1 (13)
N1—C7—H7	117.6	C20—N4—H4A	122.6 (14)
C6—C7—H7	117.6	N4—C21—C22	125.08 (14)
C7—N1—C8	123.40 (14)	N4—C21—H21	117.5
C7—N1—H1A	115.1 (15)	C22—C21—H21	117.5
C8—N1—H1A	121.5 (15)	C21—C22—C23	118.28 (14)
N1—C8—C9	109.29 (13)	C21—C22—C27	120.90 (14)
N1—C8—H8A	109.8	C23—C22—C27	120.76 (14)
C9—C8—H8A	109.8	C24—C23—C22	120.93 (15)
N1—C8—H8B	109.8	C24—C23—H23	119.5
C9—C8—H8B	109.8	C22—C23—H23	119.5
H8A—C8—H8B	108.3	C23—C24—C25	119.02 (16)
N2—C9—C8	110.25 (13)	C23—C24—H24	120.5
N2—C9—H9A	109.6	C25—C24—H24	120.5

supplementary materials

C8—C9—H9A	109.6	C26—C25—C24	121.57 (16)
N2—C9—H9B	109.6	C26—C25—H25	119.2
C8—C9—H9B	109.6	C24—C25—H25	119.2
H9A—C9—H9B	108.1	C25—C26—C27	121.68 (16)
C19—N2—C9	112.74 (12)	C25—C26—H26	119.2
C19—N2—C10	112.92 (12)	C27—C26—H26	119.2
C9—N2—C10	111.41 (12)	O3—C27—C26	122.55 (14)
N2—C10—C11	111.72 (13)	O3—C27—C22	121.43 (14)
N2—C10—H10A	109.3	C26—C27—C22	116.02 (14)
C11—C10—H10A	109.3	C27—O3—Ca1	150.62 (10)
N2—C10—H10B	109.3	N5—C28—C29	178.9 (3)
C11—C10—H10B	109.3	C28—C29—H29A	109.5
H10A—C10—H10B	107.9	C28—C29—H29B	109.5
N3—C11—C10	110.31 (13)	H29A—C29—H29B	109.5
N3—C11—H11A	109.6	C28—C29—H29C	109.5
C10—C11—H11A	109.6	H29A—C29—H29C	109.5
N3—C11—H11B	109.6	H29B—C29—H29C	109.5
C10—C11—H11B	109.6	O7—C11—O4	109.43 (13)
H11A—C11—H11B	108.1	O7—C11—O6	109.65 (12)
C12—N3—C11	123.08 (15)	O4—C11—O6	111.50 (14)
C12—N3—H3A	113.4 (14)	O7—C11—O5	109.34 (11)
C11—N3—H3A	123.3 (14)	O4—C11—O5	109.00 (12)
N3—C12—C13	124.35 (15)	O6—C11—O5	107.88 (10)

Symmetry codes: (i) $-x+1, -y, -z+1$.

Hydrogen-bond geometry (\AA , $^\circ$)

Cg1 is the centroid of the C1–C6 ring.

$D-H\cdots A$	$D-H$	$H\cdots A$	$D\cdots A$	$D-H\cdots A$
N1—H1A \cdots O1	0.85 (2)	1.98 (2)	2.6597 (17)	136 (2)
N3—H3A \cdots O2	0.86 (2)	1.91 (2)	2.6256 (18)	140 (2)
N4—H4A \cdots O3	0.83 (2)	1.98 (2)	2.6449 (17)	137 (2)
C3—H3 \cdots O7 ⁱⁱ	0.95	2.41	3.316 (2)	160
C20—H20A \cdots O5 ⁱⁱⁱ	0.99	2.50	3.380 (2)	148
C10—H10A \cdots O5 ^{iv}	0.99	2.50	3.271 (2)	134
C21—H21 \cdots O5 ⁱⁱⁱ	0.95	2.58	3.259 (2)	129
C29—H29B \cdots O6 ^v	0.98	2.58	3.555 (3)	173
C26—H26 \cdots Cg1 ⁱ	0.95	2.68	3.489 (2)	143

Symmetry codes: (ii) $-x+1, -y+1, -z+1$; (iii) $x-1, y-1, z$; (iv) $x-1/2, y-1, -z+3/2$; (v) $-x+3/2, y-1/2, z$; (i) $-x+1, -y, -z+1$.

Fig. 1




Fig. 2

