

Tetrakis(μ_2 -2-methyl-3,5-dinitrobenzoato- κ^2 O¹:O¹)bis[aquacopper(II)]tetrahydrate

Muhammad Danish,^a Sabiha Ghafoor,^a M. Nawaz Tahir,^{b,*} Nazir Ahmad^a and Mehwish Nisa^a

^aDepartment of Chemistry, University of Sargodha, Sargodha, Pakistan, and

^bDepartment of Physics, University of Sargodha, Sargodha, Pakistan

Correspondence e-mail: dmntahir_uos@yahoo.com


Received 4 January 2011; accepted 5 January 2011

Key indicators: single-crystal X-ray study; $T = 296$ K; mean $\sigma(C-C) = 0.003$ Å; R factor = 0.034; wR factor = 0.088; data-to-parameter ratio = 15.8.

The title compound, $[Cu_2(C_8H_5N_2O_6)_4(H_2O)_2] \cdot 4H_2O$, forms a centrosymmetric paddle-wheel-type dimer with an intramolecular Cu···Cu distance of 2.6540 (4) Å. The Cu^{II} atom is in a square-pyramidal coordination environment formed by four O atoms of four carboxylate groups and one water molecule, which is located in the apical position. The carboxylate groups are twisted relative to the benzene rings by 11.09 (16) and 45.55 (19)°. The nitro groups are not coplanar with the parent aromatic rings [dihedral angles = 16.2 (3)–51.45 (14)°]. O—H···O hydrogen bonds between the coordinated water molecules and one of the nitro groups, as well as π – π stacking interactions [centroid–centroid distance = 3.5764 (12) Å] between the benzene rings, assemble the complex molecules into a one-dimensional polymeric structure which is further extended into a three-dimensional polymeric network via O—H···O hydrogen bonds involving the water molecules of crystallization.

Related literature

For related crystal structures, see: Chen *et al.* (2007); Danish *et al.* (2010); Moncol *et al.* (2006); Stachova *et al.* (2004); Viossat *et al.* (2005).



Experimental

Crystal data

$[Cu_2(C_8H_5N_2O_6)_4(H_2O)_2] \cdot 4H_2O$	$V = 2209.75$ (12) Å ³
$M_r = 1135.76$	$Z = 2$
Monoclinic, $P2_1/n$	Mo $K\alpha$ radiation
$a = 8.9757$ (3) Å	$\mu = 1.08$ mm ⁻¹
$b = 22.5582$ (8) Å	$T = 296$ K
$c = 11.2698$ (3) Å	$0.30 \times 0.24 \times 0.22$ mm
$\beta = 104.443$ (1)°	

Data collection

Bruker Kappa APEXII CCD diffractometer	21133 measured reflections
Absorption correction: multi-scan (<i>SADABS</i> ; Bruker, 2005)	5449 independent reflections
$T_{\min} = 0.743$, $T_{\max} = 0.782$	4344 reflections with $I > 2\sigma(I)$
	$R_{\text{int}} = 0.035$

Refinement

$R[F^2 > 2\sigma(F^2)] = 0.034$	H atoms treated by a mixture of independent and constrained refinement
$wR(F^2) = 0.088$	$\Delta\rho_{\text{max}} = 0.33$ e Å ⁻³
$S = 1.02$	$\Delta\rho_{\text{min}} = -0.28$ e Å ⁻³
5449 reflections	
345 parameters	
6 restraints	

Table 1
Selected bond lengths (Å).

Cu1—O1	1.9616 (16)	Cu1—O13	2.0914 (19)
Cu1—O7	1.9749 (16)	Cu1—O2 ⁱ	1.9595 (16)
Cu1—O8	1.9637 (15)		

Symmetry code: (i) $-x, -y, -z + 2$.

Table 2
Hydrogen-bond geometry (Å, °).

$D-H \cdots A$	$D-H$	$H \cdots A$	$D \cdots A$	$D-H \cdots A$
O13—H13A···O14 ⁱⁱ	0.80 (3)	1.84 (3)	2.641 (3)	175 (3)
O13—H13B···O10 ⁱⁱⁱ	0.77 (2)	2.22 (2)	2.838 (2)	138 (3)
O14—H14A···O15 ^{iv}	0.81 (3)	1.95 (3)	2.758 (4)	172 (3)
O14—H14B···O7	0.80 (3)	2.51 (4)	3.182 (3)	143 (4)
O15—H15A···O12 ^v	0.83 (4)	2.16 (4)	2.953 (4)	161 (4)
O15—H15B···O5 ^{vi}	0.84 (3)	2.17 (2)	2.996 (4)	168 (4)
O15—H15B···O6 ^{vi}	0.84 (3)	2.60 (4)	3.273 (4)	138 (4)

Symmetry codes: (ii) $-x + 1, -y, -z + 2$; (iii) $x, y, z + 1$; (iv) $x - \frac{1}{2}, -y + \frac{1}{2}, z + \frac{1}{2}$; (v) $x + 1, y, z$; (vi) $x, y, z - 1$.

Data collection: *APEX2* (Bruker, 2009); cell refinement: *SAINT* (Bruker, 2009); data reduction: *SAINT*; program(s) used to solve structure: *SHELXS97* (Sheldrick, 2008); program(s) used to refine structure: *SHELXL97* (Sheldrick, 2008); molecular graphics: *ORTEP-3 for Windows* (Farrugia, 1997) and *PLATON* (Spek, 2009); software used to prepare material for publication: *WinGX* (Farrugia, 1999) and *PLATON*.

The authors acknowledge the provision of funds for the purchase of the diffractometer and encouragement by Dr Muhammad Akram Chaudhary, Vice Chancellor, University of Sargodha, Pakistan.

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: GK2339).

References

- Bruker (2005). *SADABS*. Bruker AXS Inc., Madison, Wisconsin, USA.
- Bruker (2009). *APEX2* and *SAINT*. Bruker AXS Inc., Madison, Wisconsin, USA.
- Chen, P.-G., Gao, S. & Ng, S. W. (2007). *Acta Cryst. E* **63**, m2617.
- Danish, M., Saleem, I., Tahir, M. N., Ahmad, N. & Raza, A. R. (2010). *Acta Cryst. E* **66**, m528.
- Farrugia, L. J. (1997). *J. Appl. Cryst.* **30**, 565.
- Farrugia, L. J. (1999). *J. Appl. Cryst.* **32**, 837–838.
- Moncol, J., Kavalírová, J., Lis, T. & Valigura, D. (2006). *Acta Cryst. E* **62**, m3217–m3219.
- Sheldrick, G. M. (2008). *Acta Cryst. A* **64**, 112–122.
- Spek, A. L. (2009). *Acta Cryst. D* **65**, 148–155.
- Stachova, P., Valigura, D., Koman, M., Melnik, M., Korabik, M., Mrozinski, J. & Glowiacz, T. (2004). *Polyhedron*, **23**, 1303–1308.
- Viossat, B., Greenaway, F. T., Morgant, G., Daran, J.-C., Dung, N.-H. & Sorenson, J. R. J. (2005). *J. Inorg. Biochem.* **99**, 355–367.

supplementary materials

Acta Cryst. (2011). E67, m168-m169 [doi:10.1107/S1600536811000547]

Tetrakis(μ_2 -2-methyl-3,5-dinitrobenzoato- $\kappa^2O^1:O^1'$)bis[aquacopper(II)] tetrahydrate

M. Danish, S. Ghafoor, M. N. Tahir, N. Ahmad and M. Nisa

Comment

Recently we have reported the synthesis and crystal structure of tetrakis(μ_2 -2-methylbenzoato- $\kappa^2O:O'$)bis[(methanol-O)copper(II)]. In continuation to our interest with the metal carboxylate chemistry, the title compound (I), (Fig. 1) is being reported here.

The crystal structure of (II) *i.e.*, diaqua-tetrakis(μ_2 -2,3,5-tri-iodobenzoato- $\kappa^2O:O'$)-di-copper(ii) bis(methanol- κO)-tetrakis(μ_2 -2,3,5-tri-iodobenzoato- $\kappa^2O:O'$)-di-copper(ii) methanol solvate (Chen *et al.*, 2007), (III) *i.e.*, tetrakis(μ_2 -2-nitrobenzoato- O,O')-bis(aqua-copper(ii)) ethanol solvate (Moncol *et al.*, 2006), (IV) *i.e.*, tetrakis(μ_2 -2-(3-trifluoromethylphenyl)aminonicotinato)-diaqua-di-copper(ii) *N,N*-dimethylformamide solvate (Viessat *et al.*, 2005) and (V) *i.e.*, diaqua-tetrakis(2-nitrobenzoato- O,O')-di-copper(ii) dihydrate (Stachova *et al.*, 2004) have been published which are related to the title compound (I) due to the coordination around copper.

The structure of (I) is centrosymmetric with a square pyramidal coordination around the Cu-atoms. Four oxygen atoms ($O1/O2^i/O7/O8$; $i = -x, -y, -z + 2$) from four carboxylate groups are in plane [r. m.s deviation of 0.0014 Å] and coordinated to copper(II) with bond distance of 1.9593 (15)–1.9749 (15) Å. The Cu—O [2.0909 (17) Å] bond of water molecule is longer as compared to Cu—O bonds of carboxylate groups. The Cu atom is at a distance of 0.2189 (8) Å from the mean square plane of carboxylate O-atoms. The separation of Cu—Cu is 2.6540 (4) Å. These values are in agreement with the reported structures [Table 1]. The toluene groups A (C2—C8) and B (C10—C16) are almost planar with r. m.s deviation of 0.0524 Å and 0.0363 Å and are nearly perpendicular to each other. The dihedral angle between A/B is 89.33 (6)°. The carboxylate moiety C (O1/C1/O2), nitro groups D (O3/N1/O4) and E (O5/N2/O6) are oriented at dihedral angles of 11.09 (16), 51.45 (14) and 45.30 (33)°, respectively with the parent toluene (A) moiety. In the other ligand, the carboxylate moiety F (O7/C9/O8^j), nitro groups G (O9/N3/O10) and H (O11/N4/O12) are oriented at dihedral angles of 45.55 (19), 40.43 (23) and 16.20 (34)°, respectively with the parent toluene (B) moiety. The molecules are stabilized in the form of three-dimensional polymeric network due to strong H-bonding (Table 2, Fig. 2, Fig. 3). There also exists a $\pi\cdots\pi$ interaction between the benzene rings (C10—C15) with centroid-to- centroid distance of 3.5764 (12) Å.

Experimental

Aqueous solutions of sodium salt of 3,5-dinitro-*o*-toluic acid (0.496 g, 2.0 mol) and copper chloride dihydrate (0.170 g, 1.0 mmol) were prepared separately. Both solutions were mixed in 100 ml round-bottom flask at room temperature. On mixing dirty green precipitate was formed at the begining and it disappeared on continuous stirring of the reaction mixture for 4 h. The reaction mixture was filtered and the filtrate was concentrated by heating for few minutes. The concentrated solution was kept at room temperature for 72 h to afford greenish blue prisms of of the title compound.

supplementary materials

Refinement

The coordinates of H-atoms of water molecules were refined with the distance restraints imposed on the O-H distances and with $U_{\text{iso}}(\text{H}) = 1.2 U_{\text{eq}}(\text{O})$. The C-bound H atoms were positioned geometrically ($\text{C}-\text{H} = 0.93\text{--}0.96 \text{\AA}$) and refined as riding with $U_{\text{iso}}(\text{H}) = x U_{\text{eq}}(\text{C})$, where $x = 1.5$ for methyl and $x = 1.2$ for other H-atoms.

Figures




Fig. 1. View of the title compound with the atom numbering scheme. The displacement ellipsoids are drawn at the 50% probability level. H-atoms are shown as small spheres of arbitrary radii.




Fig. 2. The partial packing (*PLATON*; Spek, 2009) which shows the H-bonding mode with neighbouring molecules through dotted lines except which are present in coordination sphere.




Fig. 3. The partial packing (*PLATON*; Spek, 2009) which shows that molecules form three-dimensional polymeric network due to strong intermolecular H-bondings.

Tetrakis(μ_2 -2-methyl-3,5-dinitrobenzoato- $\kappa^2\text{O}^1:\text{O}^1'$)bis[aquacopper(II)] tetrahydrate

Crystal data

$[\text{Cu}_2(\text{C}_8\text{H}_5\text{N}_2\text{O}_6)_4(\text{H}_2\text{O})_2]\cdot 4\text{H}_2\text{O}$	$F(000) = 1156$
$M_r = 1135.76$	$D_x = 1.707 \text{ Mg m}^{-3}$
Monoclinic, $P2_1/n$	Mo $K\alpha$ radiation, $\lambda = 0.71073 \text{ \AA}$
Hall symbol: -P 2yn	Cell parameters from 4344 reflections
$a = 8.9757 (3) \text{ \AA}$	$\theta = 2.1\text{--}28.3^\circ$
$b = 22.5582 (8) \text{ \AA}$	$\mu = 1.08 \text{ mm}^{-1}$
$c = 11.2698 (3) \text{ \AA}$	$T = 296 \text{ K}$
$\beta = 104.443 (1)^\circ$	Prism, blue
$V = 2209.75 (12) \text{ \AA}^3$	$0.30 \times 0.24 \times 0.22 \text{ mm}$
$Z = 2$	

Data collection

Bruker Kappa APEXII CCD diffractometer	5449 independent reflections
Radiation source: fine-focus sealed tube graphite	4344 reflections with $I > 2\sigma(I)$
Detector resolution: 7.50 pixels mm ⁻¹	$R_{\text{int}} = 0.035$
ω scans	$\theta_{\text{max}} = 28.3^\circ$, $\theta_{\text{min}} = 2.1^\circ$
Absorption correction: multi-scan (<i>SADABS</i> ; Bruker, 2005)	$h = -11 \rightarrow 11$
$T_{\text{min}} = 0.743$, $T_{\text{max}} = 0.782$	$k = -30 \rightarrow 30$
21133 measured reflections	$l = -15 \rightarrow 9$

Refinement

Refinement on F^2	Primary atom site location: structure-invariant direct methods
Least-squares matrix: full	Secondary atom site location: difference Fourier map
$R[F^2 > 2\sigma(F^2)] = 0.034$	Hydrogen site location: inferred from neighbouring sites
$wR(F^2) = 0.088$	H atoms treated by a mixture of independent and constrained refinement
$S = 1.02$	$w = 1/[\sigma^2(F_o^2) + (0.0398P)^2 + 0.8547P]$ where $P = (F_o^2 + 2F_c^2)/3$
5449 reflections	$(\Delta/\sigma)_{\text{max}} = 0.001$
345 parameters	$\Delta\rho_{\text{max}} = 0.33 \text{ e \AA}^{-3}$
6 restraints	$\Delta\rho_{\text{min}} = -0.28 \text{ e \AA}^{-3}$

Special details

Geometry. Bond distances, angles etc. have been calculated using the rounded fractional coordinates. All su's are estimated from the variances of the (full) variance-covariance matrix. The cell e.s.d.'s are taken into account in the estimation of distances, angles and torsion angles

Refinement. Refinement of F^2 against ALL reflections. The weighted R -factor wR and goodness of fit S are based on F^2 , conventional R -factors R are based on F , with F set to zero for negative F^2 . The threshold expression of $F^2 > \sigma(F^2)$ is used only for calculating R -factors(gt) etc. and is not relevant to the choice of reflections for refinement. R -factors based on F^2 are statistically about twice as large as those based on F , and R -factors based on ALL data will be even larger.

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (Å²)

	<i>x</i>	<i>y</i>	<i>z</i>	$U_{\text{iso}}^*/U_{\text{eq}}$
Cu1	0.14133 (3)	-0.01627 (1)	1.05844 (2)	0.0264 (1)
O1	0.17167 (19)	0.06719 (7)	1.10736 (17)	0.0528 (6)
O2	-0.06417 (19)	0.09368 (7)	1.00866 (15)	0.0474 (5)
O3	-0.1238 (3)	0.32675 (9)	0.9849 (2)	0.0826 (9)
O4	-0.0417 (3)	0.36016 (8)	1.1678 (2)	0.0701 (8)
O5	0.5065 (2)	0.29719 (9)	1.30010 (18)	0.0624 (7)

supplementary materials

O6	0.5609 (2)	0.20680 (10)	1.2665 (3)	0.0901 (9)
O7	0.19256 (18)	0.00393 (8)	0.90275 (14)	0.0461 (5)
O8	0.04616 (18)	-0.02830 (8)	1.19572 (13)	0.0445 (5)
O9	0.2910 (2)	-0.08610 (9)	0.45982 (19)	0.0680 (8)
O10	0.3870 (2)	-0.01075 (10)	0.38692 (16)	0.0623 (7)
O11	0.0204 (3)	0.15433 (11)	0.34615 (17)	0.0807 (9)
O12	-0.0072 (3)	0.18886 (9)	0.51609 (19)	0.0792 (9)
O13	0.3664 (2)	-0.04335 (10)	1.14010 (15)	0.0592 (7)
N1	-0.0457 (3)	0.32313 (9)	1.0898 (2)	0.0496 (7)
N2	0.4706 (2)	0.24745 (10)	1.2591 (2)	0.0518 (7)
N3	0.3119 (2)	-0.03328 (11)	0.45226 (17)	0.0472 (7)
N4	0.0329 (3)	0.15041 (10)	0.45558 (18)	0.0510 (7)
C1	0.0668 (2)	0.10469 (9)	1.07283 (18)	0.0321 (6)
C2	0.1073 (2)	0.16812 (8)	1.10974 (17)	0.0309 (6)
C3	-0.0028 (3)	0.21412 (9)	1.08940 (18)	0.0347 (6)
C4	0.0571 (3)	0.27102 (9)	1.1212 (2)	0.0373 (6)
C5	0.2074 (3)	0.28347 (10)	1.1774 (2)	0.0400 (7)
C6	0.3078 (3)	0.23623 (10)	1.19871 (19)	0.0377 (7)
C7	0.2606 (3)	0.17952 (9)	1.16425 (18)	0.0350 (6)
C8	-0.1731 (3)	0.20411 (11)	1.0466 (3)	0.0523 (8)
C9	0.0935 (2)	0.02030 (8)	0.81066 (18)	0.0308 (6)
C10	0.1378 (2)	0.03256 (9)	0.69218 (17)	0.0303 (6)
C11	0.2268 (2)	-0.00702 (9)	0.64135 (18)	0.0333 (6)
C12	0.2367 (2)	0.00750 (10)	0.52214 (19)	0.0353 (6)
C13	0.1764 (2)	0.05764 (10)	0.45945 (18)	0.0379 (7)
C14	0.0982 (2)	0.09566 (10)	0.51708 (18)	0.0355 (6)
C15	0.0760 (2)	0.08360 (9)	0.63150 (18)	0.0341 (6)
C16	0.3062 (3)	-0.05950 (12)	0.7111 (2)	0.0517 (8)
O14	0.4622 (3)	0.09823 (11)	0.9851 (3)	0.0827 (10)
O15	0.8508 (3)	0.29497 (14)	0.3805 (4)	0.1215 (14)
H5	0.24001	0.32188	1.20013	0.0480*
H7	0.33171	0.14872	1.17755	0.0421*
H8A	-0.20137	0.19949	0.95920	0.0784*
H8B	-0.22654	0.23750	1.06903	0.0784*
H8C	-0.20028	0.16893	1.08435	0.0784*
H13	0.18803	0.06549	0.38126	0.0455*
H13A	0.419 (3)	-0.0616 (13)	1.105 (2)	0.0711*
H13B	0.389 (4)	-0.0520 (14)	1.2082 (17)	0.0711*
H15	0.01980	0.10960	0.66751	0.0409*
H16A	0.34251	-0.04933	0.79617	0.0776*
H16B	0.39179	-0.07080	0.67924	0.0776*
H16C	0.23523	-0.09198	0.70265	0.0776*
H14A	0.428 (5)	0.1281 (12)	0.948 (3)	0.0993*
H14B	0.410 (4)	0.0722 (14)	1.000 (4)	0.0993*
H15A	0.910 (5)	0.2681 (17)	0.414 (4)	0.1457*
H15B	0.755 (2)	0.291 (2)	0.352 (4)	0.1457*

Atomic displacement parameters (\AA^2)

	U^{11}	U^{22}	U^{33}	U^{12}	U^{13}	U^{23}
Cu1	0.0288 (1)	0.0251 (1)	0.0263 (1)	0.0023 (1)	0.0088 (1)	0.0015 (1)
O1	0.0498 (9)	0.0261 (8)	0.0741 (12)	0.0047 (7)	-0.0001 (8)	-0.0090 (8)
O2	0.0451 (9)	0.0282 (8)	0.0629 (10)	0.0005 (7)	0.0023 (8)	-0.0085 (7)
O3	0.0964 (17)	0.0585 (13)	0.0761 (14)	0.0322 (12)	-0.0102 (12)	0.0002 (11)
O4	0.0857 (15)	0.0430 (11)	0.0871 (14)	0.0112 (10)	0.0318 (12)	-0.0168 (10)
O5	0.0623 (12)	0.0517 (11)	0.0660 (12)	-0.0168 (9)	0.0024 (9)	-0.0118 (9)
O6	0.0472 (12)	0.0617 (14)	0.147 (2)	0.0052 (10)	-0.0027 (13)	-0.0153 (14)
O7	0.0371 (8)	0.0709 (11)	0.0326 (8)	0.0032 (8)	0.0129 (7)	0.0138 (8)
O8	0.0399 (9)	0.0664 (11)	0.0310 (8)	0.0109 (8)	0.0160 (6)	0.0103 (7)
O9	0.0780 (14)	0.0575 (13)	0.0740 (13)	0.0053 (11)	0.0294 (11)	-0.0234 (10)
O10	0.0497 (10)	0.1042 (16)	0.0394 (9)	-0.0029 (10)	0.0233 (8)	-0.0130 (10)
O11	0.1012 (17)	0.1012 (18)	0.0450 (10)	0.0273 (14)	0.0284 (11)	0.0341 (11)
O12	0.126 (2)	0.0554 (12)	0.0644 (13)	0.0282 (13)	0.0394 (13)	0.0195 (10)
O13	0.0469 (10)	0.0925 (15)	0.0348 (9)	0.0330 (10)	0.0037 (8)	-0.0047 (10)
N1	0.0556 (13)	0.0312 (10)	0.0638 (14)	0.0042 (9)	0.0181 (11)	-0.0027 (10)
N2	0.0475 (12)	0.0485 (12)	0.0554 (13)	-0.0071 (10)	0.0054 (10)	-0.0010 (10)
N3	0.0392 (10)	0.0671 (15)	0.0365 (10)	0.0007 (10)	0.0116 (8)	-0.0167 (10)
N4	0.0581 (13)	0.0568 (13)	0.0418 (11)	0.0040 (10)	0.0192 (10)	0.0180 (10)
C1	0.0438 (12)	0.0261 (10)	0.0300 (10)	-0.0011 (8)	0.0162 (9)	-0.0007 (8)
C2	0.0423 (11)	0.0244 (9)	0.0288 (9)	-0.0011 (8)	0.0144 (8)	-0.0003 (8)
C3	0.0439 (12)	0.0298 (10)	0.0324 (10)	0.0012 (9)	0.0134 (9)	-0.0002 (8)
C4	0.0473 (12)	0.0274 (10)	0.0387 (11)	0.0048 (9)	0.0133 (10)	0.0005 (9)
C5	0.0531 (13)	0.0263 (10)	0.0408 (12)	-0.0050 (9)	0.0123 (10)	-0.0014 (9)
C6	0.0409 (12)	0.0357 (11)	0.0357 (11)	-0.0043 (9)	0.0081 (9)	-0.0010 (9)
C7	0.0417 (12)	0.0291 (10)	0.0361 (11)	0.0041 (9)	0.0131 (9)	0.0011 (9)
C8	0.0424 (13)	0.0396 (13)	0.0738 (17)	0.0033 (10)	0.0125 (12)	-0.0054 (12)
C9	0.0393 (11)	0.0265 (10)	0.0292 (10)	-0.0038 (8)	0.0137 (8)	-0.0023 (8)
C10	0.0339 (10)	0.0323 (10)	0.0264 (9)	-0.0059 (8)	0.0109 (8)	-0.0009 (8)
C11	0.0328 (10)	0.0353 (11)	0.0337 (10)	-0.0048 (8)	0.0121 (8)	-0.0043 (9)
C12	0.0330 (10)	0.0435 (12)	0.0319 (10)	-0.0068 (9)	0.0130 (8)	-0.0106 (9)
C13	0.0374 (11)	0.0503 (13)	0.0278 (10)	-0.0104 (10)	0.0115 (8)	-0.0022 (9)
C14	0.0378 (11)	0.0396 (12)	0.0297 (10)	-0.0058 (9)	0.0096 (8)	0.0041 (9)
C15	0.0378 (11)	0.0350 (11)	0.0318 (10)	-0.0043 (9)	0.0130 (9)	-0.0016 (9)
C16	0.0622 (16)	0.0480 (14)	0.0484 (14)	0.0116 (12)	0.0202 (12)	0.0036 (12)
O14	0.0699 (16)	0.0818 (18)	0.111 (2)	0.0008 (13)	0.0501 (14)	-0.0127 (15)
O15	0.0781 (19)	0.086 (2)	0.186 (3)	-0.0088 (16)	0.006 (2)	0.035 (2)

Geometric parameters (\AA , $^\circ$)

Cu1—O1	1.9616 (16)	C1—C2	1.509 (3)
Cu1—O7	1.9749 (16)	C2—C3	1.412 (3)
Cu1—O8	1.9637 (15)	C2—C7	1.384 (3)
Cu1—O13	2.0914 (19)	C3—C4	1.403 (3)
Cu1—O2 ⁱ	1.9595 (16)	C3—C8	1.501 (4)

supplementary materials

O1—C1	1.253 (3)	C4—C5	1.369 (4)
O2—C1	1.242 (3)	C5—C6	1.378 (3)
O3—N1	1.218 (3)	C6—C7	1.373 (3)
O4—N1	1.207 (3)	C9—C10	1.511 (3)
O5—N2	1.226 (3)	C10—C15	1.383 (3)
O6—N2	1.213 (3)	C10—C11	1.411 (3)
O7—C9	1.243 (2)	C11—C12	1.407 (3)
O8—C9 ⁱ	1.251 (2)	C11—C16	1.499 (3)
O9—N3	1.213 (3)	C12—C13	1.371 (3)
O10—N3	1.226 (3)	C13—C14	1.370 (3)
O11—N4	1.213 (3)	C14—C15	1.380 (3)
O12—N4	1.212 (3)	C5—H5	0.9300
O13—H13B	0.77 (2)	C7—H7	0.9300
O13—H13A	0.80 (3)	C8—H8A	0.9600
O14—H14A	0.81 (3)	C8—H8B	0.9600
O14—H14B	0.80 (3)	C8—H8C	0.9600
O15—H15A	0.83 (4)	C13—H13	0.9300
O15—H15B	0.84 (3)	C15—H15	0.9300
N1—C4	1.482 (3)	C16—H16B	0.9600
N2—C6	1.472 (3)	C16—H16C	0.9600
N3—C12	1.479 (3)	C16—H16A	0.9600
N4—C14	1.465 (3)		
O1—Cu1—O7	88.94 (7)	C4—C5—C6	116.6 (2)
O1—Cu1—O8	88.19 (7)	C5—C6—C7	122.0 (2)
O1—Cu1—O13	96.06 (8)	N2—C6—C7	119.5 (2)
O1—Cu1—O2 ⁱ	167.09 (7)	N2—C6—C5	118.6 (2)
O7—Cu1—O8	167.24 (7)	C2—C7—C6	120.1 (2)
O7—Cu1—O13	92.58 (7)	O8 ⁱ —C9—C10	113.81 (17)
O2 ⁱ —Cu1—O7	90.06 (7)	O7—C9—C10	120.09 (17)
O8—Cu1—O13	100.09 (7)	O7—C9—O8 ⁱ	126.09 (19)
O2 ⁱ —Cu1—O8	89.98 (7)	C11—C10—C15	121.60 (18)
O2 ⁱ —Cu1—O13	96.84 (8)	C9—C10—C15	115.34 (17)
Cu1—O1—C1	121.32 (15)	C9—C10—C11	122.91 (17)
Cu1 ⁱ —O2—C1	126.52 (14)	C10—C11—C16	122.00 (18)
Cu1—O7—C9	122.28 (14)	C10—C11—C12	114.66 (18)
Cu1—O8—C9 ⁱ	124.16 (13)	C12—C11—C16	123.33 (19)
Cu1—O13—H13A	123.8 (17)	N3—C12—C13	114.37 (18)
Cu1—O13—H13B	120 (3)	C11—C12—C13	124.97 (19)
H13A—O13—H13B	109 (3)	N3—C12—C11	120.64 (19)
H14A—O14—H14B	124 (4)	C12—C13—C14	117.07 (19)
H15A—O15—H15B	125 (4)	C13—C14—C15	122.0 (2)
O3—N1—O4	124.5 (2)	N4—C14—C13	119.68 (19)
O4—N1—C4	118.1 (2)	N4—C14—C15	118.31 (19)
O3—N1—C4	117.3 (2)	C10—C15—C14	119.52 (18)
O5—N2—C6	118.0 (2)	C6—C5—H5	122.00
O6—N2—C6	118.1 (2)	C4—C5—H5	122.00
O5—N2—O6	123.9 (2)	C2—C7—H7	120.00

O9—N3—O10	124.6 (2)	C6—C7—H7	120.00
O10—N3—C12	117.0 (2)	C3—C8—H8A	109.00
O9—N3—C12	118.34 (19)	H8A—C8—H8B	110.00
O12—N4—C14	118.49 (19)	H8A—C8—H8C	109.00
O11—N4—O12	123.7 (2)	H8B—C8—H8C	109.00
O11—N4—C14	117.8 (2)	C3—C8—H8B	109.00
O1—C1—C2	116.35 (17)	C3—C8—H8C	109.00
O1—C1—O2	125.1 (2)	C12—C13—H13	121.00
O2—C1—C2	118.56 (18)	C14—C13—H13	121.00
C1—C2—C7	116.30 (17)	C10—C15—H15	120.00
C3—C2—C7	120.91 (18)	C14—C15—H15	120.00
C1—C2—C3	122.78 (17)	C11—C16—H16B	109.00
C4—C3—C8	121.0 (2)	C11—C16—H16C	109.00
C2—C3—C8	123.86 (19)	H16A—C16—H16C	109.00
C2—C3—C4	115.0 (2)	H16B—C16—H16C	109.00
N1—C4—C3	119.3 (2)	H16A—C16—H16B	109.00
N1—C4—C5	115.5 (2)	C11—C16—H16A	109.00
C3—C4—C5	125.2 (2)		
O7—Cu1—O1—C1	83.36 (17)	O2—C1—C2—C3	9.2 (3)
O8—Cu1—O1—C1	−84.20 (17)	O2—C1—C2—C7	−170.12 (19)
O13—Cu1—O1—C1	175.84 (17)	C1—C2—C3—C4	−175.44 (18)
O1—Cu1—O7—C9	−88.62 (17)	C1—C2—C3—C8	9.3 (3)
O13—Cu1—O7—C9	175.36 (17)	C7—C2—C3—C4	3.8 (3)
O2 ⁱ —Cu1—O7—C9	78.51 (17)	C7—C2—C3—C8	−171.5 (2)
O1—Cu1—O8—C9 ⁱ	87.39 (18)	C1—C2—C7—C6	178.87 (19)
O13—Cu1—O8—C9 ⁱ	−176.78 (17)	C3—C2—C7—C6	−0.4 (3)
O2 ⁱ —Cu1—O8—C9 ⁱ	−79.83 (17)	C2—C3—C4—N1	172.73 (19)
O7—Cu1—O2 ⁱ —C1 ⁱ	−84.61 (18)	C2—C3—C4—C5	−5.4 (3)
O8—Cu1—O2 ⁱ —C1 ⁱ	82.62 (18)	C8—C3—C4—N1	−11.8 (3)
O13—Cu1—O2 ⁱ —C1 ⁱ	−177.22 (17)	C8—C3—C4—C5	170.1 (2)
Cu1—O1—C1—O2	1.7 (3)	N1—C4—C5—C6	−175.0 (2)
Cu1—O1—C1—C2	−176.21 (13)	C3—C4—C5—C6	3.2 (4)
Cu1 ⁱ —O2—C1—O1	−1.3 (3)	C4—C5—C6—N2	−179.9 (2)
Cu1 ⁱ —O2—C1—C2	176.49 (13)	C4—C5—C6—C7	0.8 (3)
Cu1—O7—C9—C10	−177.24 (13)	N2—C6—C7—C2	178.63 (19)
Cu1—O7—C9—O8 ⁱ	2.8 (3)	C5—C6—C7—C2	−2.0 (3)
Cu1—O8—C9 ⁱ —O7 ⁱ	0.1 (3)	O7—C9—C10—C11	48.5 (3)
Cu1—O8—C9 ⁱ —C10 ⁱ	−179.93 (13)	O7—C9—C10—C15	−135.9 (2)
O3—N1—C4—C3	−49.7 (3)	O8 ⁱ —C9—C10—C11	−131.5 (2)
O3—N1—C4—C5	128.6 (3)	O8 ⁱ —C9—C10—C15	44.1 (2)
O4—N1—C4—C3	133.4 (3)	C9—C10—C11—C12	170.77 (18)
O4—N1—C4—C5	−48.3 (3)	C9—C10—C11—C16	−10.4 (3)
O5—N2—C6—C5	8.0 (3)	C15—C10—C11—C12	−4.6 (3)
O5—N2—C6—C7	−172.6 (2)	C15—C10—C11—C16	174.3 (2)
O6—N2—C6—C5	−172.8 (3)	C9—C10—C15—C14	−174.02 (17)
O6—N2—C6—C7	6.6 (3)	C11—C10—C15—C14	1.7 (3)

supplementary materials

O9—N3—C12—C11	39.4 (3)	C10—C11—C12—N3	−173.40 (18)
O9—N3—C12—C13	−138.8 (2)	C10—C11—C12—C13	4.5 (3)
O10—N3—C12—C11	−143.2 (2)	C16—C11—C12—N3	7.7 (3)
O10—N3—C12—C13	38.7 (3)	C16—C11—C12—C13	−174.4 (2)
O11—N4—C14—C13	15.9 (4)	N3—C12—C13—C14	176.69 (18)
O11—N4—C14—C15	−162.9 (2)	C11—C12—C13—C14	−1.3 (3)
O12—N4—C14—C13	−165.0 (2)	C12—C13—C14—N4	179.2 (2)
O12—N4—C14—C15	16.2 (3)	C12—C13—C14—C15	−2.1 (3)
O1—C1—C2—C3	−172.83 (19)	N4—C14—C15—C10	−179.30 (19)
O1—C1—C2—C7	7.9 (3)	C13—C14—C15—C10	1.9 (3)


Symmetry codes: (i) $-x, -y, -z+2$.

Hydrogen-bond geometry (\AA , $^\circ$)

$D\cdots H$	$H\cdots A$	$D\cdots A$	$D\cdots H\cdots A$
O13—H13A ⁱⁱ ···O14 ⁱⁱ	0.80 (3)	1.84 (3)	2.641 (3)
O13—H13B ⁱⁱⁱ ···O10 ⁱⁱⁱ	0.77 (2)	2.22 (2)	2.838 (2)
O14—H14A ^{iv} ···O15 ^{iv}	0.81 (3)	1.95 (3)	2.758 (4)
O14—H14B ^{iv} ···O7	0.80 (3)	2.51 (4)	3.182 (3)
O15—H15A ^v ···O12 ^v	0.83 (4)	2.16 (4)	2.953 (4)
O15—H15B ^v ···O5 ^{vi}	0.84 (3)	2.17 (2)	2.996 (4)
O15—H15B ^v ···O6 ^{vi}	0.84 (3)	2.60 (4)	3.273 (4)
C15—H15 ^{vii} ···O5 ^{vii}	0.93	2.60	3.438 (3)

Symmetry codes: (ii) $-x+1, -y, -z+2$; (iii) $x, y, z+1$; (iv) $x-1/2, -y+1/2, z+1/2$; (v) $x+1, y, z$; (vi) $x, y, z-1$; (vii) $x-1/2, -y+1/2, z-1/2$.

Fig. 1



supplementary materials

Fig. 2




Fig. 3

