

# Benzene-1,3-diammonium bis(pyridine-2,6-dicarboxylato)nickelate(II) pentahydrate

Hoda Pasdar,<sup>a</sup> Saghi Sadat Kashani,<sup>a\*</sup> Hossein Aghabozorg<sup>a</sup> and Behrouz Notash<sup>b</sup>

<sup>a</sup>Department of Chemistry, Islamic Azad University, North Tehran Branch, Tehran, Iran, and <sup>b</sup>Department of Chemistry, Shahid Beheshti University, G. C., Evin, Tehran 1983963113, Iran

Correspondence e-mail: s.sadatkashani@gmail.com


Received 5 December 2010; accepted 26 December 2010

Key indicators: single-crystal X-ray study;  $T = 298$  K; mean  $\sigma(\text{C}-\text{C}) = 0.004$  Å; R factor = 0.054; wR factor = 0.086; data-to-parameter ratio = 16.4.

In the title compound,  $(\text{C}_6\text{H}_{10}\text{N}_2)[\text{Ni}(\text{C}_7\text{H}_3\text{NO}_4)_2] \cdot 5\text{H}_2\text{O}$ , the  $\text{Ni}^{\text{II}}$  ion is six-coordinated by two N and four O atoms from two pyridine-2,6-dicarboxylate ligands in a distorted octahedral fashion. The crystal packing is stabilized by intermolecular  $\text{O}-\text{H} \cdots \text{O}$  and  $\text{N}-\text{H} \cdots \text{O}$  and weak  $\text{C}-\text{H} \cdots \text{O}$  hydrogen bonds and  $\pi-\pi$  interactions [centroid-centroid distances = 3.4669 (19) and 3.764 (2) Å].

## Related literature

For background to proton-transfer compounds, see: Aghabozorg *et al.* (2008). For related structures, see: Aghabozorg *et al.* (2009); Beatty *et al.* (2002); Dobrzycki & Woźniak (2008); Attar Gharamaleki *et al.* (2009); Imaz *et al.* (2007); MacDonald *et al.* (2000, 2004); Sharif *et al.* (2007).



## Experimental

### Crystal data

$(\text{C}_6\text{H}_{10}\text{N}_2)[\text{Ni}(\text{C}_7\text{H}_3\text{NO}_4)_2] \cdot 5\text{H}_2\text{O}$   
 $M_r = 589.14$   
 Monoclinic,  $P2_1/n$

$a = 7.5331$  (15) Å  
 $b = 18.085$  (4) Å  
 $c = 18.578$  (4) Å

$\beta = 100.90$  (3)°  
 $V = 2485.3$  (9) Å<sup>3</sup>  
 $Z = 4$   
 Mo  $K\alpha$  radiation

$\mu = 0.86$  mm<sup>-1</sup>  
 $T = 298$  K  
 $0.20 \times 0.15 \times 0.10$  mm

### Data collection

Stoe IPDS II diffractometer  
 Absorption correction: numerical  
 ( $X\text{-RED}$  and  $X\text{-SHAPE}$ ; Stoe & Cie, 2005)  
 $T_{\text{min}} = 0.855$ ,  $T_{\text{max}} = 0.920$

18439 measured reflections  
 6687 independent reflections  
 3608 reflections with  $I > 2\sigma(I)$   
 $R_{\text{int}} = 0.090$

### Refinement

$R[F^2 > 2\sigma(F^2)] = 0.054$   
 $wR(F^2) = 0.086$   
 $S = 0.92$   
 6687 reflections  
 407 parameters  
 4 restraints

H atoms treated by a mixture of independent and constrained refinement  
 $\Delta\rho_{\text{max}} = 0.39$  e Å<sup>-3</sup>  
 $\Delta\rho_{\text{min}} = -0.36$  e Å<sup>-3</sup>

**Table 1**

Hydrogen-bond geometry (Å, °).

$D-\text{H} \cdots A$	$D-\text{H}$	$\text{H} \cdots A$	$D \cdots A$	$D-\text{H} \cdots A$
$\text{C18}-\text{H18} \cdots \text{O2}^{\text{i}}$	0.93	2.48	3.118 (4)	126
$\text{C3}-\text{H3} \cdots \text{O7}^{\text{ii}}$	0.93	2.56	3.284 (4)	135
$\text{O13}-\text{H13B} \cdots \text{O9}^{\text{iii}}$	0.82 (5)	2.14 (5)	2.945 (6)	166 (5)
$\text{O13}-\text{H13A} \cdots \text{O6}^{\text{iv}}$	0.76 (4)	2.05 (4)	2.776 (5)	158 (6)
$\text{O12}-\text{H12B} \cdots \text{O13}^{\text{v}}$	0.79 (4)	2.02 (4)	2.808 (5)	176 (8)
$\text{O12}-\text{H12A} \cdots \text{O3}^{\text{vi}}$	0.89 (6)	1.95 (6)	2.838 (4)	174 (5)
$\text{O11}-\text{H11B} \cdots \text{O9}$	0.75 (3)	2.08 (3)	2.829 (4)	173 (5)
$\text{O11}-\text{H11A} \cdots \text{O7}$	0.79 (4)	2.03 (4)	2.798 (4)	163 (4)
$\text{O10}-\text{H10B} \cdots \text{O2}$	0.97 (5)	1.80 (5)	2.750 (4)	168 (5)
$\text{O10}-\text{H10A} \cdots \text{O12}^{\text{vii}}$	0.79 (3)	2.20 (5)	2.880 (5)	146 (6)
$\text{O9}-\text{H9B} \cdots \text{O4}^{\text{ii}}$	0.85 (5)	2.07 (5)	2.918 (4)	172 (5)
$\text{O9}-\text{H9A} \cdots \text{O10}$	0.86 (5)	1.91 (5)	2.772 (5)	176 (4)
$\text{N4}-\text{H4C} \cdots \text{O6}^{\text{viii}}$	0.97 (4)	1.78 (4)	2.730 (4)	167 (3)
$\text{N4}-\text{H4B} \cdots \text{O4}^{\text{vi}}$	0.82 (3)	2.05 (4)	2.854 (4)	168 (3)
$\text{N4}-\text{H4A} \cdots \text{O11}$	0.91 (4)	1.94 (4)	2.842 (4)	173 (3)
$\text{N3}-\text{H3C} \cdots \text{O1}$	0.91 (5)	1.82 (5)	2.702 (4)	164 (4)
$\text{N3}-\text{H3B} \cdots \text{O8}^{\text{vii}}$	0.93 (5)	1.85 (5)	2.773 (4)	173 (4)
$\text{N3}-\text{H3A} \cdots \text{O12}^{\text{vii}}$	0.91 (4)	2.07 (4)	2.899 (5)	152 (4)

Symmetry codes: (i)  $x - \frac{1}{2}, -y + \frac{3}{2}, z - \frac{1}{2}$ ; (ii)  $-x + 1, -y + 2, -z + 2$ ; (iii)  $x + \frac{1}{2}, -y + \frac{3}{2}, z - \frac{1}{2}$ ; (iv)  $-x + 2, -y + 2, -z + 1$ ; (v)  $x - \frac{1}{2}, -y + \frac{3}{2}, z + \frac{1}{2}$ ; (vi)  $-x + \frac{1}{2}, y - \frac{1}{2}, -z + \frac{3}{2}$ ; (vii)  $x + 1, y, z$ ; (viii)  $-x + \frac{3}{2}, y - \frac{1}{2}, -z + \frac{3}{2}$ .

Data collection:  $X\text{-AREA}$  (Stoe & Cie, 2005); cell refinement:  $X\text{-AREA}$ ; data reduction:  $X\text{-AREA}$ ; program(s) used to solve structure:  $SHELXS97$  (Sheldrick, 2008); program(s) used to refine structure:  $SHELXL97$  (Sheldrick, 2008); molecular graphics:  $ORTEP-3$  for Windows (Farrugia, 1997); software used to prepare material for publication:  $WinGX$  (Farrugia, 1999).

The authors are grateful to the Islamic Azad University, North Branch, for financial support of this work.

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: BT5430).

## References

- Aghabozorg, H., Manteghi, F. & Sheshmani, S. (2008). *J. Iran. Chem. Soc.* **5**, 184–227.  
 Aghabozorg, H., Sadr-khanlou, E., Shokrollahi, A., Ghaedi, M. & Shamsipur, M. (2009). *J. Iran. Chem. Soc.* **6**, 55–70.  
 Attar Gharamaleki, J., Aghabozorg, H., Derikvand, Z. & Yousefi, M. (2009). *Acta Cryst.* **E65**, m824–m825.

- Beatty, A. M., Granger, K. E. & Simpson, A. E. (2002). *Chem. Eur. J.* **8**, 3254–3259.
- Dobrzycki, L. & Woźniak, K. (2008). *CrystEngComm*, **10**, 577–589.
- Farrugia, L. J. (1997). *J. Appl. Cryst.* **30**, 565.
- Farrugia, L. J. (1999). *J. Appl. Cryst.* **32**, 837–838.
- Imaz, I., Thillet, A. & Sutter, J. P. (2007). *Cryst. Growth Des.* **7**, 1753–1761.
- MacDonald, J. C., Dorrestein, P., Pilley, M. M., Foote, M. M., Lundburg, J. L., Henning, R. W., Schultz, A. J. & Manson, J. L. (2000). *J. Am. Chem. Soc.* **122**, 11692–11702.
- MacDonald, J. C., Luo, T.-J. M. & Palmore, G. T. R. (2004). *Cryst. Growth Des.* **4**, 1203–1209.
- Sharif, M. A., Aghajani, Z. & Aghabozorg, H. (2007). *Anal. Sci.* **23**, x147–x148.
- Sheldrick, G. M. (2008). *Acta Cryst.* **A64**, 112–122.
- Stoe & Cie (2005). *X-AREA, X-RED and X-SHAPE*. Stoe & Cie, Darmstadt, Germany.

**supplementary materials**

*Acta Cryst.* (2011). E67, m193-m194 [ doi:10.1107/S1600536810054371 ]

## Benzene-1,3-diammonium bis(pyridine-2,6-dicarboxylato)nickelate(II) pentahydrate

H. Pasdar, S. Sadat Kashani, H. Aghabozorg and B. Notash

### Comment

Pyridine-2,6-dicarboxylic acid (pydcH<sub>2</sub>) was commonly used as an acid in proton transfer systems (Aghabozorg *et al.* 2008). It has been reported that several proton transfer systems containing the anionic [Ni(pydc)<sub>2</sub>]<sup>2-</sup> moiety and different cationic parts (Aghabozorg *et al.* 2009; Attar Gharamaleki *et al.* 2009; MacDonald *et al.* 2000; MacDonald *et al.* 2004; Sharif *et al.* 2007). In addition, the formation of mono (Beatty *et al.* 2002) and diprotonated benzene-1,3-diamine (Dobrzycki & Woźniak 2008; Imaz *et al.* 2007) have been observed previously.

In the title compound, (bdaH<sub>2</sub>)[Ni(pydc)<sub>2</sub>].5H<sub>2</sub>O, the anionic part is comprised of a Ni<sup>II</sup> ion which is six-coordinated by two pyridyl nitrogen and four oxygen atoms from pydc ligands. The Ni(II) ion has a distorted octahedral geometry. (Fig. 1). Bond lengths for Ni—O and Ni—N and angles (Table 1) are in normal ranges (Aghabozorg *et al.* 2009; Attar Gharamaleki *et al.* 2009; Sharif *et al.* 2007). Crystal packing is stabilized by intermolecular O—H...O, N—H...O and weak C—H...O intermolecular hydrogen bonds which formed between [Ni(pydc)<sub>2</sub>]<sup>2-</sup>, (bdaH<sub>2</sub>)<sup>2+</sup> and water molecules (Fig. 2 & Table 2). There are also  $\pi$ - $\pi$  interactions between pyridine rings of pydc and between benzene ring of (bdaH<sub>2</sub>)<sup>2+</sup> and pyridine ring of pydc molecule by distances Cg5...Cg5<sup>ix</sup> and Cg6...Cg7 of 3.4669 (19) and 3.764 (2) Å, respectively. [Cg5, Cg6 and Cg7 are centroids of N1/C1—C6, N2/C8—C12 and C15—C20 rings, respectively. Symmetry code: (ix) 1 - x, 2 - y, 2 - z]. Intermolecular  $\pi$ - $\pi$  interactions are shown in Fig. 3.

### Experimental

A solution of pyridine-2,6-dicarboxylic acid (pydcH<sub>2</sub>) (162 mg, 0.9 mmol) in 15 ml water was added to a solution of benzene-1,3-diamine (bda) (108 mg, 0.6 mmol) in 12 ml water and stirred for half an hour. Then a solution of NiCl<sub>2</sub>.6H<sub>2</sub>O (7 mg, 0.7 mmol) in 5 ml water was added to the solution of pydcH<sub>2</sub> and bda. The resulted solution was stirred for 2 hrs and green crystals of the title compound were obtained after one week which were suitable for X-ray analysis (m.p 200°C).

### Refinement

The hydrogen atoms of the water molecules and of the diammonium groups were found in a difference Fourier map and refined isotropically. The O—H bonds to H10A, H11B, H12B and H13A were refined with a distance restraint of 0.82 (4) Å. The C—H protons were positioned geometrically and refined as riding atoms with C—H = 0.93 Å and  $U_{iso}(H) = 1.2U_{eq}(C)$ .

## Figures




Fig. 1. The molecular structure of the title compound with displacement ellipsoids drawn at 30% probability level.




Fig. 2. Packing diagram of the title compound viewed down the *a*-axis. The intermolecular O—H...O, N—H...O and C—H...O hydrogen bonds are shown as blue dashed lines.




Fig. 3. Packing diagram of the title compound viewed down the *a*-axis showing intermolecular  $\pi$ - $\pi$  interactions (dashed lines). Hydrogen atoms and water molecules have been omitted for clarity.

## Benzene-1,3-diammonium bis(pyridine-2,6-dicarboxylato)nickelate(II) pentahydrate

### Crystal data

(C<sub>6</sub>H<sub>10</sub>N<sub>2</sub>)[Ni(C<sub>7</sub>H<sub>3</sub>NO<sub>4</sub>)<sub>2</sub>]·5H<sub>2</sub>O

*M<sub>r</sub>* = 589.14

Monoclinic, *P*2<sub>1</sub>/*n*

Hall symbol: -*P* 2yn

*a* = 7.5331 (15) Å

*b* = 18.085 (4) Å

*c* = 18.578 (4) Å

$\beta$  = 100.90 (3)°

*V* = 2485.3 (9) Å<sup>3</sup>

*Z* = 4

*F*(000) = 1224.0

*D<sub>x</sub>* = 1.574 Mg m<sup>-3</sup>

Mo *K* $\alpha$  radiation,  $\lambda$  = 0.71073 Å

Cell parameters from 6687 reflections

$\theta$  = 2.2–29.2°

$\mu$  = 0.86 mm<sup>-1</sup>

*T* = 298 K

Needle, green

0.2 × 0.15 × 0.1 mm

*Data collection*

Stoe IPDS II diffractometer	6687 independent reflections
Radiation source: fine-focus sealed tube graphite	3608 reflections with $I > 2\sigma(I)$ $R_{\text{int}} = 0.090$
Detector resolution: 0.15 mm pixels $\text{mm}^{-1}$	$\theta_{\text{max}} = 29.2^\circ$ , $\theta_{\text{min}} = 2.2^\circ$
rotation method scans	$h = -10 \rightarrow 9$
Absorption correction: numerical [shape of crystal determined optically ( <i>X-RED</i> and <i>X-SHAPE</i> (Stoe & Cie, 2005))]	
$T_{\text{min}} = 0.855$ , $T_{\text{max}} = 0.920$	$l = -25 \rightarrow 25$
18439 measured reflections	

*Refinement*

Refinement on $F^2$	Primary atom site location: structure-invariant direct methods
Least-squares matrix: full	Secondary atom site location: difference Fourier map
$R[F^2 > 2\sigma(F^2)] = 0.054$	Hydrogen site location: inferred from neighbouring sites
$wR(F^2) = 0.086$	H atoms treated by a mixture of independent and constrained refinement
$S = 0.92$	$w = 1/[\sigma^2(F_o^2) + (0.0238P)^2]$ where $P = (F_o^2 + 2F_c^2)/3$
6687 reflections	$(\Delta/\sigma)_{\text{max}} = 0.001$
407 parameters	$\Delta\rho_{\text{max}} = 0.39 \text{ e } \text{\AA}^{-3}$
4 restraints	$\Delta\rho_{\text{min}} = -0.36 \text{ e } \text{\AA}^{-3}$

*Special details*

**Geometry.** All e.s.d.'s (except the e.s.d. in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell e.s.d.'s are taken into account individually in the estimation of e.s.d.'s in distances, angles and torsion angles; correlations between e.s.d.'s in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell e.s.d.'s is used for estimating e.s.d.'s involving l.s. planes.

**Refinement.** Refinement of  $F^2$  against ALL reflections. The weighted  $R$ -factor  $wR$  and goodness of fit  $S$  are based on  $F^2$ , conventional  $R$ -factors  $R$  are based on  $F$ , with  $F$  set to zero for negative  $F^2$ . The threshold expression of  $F^2 > \sigma(F^2)$  is used only for calculating  $R$ -factors(gt) etc. and is not relevant to the choice of reflections for refinement.  $R$ -factors based on  $F^2$  are statistically about twice as large as those based on  $F$ , and  $R$ -factors based on ALL data will be even larger.

*Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters ( $\text{\AA}^2$ )*

	<i>x</i>	<i>y</i>	<i>z</i>	$U_{\text{iso}}^*/U_{\text{eq}}$
O12	0.0352 (4)	0.65115 (16)	0.82305 (17)	0.0522 (8)
Ni1	0.64385 (6)	0.98344 (2)	0.81233 (2)	0.02316 (10)
N1	0.6858 (3)	1.00256 (12)	0.91843 (12)	0.0203 (5)
N2	0.6146 (3)	0.96578 (13)	0.70636 (12)	0.0225 (6)

## supplementary materials

---

O7	0.4041 (3)	0.91734 (12)	0.79023 (11)	0.0312 (5)
O3	0.4988 (3)	1.08346 (11)	0.81671 (11)	0.0305 (5)
O5	0.8722 (3)	1.03845 (12)	0.78742 (11)	0.0368 (6)
C11	0.7284 (5)	0.97699 (19)	0.59762 (16)	0.0354 (8)
H11	0.8166	0.9944	0.5730	0.042*
C12	0.7401 (4)	0.99113 (17)	0.67087 (15)	0.0262 (7)
O1	0.8002 (3)	0.89096 (11)	0.85564 (11)	0.0315 (5)
C1	0.7845 (4)	0.95527 (15)	0.96399 (15)	0.0200 (6)
C5	0.6176 (4)	1.06446 (15)	0.94246 (15)	0.0227 (7)
N3	0.9677 (4)	0.80203 (19)	0.77250 (17)	0.0290 (6)
C3	0.7580 (4)	1.03320 (17)	1.06483 (16)	0.0277 (7)
H3	0.7827	1.0437	1.1147	0.033*
C4	0.6531 (4)	1.08166 (17)	1.01596 (16)	0.0275 (7)
H4	0.6078	1.1249	1.0328	0.033*
C2	0.8252 (4)	0.96944 (16)	1.03883 (15)	0.0272 (7)
H2	0.8960	0.9368	1.0707	0.033*
C8	0.4727 (4)	0.92723 (16)	0.67186 (16)	0.0255 (7)
C15	0.8119 (4)	0.77521 (16)	0.71966 (16)	0.0241 (7)
C7	0.5013 (4)	1.10803 (16)	0.88110 (17)	0.0256 (7)
C20	0.6666 (4)	0.74641 (17)	0.74619 (17)	0.0261 (7)
H20	0.6665	0.7454	0.7962	0.031*
C16	0.8142 (4)	0.77784 (17)	0.64577 (17)	0.0289 (7)
H16	0.9119	0.7982	0.6287	0.035*
O4	0.4185 (3)	1.16332 (11)	0.89702 (12)	0.0322 (5)
O2	0.9292 (4)	0.83930 (13)	0.96180 (12)	0.0431 (6)
O8	0.2157 (3)	0.86339 (14)	0.69685 (13)	0.0467 (7)
O6	1.0038 (3)	1.06724 (13)	0.69308 (13)	0.0468 (7)
C19	0.5220 (4)	0.71920 (16)	0.69634 (16)	0.0238 (7)
N4	0.3701 (4)	0.68770 (17)	0.72430 (17)	0.0275 (6)
C18	0.5215 (4)	0.72028 (17)	0.62201 (17)	0.0303 (8)
H18	0.4235	0.7015	0.5890	0.036*
C6	0.8438 (4)	0.88880 (17)	0.92559 (16)	0.0266 (7)
C14	0.8847 (4)	1.03584 (17)	0.72077 (17)	0.0311 (8)
C17	0.6668 (5)	0.74933 (19)	0.59727 (17)	0.0369 (8)
H17	0.6669	0.7500	0.5472	0.044*
C9	0.4516 (5)	0.91218 (18)	0.59783 (17)	0.0334 (8)
H9	0.3513	0.8863	0.5734	0.040*
C10	0.5841 (5)	0.93673 (19)	0.56106 (18)	0.0394 (9)
H10	0.5756	0.9260	0.5116	0.047*
C13	0.3517 (4)	0.90100 (17)	0.72284 (16)	0.0271 (7)
O11	0.3389 (4)	0.77842 (17)	0.84618 (15)	0.0402 (7)
O9	0.5261 (5)	0.72860 (19)	0.98420 (18)	0.0583 (9)
O10	0.8429 (5)	0.6926 (2)	0.93695 (19)	0.0649 (9)
O13	0.8277 (5)	0.9051 (2)	0.4247 (2)	0.0646 (9)
H11A	0.360 (6)	0.820 (2)	0.839 (2)	0.057 (15)*
H11B	0.381 (6)	0.765 (2)	0.8838 (19)	0.062 (15)*
H12B	0.114 (8)	0.633 (4)	0.852 (3)	0.15 (3)*
H13A	0.879 (7)	0.923 (3)	0.398 (3)	0.11 (3)*
H12A	0.025 (7)	0.627 (3)	0.781 (3)	0.103 (19)*

H9B	0.552 (7)	0.761 (3)	1.018 (3)	0.09 (2)*
H9A	0.628 (7)	0.718 (2)	0.972 (2)	0.067 (17)*
H10A	0.857 (8)	0.670 (3)	0.902 (2)	0.10 (2)*
H13B	0.894 (7)	0.873 (3)	0.446 (3)	0.08 (2)*
H10B	0.882 (7)	0.744 (3)	0.940 (3)	0.095 (18)*
H3A	1.023 (6)	0.763 (2)	0.797 (2)	0.063 (14)*
H3B	1.051 (6)	0.826 (2)	0.750 (2)	0.056 (13)*
H3C	0.932 (6)	0.834 (2)	0.805 (2)	0.061 (13)*
H4C	0.398 (5)	0.641 (2)	0.750 (2)	0.054 (12)*
H4A	0.351 (5)	0.716 (2)	0.763 (2)	0.045 (11)*
H4B	0.279 (5)	0.6851 (18)	0.6925 (19)	0.034 (10)*

*Atomic displacement parameters ( $\text{\AA}^2$ )*

	$U^{11}$	$U^{22}$	$U^{33}$	$U^{12}$	$U^{13}$	$U^{23}$
O12	0.064 (2)	0.0460 (17)	0.0440 (17)	0.0030 (15)	0.0021 (16)	-0.0136 (14)
Ni1	0.0261 (2)	0.02501 (18)	0.01821 (16)	-0.0019 (2)	0.00388 (14)	-0.00170 (18)
N1	0.0188 (12)	0.0206 (14)	0.0222 (12)	-0.0019 (9)	0.0056 (10)	-0.0015 (9)
N2	0.0244 (13)	0.0238 (14)	0.0202 (12)	-0.0022 (10)	0.0065 (10)	-0.0012 (10)
O7	0.0347 (13)	0.0370 (13)	0.0238 (11)	-0.0078 (10)	0.0100 (10)	-0.0045 (9)
O3	0.0346 (13)	0.0301 (12)	0.0247 (11)	0.0043 (10)	0.0004 (10)	0.0004 (9)
O5	0.0391 (14)	0.0434 (14)	0.0279 (11)	-0.0156 (11)	0.0059 (10)	-0.0076 (10)
C11	0.047 (2)	0.0370 (18)	0.0253 (15)	-0.0073 (18)	0.0145 (14)	-0.0003 (15)
C12	0.0315 (16)	0.0238 (16)	0.0247 (14)	-0.0044 (14)	0.0087 (12)	0.0014 (13)
O1	0.0398 (14)	0.0310 (12)	0.0236 (11)	0.0110 (10)	0.0055 (10)	-0.0036 (9)
C1	0.0137 (14)	0.0242 (15)	0.0232 (14)	-0.0027 (12)	0.0065 (12)	0.0009 (12)
C5	0.0205 (15)	0.0215 (15)	0.0259 (15)	-0.0013 (12)	0.0037 (13)	0.0000 (12)
N3	0.0201 (15)	0.0389 (18)	0.0281 (15)	-0.0029 (14)	0.0049 (13)	-0.0087 (14)
C3	0.0271 (16)	0.0371 (19)	0.0194 (14)	-0.0066 (14)	0.0055 (13)	-0.0057 (12)
C4	0.0247 (16)	0.0303 (17)	0.0298 (16)	-0.0002 (13)	0.0108 (13)	-0.0110 (13)
C2	0.0287 (16)	0.0288 (18)	0.0222 (14)	-0.0009 (13)	0.0003 (13)	0.0027 (12)
C8	0.0308 (18)	0.0221 (15)	0.0228 (15)	-0.0016 (13)	0.0033 (13)	0.0001 (12)
C15	0.0191 (15)	0.0245 (16)	0.0275 (16)	0.0021 (13)	0.0013 (13)	-0.0072 (13)
C7	0.0231 (16)	0.0222 (16)	0.0300 (17)	-0.0034 (13)	0.0012 (13)	-0.0005 (13)
C20	0.0230 (16)	0.0331 (17)	0.0223 (15)	-0.0011 (14)	0.0045 (13)	-0.0028 (13)
C16	0.0272 (17)	0.0323 (18)	0.0291 (17)	-0.0044 (14)	0.0102 (14)	-0.0026 (14)
O4	0.0273 (12)	0.0288 (12)	0.0397 (13)	0.0071 (10)	0.0046 (10)	-0.0025 (10)
O2	0.0580 (17)	0.0336 (13)	0.0354 (13)	0.0184 (12)	0.0028 (12)	0.0041 (10)
O8	0.0359 (15)	0.0651 (17)	0.0383 (14)	-0.0234 (13)	0.0050 (12)	-0.0055 (12)
O6	0.0472 (16)	0.0525 (16)	0.0451 (15)	-0.0244 (13)	0.0199 (12)	-0.0063 (12)
C19	0.0195 (15)	0.0224 (15)	0.0298 (16)	0.0021 (12)	0.0055 (13)	-0.0021 (13)
N4	0.0202 (15)	0.0293 (16)	0.0323 (16)	-0.0033 (12)	0.0029 (13)	-0.0012 (13)
C18	0.0294 (18)	0.0313 (18)	0.0282 (16)	-0.0047 (14)	0.0001 (14)	-0.0062 (13)
C6	0.0266 (17)	0.0269 (17)	0.0266 (16)	-0.0008 (14)	0.0056 (14)	-0.0010 (13)
C14	0.0299 (18)	0.0325 (19)	0.0337 (17)	-0.0066 (14)	0.0132 (14)	-0.0037 (14)
C17	0.042 (2)	0.047 (2)	0.0215 (16)	-0.0046 (17)	0.0054 (16)	-0.0021 (15)
C9	0.036 (2)	0.0371 (19)	0.0248 (16)	-0.0068 (15)	0.0012 (15)	-0.0032 (14)
C10	0.054 (2)	0.044 (2)	0.0226 (17)	-0.0075 (18)	0.0128 (16)	-0.0024 (15)



## supplementary materials

---

C13	0.0239 (17)	0.0287 (17)	0.0299 (17)	-0.0004 (14)	0.0082 (14)	0.0003 (13)
O11	0.0474 (17)	0.0362 (16)	0.0357 (16)	-0.0017 (13)	0.0051 (13)	0.0008 (13)
O9	0.059 (2)	0.068 (2)	0.0464 (19)	0.0053 (18)	0.0066 (17)	-0.0100 (16)
O10	0.077 (2)	0.058 (2)	0.063 (2)	0.0039 (18)	0.0247 (18)	-0.0084 (18)
O13	0.060 (2)	0.082 (3)	0.059 (2)	-0.001 (2)	0.0275 (19)	0.0045 (19)

### *Geometric parameters (Å, °)*

O12—H12B	0.79 (4)	C2—H2	0.9300
O12—H12A	0.89 (6)	C8—C9	1.381 (4)
Ni1—N2	1.965 (2)	C8—C13	1.510 (5)
Ni1—N1	1.967 (2)	C15—C16	1.377 (4)
Ni1—O5	2.113 (2)	C15—C20	1.384 (4)
Ni1—O1	2.115 (2)	C7—O4	1.244 (4)
Ni1—O3	2.123 (2)	C20—C19	1.380 (4)
Ni1—O7	2.140 (2)	C20—H20	0.9300
N1—C1	1.328 (3)	C16—C17	1.390 (4)
N1—C5	1.343 (4)	C16—H16	0.9300
N2—C12	1.332 (4)	O2—C6	1.226 (3)
N2—C8	1.334 (4)	O8—C13	1.248 (4)
O7—C13	1.274 (3)	O6—C14	1.251 (4)
O3—C7	1.273 (4)	C19—C18	1.381 (4)
O5—C14	1.260 (4)	C19—N4	1.459 (4)
C11—C12	1.371 (4)	N4—H4C	0.97 (4)
C11—C10	1.376 (5)	N4—H4A	0.91 (4)
C11—H11	0.9300	N4—H4B	0.82 (3)
C12—C14	1.522 (4)	C18—C17	1.370 (5)
O1—C6	1.279 (3)	C18—H18	0.9300
C1—C2	1.390 (4)	C17—H17	0.9300
C1—C6	1.508 (4)	C9—C10	1.385 (5)
C5—C4	1.376 (4)	C9—H9	0.9300
C5—C7	1.520 (4)	C10—H10	0.9300
N3—C15	1.463 (4)	O11—H11A	0.79 (4)
N3—H3A	0.91 (4)	O11—H11B	0.75 (3)
N3—H3B	0.93 (5)	O9—H9B	0.85 (5)
N3—H3C	0.91 (5)	O9—H9A	0.86 (5)
C3—C2	1.383 (4)	O10—H10A	0.79 (3)
C3—C4	1.395 (4)	O10—H10B	0.97 (5)
C3—H3	0.9300	O13—H13A	0.76 (4)
C4—H4	0.9300	O13—H13B	0.82 (5)
H12B—O12—H12A	109 (6)	C3—C2—H2	120.7
N2—Ni1—N1	177.13 (11)	C1—C2—H2	120.7
N2—Ni1—O5	78.39 (10)	N2—C8—C9	120.8 (3)
N1—Ni1—O5	98.84 (9)	N2—C8—C13	112.7 (3)
N2—Ni1—O1	101.62 (9)	C9—C8—C13	126.4 (3)
N1—Ni1—O1	77.61 (9)	C16—C15—C20	121.9 (3)
O5—Ni1—O1	92.17 (9)	C16—C15—N3	119.8 (3)
N2—Ni1—O3	102.47 (9)	C20—C15—N3	118.3 (3)
N1—Ni1—O3	78.34 (9)	O4—C7—O3	125.8 (3)

O5—Ni1—O3	93.06 (9)	O4—C7—C5	118.9 (3)
O1—Ni1—O3	155.91 (8)	O3—C7—C5	115.3 (3)
N2—Ni1—O7	77.66 (10)	C19—C20—C15	118.2 (3)
N1—Ni1—O7	105.08 (9)	C19—C20—H20	120.9
O5—Ni1—O7	156.01 (8)	C15—C20—H20	120.9
O1—Ni1—O7	91.16 (9)	C15—C16—C17	118.3 (3)
O3—Ni1—O7	93.54 (9)	C15—C16—H16	120.8
C1—N1—C5	121.9 (2)	C17—C16—H16	120.8
C1—N1—Ni1	119.34 (19)	C20—C19—C18	121.2 (3)
C5—N1—Ni1	118.74 (18)	C20—C19—N4	118.2 (3)
C12—N2—C8	121.4 (3)	C18—C19—N4	120.5 (3)
C12—N2—Ni1	118.97 (19)	C19—N4—H4C	114 (2)
C8—N2—Ni1	119.6 (2)	C19—N4—H4A	108 (2)
C13—O7—Ni1	114.3 (2)	H4C—N4—H4A	99 (3)
C7—O3—Ni1	114.67 (18)	C19—N4—H4B	112 (3)
C14—O5—Ni1	114.38 (19)	H4C—N4—H4B	112 (3)
C12—C11—C10	119.0 (3)	H4A—N4—H4B	111 (3)
C12—C11—H11	120.5	C17—C18—C19	119.3 (3)
C10—C11—H11	120.5	C17—C18—H18	120.3
N2—C12—C11	120.6 (3)	C19—C18—H18	120.3
N2—C12—C14	112.0 (2)	O2—C6—O1	125.9 (3)
C11—C12—C14	127.4 (3)	O2—C6—C1	119.6 (3)
C6—O1—Ni1	115.35 (19)	O1—C6—C1	114.5 (3)
N1—C1—C2	120.7 (3)	O6—C14—O5	125.4 (3)
N1—C1—C6	113.1 (2)	O6—C14—C12	118.5 (3)
C2—C1—C6	126.2 (3)	O5—C14—C12	116.1 (3)
N1—C5—C4	120.3 (3)	C18—C17—C16	121.1 (3)
N1—C5—C7	112.7 (2)	C18—C17—H17	119.5
C4—C5—C7	127.1 (3)	C16—C17—H17	119.5
C15—N3—H3A	108 (3)	C8—C9—C10	118.0 (3)
C15—N3—H3B	112 (2)	C8—C9—H9	121.0
H3A—N3—H3B	108 (4)	C10—C9—H9	121.0
C15—N3—H3C	111 (3)	C11—C10—C9	120.1 (3)
H3A—N3—H3C	109 (4)	C11—C10—H10	119.9
H3B—N3—H3C	109 (4)	C9—C10—H10	119.9
C2—C3—C4	119.8 (3)	O8—C13—O7	125.8 (3)
C2—C3—H3	120.1	O8—C13—C8	118.5 (3)
C4—C3—H3	120.1	O7—C13—C8	115.6 (3)
C5—C4—C3	118.9 (3)	H11A—O11—H11B	113 (4)
C5—C4—H4	120.6	H9B—O9—H9A	105 (5)
C3—C4—H4	120.6	H10A—O10—H10B	117 (6)
C3—C2—C1	118.5 (3)	H13A—O13—H13B	106 (6)
N2—Ni1—N1—C1	-74 (2)	Ni1—N1—C5—C4	-176.3 (2)
O5—Ni1—N1—C1	-89.1 (2)	C1—N1—C5—C7	-177.0 (3)
O1—Ni1—N1—C1	1.2 (2)	Ni1—N1—C5—C7	4.9 (3)
O3—Ni1—N1—C1	179.6 (2)	N1—C5—C4—C3	-1.0 (5)
O7—Ni1—N1—C1	89.0 (2)	C7—C5—C4—C3	177.5 (3)
N2—Ni1—N1—C5	104 (2)	C2—C3—C4—C5	0.4 (5)
O5—Ni1—N1—C5	89.0 (2)	C4—C3—C2—C1	-0.4 (5)

## supplementary materials

O1—Ni1—N1—C5	179.3 (2)	N1—C1—C2—C3	1.1 (4)
O3—Ni1—N1—C5	-2.3 (2)	C6—C1—C2—C3	-179.5 (3)
O7—Ni1—N1—C5	-92.9 (2)	C12—N2—C8—C9	0.2 (4)
N1—Ni1—N2—C12	-14 (2)	Ni1—N2—C8—C9	-178.1 (2)
O5—Ni1—N2—C12	2.1 (2)	C12—N2—C8—C13	177.5 (3)
O1—Ni1—N2—C12	-87.8 (2)	Ni1—N2—C8—C13	-0.8 (3)
O3—Ni1—N2—C12	92.7 (2)	Ni1—O3—C7—O4	-176.1 (3)
O7—Ni1—N2—C12	-176.4 (2)	Ni1—O3—C7—C5	4.0 (3)
N1—Ni1—N2—C8	164.8 (19)	N1—C5—C7—O4	174.2 (3)
O5—Ni1—N2—C8	-179.6 (2)	C4—C5—C7—O4	-4.4 (5)
O1—Ni1—N2—C8	90.5 (2)	N1—C5—C7—O3	-5.9 (4)
O3—Ni1—N2—C8	-89.0 (2)	C4—C5—C7—O3	175.5 (3)
O7—Ni1—N2—C8	1.9 (2)	C16—C15—C20—C19	0.6 (5)
N2—Ni1—O7—C13	-2.9 (2)	N3—C15—C20—C19	-178.2 (3)
N1—Ni1—O7—C13	178.0 (2)	C20—C15—C16—C17	-1.2 (5)
O5—Ni1—O7—C13	-6.6 (3)	N3—C15—C16—C17	177.6 (3)
O1—Ni1—O7—C13	-104.5 (2)	C15—C20—C19—C18	0.2 (4)
O3—Ni1—O7—C13	99.1 (2)	C15—C20—C19—N4	179.0 (3)
N2—Ni1—O3—C7	-178.4 (2)	C20—C19—C18—C17	-0.4 (5)
N1—Ni1—O3—C7	-1.2 (2)	N4—C19—C18—C17	-179.2 (3)
O5—Ni1—O3—C7	-99.6 (2)	Ni1—O1—C6—O2	177.8 (3)
O1—Ni1—O3—C7	2.7 (4)	Ni1—O1—C6—C1	-3.4 (3)
O7—Ni1—O3—C7	103.5 (2)	N1—C1—C6—O2	-176.8 (3)
N2—Ni1—O5—C14	0.6 (2)	C2—C1—C6—O2	3.7 (5)
N1—Ni1—O5—C14	179.8 (2)	N1—C1—C6—O1	4.4 (4)
O1—Ni1—O5—C14	102.0 (2)	C2—C1—C6—O1	-175.1 (3)
O3—Ni1—O5—C14	-101.6 (2)	Ni1—O5—C14—O6	176.6 (3)
O7—Ni1—O5—C14	4.2 (4)	Ni1—O5—C14—C12	-2.7 (3)
C8—N2—C12—C11	-1.2 (4)	N2—C12—C14—O6	-175.0 (3)
Ni1—N2—C12—C11	177.1 (2)	C11—C12—C14—O6	3.9 (5)
C8—N2—C12—C14	177.8 (3)	N2—C12—C14—O5	4.3 (4)
Ni1—N2—C12—C14	-3.9 (3)	C11—C12—C14—O5	-176.7 (3)
C10—C11—C12—N2	0.6 (5)	C19—C18—C17—C16	-0.2 (5)
C10—C11—C12—C14	-178.3 (3)	C15—C16—C17—C18	0.9 (5)
N2—Ni1—O1—C6	178.6 (2)	N2—C8—C9—C10	1.5 (5)
N1—Ni1—O1—C6	1.5 (2)	C13—C8—C9—C10	-175.4 (3)
O5—Ni1—O1—C6	100.0 (2)	C12—C11—C10—C9	1.1 (5)
O3—Ni1—O1—C6	-2.4 (4)	C8—C9—C10—C11	-2.1 (5)
O7—Ni1—O1—C6	-103.7 (2)	Ni1—O7—C13—O8	-179.7 (3)
C5—N1—C1—C2	-1.8 (4)	Ni1—O7—C13—C8	3.3 (3)
Ni1—N1—C1—C2	176.3 (2)	N2—C8—C13—O8	-179.0 (3)
C5—N1—C1—C6	178.8 (3)	C9—C8—C13—O8	-1.9 (5)
Ni1—N1—C1—C6	-3.2 (3)	N2—C8—C13—O7	-1.8 (4)
C1—N1—C5—C4	1.7 (4)	C9—C8—C13—O7	175.3 (3)

### Hydrogen-bond geometry ( $\text{\AA}$ , $^\circ$ )

$D-H\cdots A$	$D-H$	$H\cdots A$	$D\cdots A$	$D-H\cdots A$
C18—H18 $\cdots$ O2 <sup>i</sup>	0.93	2.48	3.118 (4)	126.

C3—H3···O7 <sup>ii</sup>	0.93	2.56	3.284 (4)	135.
O13—H13B···O9 <sup>iii</sup>	0.82 (5)	2.14 (5)	2.945 (6)	166 (5)
O13—H13A···O6 <sup>iv</sup>	0.76 (4)	2.05 (4)	2.776 (5)	158 (6)
O12—H12B···O13 <sup>v</sup>	0.79 (4)	2.02 (4)	2.808 (5)	176 (8)
O12—H12A···O3 <sup>vi</sup>	0.89 (6)	1.95 (6)	2.838 (4)	174 (5)
O11—H11B···O9	0.75 (3)	2.08 (3)	2.829 (4)	173 (5)
O11—H11A···O7	0.79 (4)	2.03 (4)	2.798 (4)	163 (4)
O10—H10B···O2	0.97 (5)	1.80 (5)	2.750 (4)	168 (5)
O10—H10A···O12 <sup>vii</sup>	0.79 (3)	2.20 (5)	2.880 (5)	146 (6)
O9—H9B···O4 <sup>ii</sup>	0.85 (5)	2.07 (5)	2.918 (4)	172 (5)
O9—H9A···O10	0.86 (5)	1.91 (5)	2.772 (5)	176 (4)
N4—H4C···O6 <sup>viii</sup>	0.97 (4)	1.78 (4)	2.730 (4)	167 (3)
N4—H4B···O4 <sup>vi</sup>	0.82 (3)	2.05 (4)	2.854 (4)	168 (3)
N4—H4A···O11	0.91 (4)	1.94 (4)	2.842 (4)	173 (3)
N3—H3C···O1	0.91 (5)	1.82 (5)	2.702 (4)	164 (4)
N3—H3B···O8 <sup>vii</sup>	0.93 (5)	1.85 (5)	2.773 (4)	173 (4)
N3—H3A···O12 <sup>vii</sup>	0.91 (4)	2.07 (4)	2.899 (5)	152 (4)

Symmetry codes: (i)  $x-1/2, -y+3/2, z-1/2$ ; (ii)  $-x+1, -y+2, -z+2$ ; (iii)  $x+1/2, -y+3/2, z-1/2$ ; (iv)  $-x+2, -y+2, -z+1$ ; (v)  $x-1/2, -y+3/2, z+1/2$ ; (vi)  $-x+1/2, y-1/2, -z+3/2$ ; (vii)  $x+1, y, z$ ; (viii)  $-x+3/2, y-1/2, -z+3/2$ .

Fig. 1




Fig. 2




Fig. 3

