

Bis[2-(hydroxyiminomethyl)phenolato]-nickel(II): a second monoclinic polymorph

Julia A. Rusanova,^{a*} Elena A. Buvaylo^a and Eduard B. Rusanov^b

^aNational Taras Shevchenko University, Department of Chemistry, Volodymyrska str. 64, 01033 Kyiv, Ukraine, and ^bInst. of Organic Chemistry, Acad. Sci. Ukraine, Murmanska Str. 5, Kiev 02094, Ukraine
Correspondence e-mail: rusanova_j@yahoo.com


Received 21 December 2010; accepted 10 January 2011

Key indicators: single-crystal X-ray study; $T = 296$ K; mean $\sigma(\text{C}-\text{C}) = 0.004$ Å; R factor = 0.030; wR factor = 0.072; data-to-parameter ratio = 11.1.

The title compound, $[\text{Ni}(\text{C}_7\text{H}_6\text{NO}_2)_2]$, (I), is a second monoclinic polymorph of the compound, (II), reported by Srivastava *et al.* [*Acta Cryst.* (1967), **22**, 922] and Mereiter [Private communication (2002) CCDC refcode NISALO01]. The bond lengths and angles are similar in both structures. The molecule in both structures lies on a crystallographic inversion center and both have an internal hydrogen bond. The title compound crystallizes in the space group $P2_1/c$ ($Z = 2$), whereas compound (II) is in the space group $P2_1/n$ ($Z = 2$) with a similar cell volume but different cell parameters. In both polymorphs, molecules are arranged in the layers but in contrast to the previously published compound (II) where the dihedral angle between the layers is 86.3° , in the title polymorph the same dihedral angle is 29.4° . The structure of (I) is stabilized by strong intramolecular $\text{O}-\text{H}\cdots\text{O}$ hydrogen bonding between the $\text{O}-\text{H}$ group and the phenolate O atom.

Related literature

For the original monoclinic polymorph, see: Srivastava *et al.* (1967); Mereiter (2002). For background to direct synthesis, see: Nesterov *et al.* (2004, 2006); Kovbasyuk *et al.* (1998); Vassilyeva *et al.* (1997). For bond-length data, see: Allen *et al.* (1987).



Experimental

Crystal data

$[\text{Ni}(\text{C}_7\text{H}_6\text{NO}_2)_2]$	$V = 649.42$ (5) Å ³
$M_r = 330.97$	$Z = 2$
Monoclinic, $P2_1/c$	Mo $K\alpha$ radiation
$a = 4.9912$ (2) Å	$\mu = 1.51$ mm ⁻¹
$b = 7.4717$ (3) Å	$T = 296$ K
$c = 17.4152$ (7) Å	$0.53 \times 0.36 \times 0.11$ mm
$\beta = 90.653$ (3)°	

Data collection

Bruker SMART APEXII diffractometer	4678 measured reflections
Absorption correction: numerical (SADABS; Sheldrick, 2002)	1342 independent reflections
$T_{\min} = 0.501$, $T_{\max} = 0.851$	1096 reflections with $I > 2\sigma(I)$
	$R_{\text{int}} = 0.028$

Refinement

$R[F^2 > 2\sigma(F^2)] = 0.030$	121 parameters
$wR(F^2) = 0.072$	All H-atom parameters refined
$S = 1.06$	$\Delta\rho_{\text{max}} = 0.28$ e Å ⁻³
1342 reflections	$\Delta\rho_{\text{min}} = -0.26$ e Å ⁻³

Table 1

Selected geometric parameters (Å, °).

N1—Ni1	1.8661 (19)	Ni1—O1	1.8292 (16)
O1—Ni1—N1	93.08 (8)		

Table 2

Hydrogen-bond geometry (Å, °).

$D-\text{H}\cdots A$	$D-\text{H}$	$\text{H}\cdots A$	$D\cdots A$	$D-\text{H}\cdots A$
$\text{O2}-\text{H2}\cdots\text{O1}^i$	0.80 (3)	1.80 (3)	2.511 (2)	146 (4)

Symmetry code: (i) $-x + 1, -y + 1, -z + 1$.

Data collection: APEX2 (Bruker, 2007); cell refinement: SAINT (Bruker, 2007); data reduction: SAINT; program(s) used to solve structure: SHELXS97 (Sheldrick, 2008); program(s) used to refine structure: SHELXL97 (Sheldrick, 2008); molecular graphics: SHELXTL (Sheldrick, 2008); software used to prepare material for publication: SHELXTL.

The authors gratefully acknowledge the Ukrainian State Fund for Fundamental Researchers (SFFR) for financial support of the Research Program (Chemistry).

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: BR2156).

References

- Allen, F. H., Kennard, O., Watson, D. G., Brammer, L., Orpen, A. G. & Taylor, R. (1987). *J. Chem. Soc. Perkin Trans. 2*, pp. S1–19.
Bruker (2007). APEX2 and SAINT. Bruker AXS Inc., Madison, Wisconsin, USA.
Kovbasyuk, L. A., Vassilyeva, O. Yu., Kokozay, V. N., Linert, W., Reedijk, J., Skelton, B. W. & Oliver, A. G. (1998). *J. Chem. Soc. Dalton Trans.* pp. 2735–2738.

- Mereiter, K. (2002). Private communication (refcode NISALO01). CCDC, Cambridge, England.
- Nesterov, D. S., Kokozay, V. N., Dyakonenko, V. V., Shishkin, O. V., Jezierska, J., Ozarowski, F., Kirillov, A. M., Kopylovich, M. N. & Pombeiro, A. J. L. (2006). *Chem. Commun.* pp. 4605–4607.
- Nesterov, D. S., Makhankova, V. G., Vassilyeva, O. Yu., Kokozay, V. N., Kovbasyuk, L. A., Skelton, B. W. & Jezierska, J. (2004). *Inorg. Chem.* **43**, 7868–7876.
- Sheldrick, G. M. (2002). *SADABS*. University of Göttingen, Germany.
- Sheldrick, G. M. (2008). *Acta Cryst.* **A64**, 112–122.
- Srivastava, R. C., Lingafelter, E. C. & Jain, P. C. (1967). *Acta Cryst.* **22**, 922–923.
- Vassilyeva, O. Yu., Kokozay, V. N., Zhukova, N. I. & Kovbasyuk, L. A. (1997). *Polyhedron*, **16**, 263–266.

supplementary materials

Acta Cryst. (2011). E67, m222-m223 [doi:10.1107/S1600536811001383]

Bis[2-(hydroxyiminomethyl)phenolato]nickel(II): a second monoclinic polymorph

J. A. Rusanova, E. A. Buvaylo and E. B. Rusanov

Comment

The cell dimensions of the title modification after transformation from $P2_1/c$ to $P2_1/n$ setting, are: $a=18.062$, $b=7.472$, $c=4.991$, $\beta=105.39$, whereas the cell dimensions of the reported compound (II) monoclinic $P2_1/n$ modification are: $a=13.830$, $b=4.880$, $c=10.200$ Å; $\beta=110.43^\circ$ (Srivastava *et al.*, 1967). The asymmetric unit of the title compound contains half molecule (Fig. 1), lying across a crystallographic inversion centre. The bond lengths and angles are within normal ranges (Allen *et al.*, 1987) and are comparable with the structure of the compound (II).

Experimental

The title compound was prepared by direct synthesis: manganese powder (0.06 g, 1 mmol), $\text{Ni}(\text{OAc})_2$ (0.25 g, 1 mmol), salicylic aldehyde (0.21 ml, 2 mmol), $\text{NH}_2\text{OH}\cdot\text{HCl}$ (0.14 g, 2 mmol), dimethylformamide (20 ml) were heated to 323–333 K and stirred magnetically for 40 min, until total dissolution of the manganese powder was observed. The transparent brown solution was allowed to stand at room temperature and brown-green crystals of the title compound suitable for X-ray analysis precipitated within few days. They were collected by filter-suction, washed with dry Pr^iOH and finally dried in *vacuo* at room temperature (yield; 0.12 g)

Refinement

The hydrogen atoms were located in difference Fourier synthesis and refined in isotropic approximation.

Figures




Fig. 1. Molecular view of the title compound with the atom-labelling scheme. Displacement ellipsoids are drawn at the 30% probability level [symmetry code: 1-x;1-y;1-z].




Fig. 2. The crystal packing of the title compound (a) and compound (II) (NISALO) (b). Hydrogen atoms are omitted for clarity.

Bis[2-(hydroxyiminomethyl)phenolato]nickel(II)

Crystal data

[Ni(C₇H₆NO₂)₂]

M_r = 330.97

Monoclinic, *P*2₁/*c*

Hall symbol: -*P* 2ybc

a = 4.9912 (2) Å

b = 7.4717 (3) Å

c = 17.4152 (7) Å

β = 90.653 (3)°

V = 649.42 (5) Å³

Z = 2

F(000) = 340

D_x = 1.693 Mg m⁻³

Mo *K*α radiation, λ = 0.71073 Å

Cell parameters from 1800 reflections

θ = 2.3–26.1°

μ = 1.51 mm⁻¹

T = 296 K

Prizm, green

0.53 × 0.36 × 0.11 mm

Data collection

Bruker SMART APEXII
diffractometer

Radiation source: fine-focus sealed tube
graphite

φ and ω scans

Absorption correction: numerical
(*SADABS*; Sheldrick, 2002)

T_{min} = 0.501, *T_{max}* = 0.851

4678 measured reflections

1342 independent reflections

1096 reflections with *I* > 2σ(*I*)

R_{int} = 0.028

θ_{max} = 26.4°, θ_{min} = 2.3°

h = -6→6

k = -9→8

l = -19→21

Refinement

Refinement on *F*²

Least-squares matrix: full

R[*F*² > 2σ(*F*²)] = 0.030

wR(*F*²) = 0.072

Primary atom site location: structure-invariant direct methods

Secondary atom site location: difference Fourier map

Hydrogen site location: inferred from neighbouring sites

All H-atom parameters refined

$S = 1.06$	$w = 1/[\sigma^2(F_o^2) + (0.030P)^2 + 0.3216P]$
1342 reflections	where $P = (F_o^2 + 2F_c^2)/3$
121 parameters	$(\Delta/\sigma)_{\max} = 0.005$
0 restraints	$\Delta\rho_{\max} = 0.28 \text{ e } \text{\AA}^{-3}$
	$\Delta\rho_{\min} = -0.26 \text{ e } \text{\AA}^{-3}$

Special details

Experimental. Numerical absorption corrections based on indexed crystal faces were applied using the Crystal Faces plugin in Bruker APEX2 software (Bruker, 2007)

Geometry. All e.s.d.'s (except the e.s.d. in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell e.s.d.'s are taken into account individually in the estimation of e.s.d.'s in distances, angles and torsion angles; correlations between e.s.d.'s in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell e.s.d.'s is used for estimating e.s.d.'s involving l.s. planes.

Refinement. Refinement of F^2 against ALL reflections. The weighted R -factor wR and goodness of fit S are based on F^2 , conventional R -factors R are based on F , with F set to zero for negative F^2 . The threshold expression of $F^2 > \sigma(F^2)$ is used only for calculating R -factors(gt) *etc.* and is not relevant to the choice of reflections for refinement. R -factors based on F^2 are statistically about twice as large as those based on F , and R -factors based on ALL data will be even larger.

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (\AA^2)

	x	y	z	$U_{\text{iso}}^*/U_{\text{eq}}$
C1	0.0820 (4)	0.3351 (3)	0.41130 (12)	0.0363 (5)
C2	0.0261 (5)	0.4911 (3)	0.36993 (13)	0.0383 (5)
C3	-0.1766 (5)	0.4888 (4)	0.31268 (15)	0.0471 (6)
C4	-0.3212 (6)	0.3375 (4)	0.29774 (16)	0.0540 (7)
C5	-0.2652 (6)	0.1845 (4)	0.33888 (16)	0.0530 (7)
C6	-0.0674 (5)	0.1815 (4)	0.39413 (15)	0.0482 (6)
C7	0.1642 (5)	0.6566 (3)	0.38506 (14)	0.0415 (6)
H3	-0.208 (5)	0.586 (3)	0.2867 (14)	0.040 (7)*
H4	-0.455 (6)	0.342 (4)	0.2600 (17)	0.071 (9)*
H5	-0.369 (6)	0.088 (4)	0.3311 (16)	0.064 (9)*
H6	-0.029 (5)	0.081 (4)	0.4202 (15)	0.048 (7)*
H7	0.106 (5)	0.757 (3)	0.3582 (13)	0.037 (6)*
H2	0.578 (7)	0.828 (5)	0.4672 (18)	0.074 (11)*
N1	0.3534 (4)	0.6733 (3)	0.43480 (11)	0.0386 (4)
Ni1	0.5000	0.5000	0.5000	0.03283 (15)
O1	0.2671 (3)	0.3269 (2)	0.46661 (9)	0.0413 (4)
O2	0.4542 (4)	0.8474 (3)	0.43860 (12)	0.0586 (5)

Atomic displacement parameters (\AA^2)

	U^{11}	U^{22}	U^{33}	U^{12}	U^{13}	U^{23}
C1	0.0337 (11)	0.0407 (13)	0.0346 (12)	-0.0023 (10)	0.0046 (9)	-0.0038 (10)
C2	0.0372 (11)	0.0425 (12)	0.0355 (12)	0.0022 (11)	0.0022 (9)	-0.0003 (11)
C3	0.0458 (14)	0.0516 (16)	0.0436 (14)	0.0049 (14)	-0.0054 (11)	0.0030 (13)

supplementary materials

C4	0.0467 (15)	0.069 (2)	0.0457 (15)	-0.0028 (15)	-0.0113 (12)	-0.0060 (15)
C5	0.0522 (16)	0.0531 (17)	0.0536 (16)	-0.0145 (14)	-0.0064 (13)	-0.0100 (14)
C6	0.0537 (16)	0.0421 (15)	0.0488 (15)	-0.0064 (13)	-0.0030 (12)	-0.0010 (12)
C7	0.0470 (14)	0.0384 (13)	0.0391 (13)	0.0027 (11)	-0.0033 (11)	0.0092 (11)
N1	0.0450 (11)	0.0306 (10)	0.0401 (10)	-0.0052 (9)	-0.0023 (9)	0.0034 (9)
Ni1	0.0366 (2)	0.0295 (2)	0.0324 (2)	-0.00221 (19)	-0.00104 (15)	0.00314 (18)
O1	0.0474 (10)	0.0329 (9)	0.0434 (9)	-0.0062 (7)	-0.0100 (7)	0.0056 (7)
O2	0.0717 (14)	0.0338 (10)	0.0698 (13)	-0.0151 (9)	-0.0272 (11)	0.0166 (9)

Geometric parameters (Å, °)

C1—O1	1.328 (3)	C5—C6	1.371 (4)
C1—C2	1.397 (3)	C5—H5	0.90 (3)
C1—C6	1.399 (3)	C6—H6	0.90 (3)
C2—C3	1.412 (3)	C7—N1	1.280 (3)
C2—C7	1.439 (3)	C7—H7	0.93 (2)
C3—C4	1.365 (4)	N1—O2	1.396 (3)
C3—H3	0.87 (2)	N1—Ni1	1.8661 (19)
C4—C5	1.376 (4)	Ni1—O1	1.8292 (16)
C4—H4	0.93 (3)	O2—H2	0.80 (3)
O1—C1—C2	123.1 (2)	C5—C6—C1	121.0 (3)
O1—C1—C6	118.8 (2)	C5—C6—H6	121.2 (17)
C2—C1—C6	118.1 (2)	C1—C6—H6	117.8 (18)
C1—C2—C3	119.4 (2)	N1—C7—C2	123.8 (2)
C1—C2—C7	122.0 (2)	N1—C7—H7	119.1 (15)
C3—C2—C7	118.5 (2)	C2—C7—H7	117.1 (15)
C4—C3—C2	121.2 (3)	C7—N1—O2	112.7 (2)
C4—C3—H3	120.3 (17)	C7—N1—Ni1	128.70 (18)
C2—C3—H3	118.6 (17)	O2—N1—Ni1	118.62 (15)
C3—C4—C5	119.0 (3)	O1 ⁱ —Ni1—O1	180.0
C3—C4—H4	118 (2)	O1 ⁱ —Ni1—N1	86.92 (8)
C5—C4—H4	123 (2)	O1—Ni1—N1	93.08 (8)
C6—C5—C4	121.3 (3)	N1 ⁱ —Ni1—N1	180.00 (9)
C6—C5—H5	120.1 (19)	C1—O1—Ni1	129.22 (15)
C4—C5—H5	118.5 (19)	N1—O2—H2	98 (3)

Symmetry codes: (i) $-x+1, -y+1, -z+1$.

Hydrogen-bond geometry (Å, °)

<i>D</i> —H... <i>A</i>	<i>D</i> —H	H... <i>A</i>	<i>D</i> ... <i>A</i>	<i>D</i> —H... <i>A</i>
O2—H2...O1 ⁱ	0.80 (3)	1.80 (3)	2.511 (2)	146 (4)

Symmetry codes: (i) $-x+1, -y+1, -z+1$.

Fig. 1




Fig. 2

