

Poly[[bis(2,2-bipyridine)bis[μ_6 -5-(carboxylatomoxy)benzene-1,3-dicarboxylato]trimanganese(II)] monohydrate]

Zhan-Guo Jiang, Jing Chen and Yun-Long Feng*

Zhejiang Key Laboratory for Reactive Chemistry on Solid Surfaces, College of Chemistry and Life Science, Zhejiang Normal University, Jinhua, Zhejiang 321004, People's Republic of China
Correspondence e-mail: sky37@zjnu.cn


Received 6 January 2011; accepted 20 January 2011

Key indicators: single-crystal X-ray study; $T = 296$ K; mean $\sigma(C-C) = 0.005$ Å; disorder in solvent or counterion; R factor = 0.036; wR factor = 0.123; data-to-parameter ratio = 11.5.

The title compound, $\{[\text{Mn}_3(\text{C}_{10}\text{H}_5\text{O}_7)_2(\text{C}_{10}\text{H}_8\text{N}_2)_2]\cdot\text{H}_2\text{O}\}_n$, was synthesized under hydrothermal conditions. Six carboxylate groups of six 5-(carboxylatomoxy)benzene-1,3-dicarboxylate anions (OABDC^{3-}) join three Mn^{II} ions into a trinuclear centrosymmetric $[\text{Mn}_3(\mu_2\text{-COO})_6]$ unit with one Mn site situated on a centre of inversion. The latter Mn^{II} ion exhibits a distorted MnO_6 coordination, whereas the other Mn^{II} ion has a trigonal-bipyramidal MnN_2O_3 coordination environment resulting from three carboxylate O atoms and the two N atoms of the bipyridine ligand. Adjacent units are linked to each other by OABDC^{3-} ligands into a layer parallel to (010). Within the layer, $\text{O}-\text{H}\cdots\text{O}$ hydrogen-bonding interactions involving the uncoordinated and half-occupied water molecule and the free carboxylate O atoms are observed. The layers stack along [010], constructing a three-dimensional structure through $\pi-\pi$ interactions between adjacent pyridine rings, with a centroid–centroid distance of 3.473 (5) Å.

Related literature

For the construction of metal-organic frameworks with polycarboxylate ligands, see: Xing *et al.* (2010); Cao *et al.* (2004, 2007).



Experimental

Crystal data

$[\text{Mn}_3(\text{C}_{10}\text{H}_5\text{O}_7)_2(\text{C}_{10}\text{H}_8\text{N}_2)_2]\cdot\text{H}_2\text{O}$	$V = 1919.41 (4)$ Å ³
$M_r = 969.48$	$Z = 2$
Monoclinic, $P2_1/n$	Mo $K\alpha$ radiation
$a = 8.5683 (1)$ Å	$\mu = 1.05$ mm ⁻¹
$b = 25.2280 (4)$ Å	$T = 296$ K
$c = 9.7685 (1)$ Å	$0.37 \times 0.19 \times 0.09$ mm
$\beta = 114.633 (1)$ °	

Data collection

Bruker APEXII area-detector diffractometer	16528 measured reflections
Absorption correction: multi-scan (SADABS; Sheldrick, 1996)	3350 independent reflections
$(SADABS; Sheldrick, 1996)$	3012 reflections with $I > 2\sigma(I)$
$T_{\min} = 0.786$, $T_{\max} = 0.907$	$R_{\text{int}} = 0.023$

Refinement

$R[F^2 > 2\sigma(F^2)] = 0.036$	H atoms treated by a mixture of independent and constrained refinement
$wR(F^2) = 0.123$	$\Delta\rho_{\max} = 1.11$ e Å ⁻³
$S = 1.10$	$\Delta\rho_{\min} = -0.47$ e Å ⁻³
3350 reflections	
292 parameters	
6 restraints	

Table 1
Selected bond lengths (Å).

$\text{Mn1}-\text{O}^{\text{i}}$	2.100 (2)	$\text{Mn1}-\text{N1}$	2.251 (3)
$\text{Mn1}-\text{O}^{\text{ii}}$	2.148 (2)	$\text{Mn2}-\text{O}^{\text{iv}}$	2.108 (2)
$\text{Mn1}-\text{O1}$	2.175 (2)	$\text{Mn2}-\text{O}^{\text{v}}$	2.159 (2)
$\text{Mn1}-\text{N2}$	2.228 (3)	$\text{Mn2}-\text{O1}$	2.322 (2)

Symmetry codes: (i) $x + 1, y, z + 1$; (ii) $x, y, z + 1$; (iii) $-x, -y, -z - 1$; (iv) $-x - 1, -y, -z - 1$.

Table 2Hydrogen-bond geometry (\AA , $^\circ$).

$D-\text{H}\cdots A$	$D-\text{H}$	$\text{H}\cdots A$	$D\cdots A$	$D-\text{H}\cdots A$
O1W—H1WB···O2 ⁱⁱⁱ	0.85 (1)	2.19 (2)	3.015 (5)	164 (6)

Symmetry code: (iii) $-x, -y, -z - 1$.

Data collection: *APEX2* (Bruker, 2006); cell refinement: *SAINT* (Bruker, 2006); data reduction: *SAINT*; program(s) used to solve structure: *SHELXS97* (Sheldrick, 2008); program(s) used to refine structure: *SHELXL97* (Sheldrick, 2008); molecular graphics: *DIAMOND* (Brandenburg, 2008); software used to prepare material for publication: *SHELXTL* (Sheldrick, 2008).

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: WM2447).

References

- Brandenburg, K. (2008). *DIAMOND*. Crystal Impact GbR, Bonn, Germany.
Bruker (2006). *APEX2* and *SAINT*. Bruker AXS Inc., Madison, Wisconsin, USA.
Cao, X.-Y., Zhang, J., Cheng, J.-K., Kang, Y. & Yao, Y.-G. (2004). *CrystEngComm*, **6**, 315–317.
Cao, X.-Y., Zhang, J., Li, Z.-J., Cheng, J.-K. & Yao, Y.-G. (2007). *CrystEngComm*, **9**, 806–814.
Sheldrick, G. M. (1996). *SADABS*. University of Göttingen, Germany.
Sheldrick, G. M. (2008). *Acta Cryst. A* **64**, 112–122.
Xing, X.-Y., Song, X.-Y., Yang, P.-P., Liu, R.-N., Li, L.-C. & Liao, D.-Z. (2010). *J. Mol. Struct.* **967**, 196–200.

supplementary materials

Acta Cryst. (2011). E67, m259-m260 [doi:10.1107/S1600536811002819]

Poly[[bis(2,2-bipyridine)bis μ_6 -5-(carboxylatomethoxy)benzene-1,3-dicarboxylato]trimanganese(II)] monohydrate]

Z.-G. Jiang, J. Chen and Y.-L. Feng

Comment

Aromatic polycarboxylic acids like 1,3,5-benzenetricarboxylate (H_3btc) and 1,2,4,5-benzenetetracarboxylate (H_4btec) are widely used to construct metal-organic frameworks (Xing *et al.*, 2010). Aromatic polycarboxylate ligands with rigid and flexible carboxyl group are less reported. 5-oxyacetato-1,3-benzenebiscarboxylic acid (H_3OABDC) is a ligand with two rigid carboxyl groups and one flexible oxyacetato group. Previous structural studies including this ligand have also been reported (Cao *et al.*, 2004, 2007).

The title compound, $[Mn_3(C_{10}H_5O_7)_2(C_{10}H_8N_2)_2]H_2O$, crystallizes with two Mn^{II} ions ($Mn1$ on a general position, $Mn2$ on a centre of inversion), one $OABDC^{3-}$ trianion, one chelating 2,2-bipyridine (bipy) ligand and one water molecule (half-occupied) in the asymmetric unit. $Mn1$ is five-coordinated by two N atoms from the bipy ligand, one O atom from one flexible carboxyl group of the $OABDC^{3-}$ anion and two O atoms from two rigid carboxyl groups of the $OABDC^{3-}$ anion to form a distorted trigonal-bipyramidal environment. $Mn2$ is six-coordinated by two O atoms from two flexible carboxyl groups of the $OABDC^{3-}$ anion and four oxygen atoms from four rigid carboxyl groups of adjacent $OABDC^{3-}$ anions to form a distorted octahedral environment (Table 1, Fig. 1).

Six $OABDC^{3-}$ anions join three Mn^{II} ions through their carboxyl groups. As a result, trinuclear $[Mn_3(\mu_2\text{-COO})_6]_n$ layers are formed in which $Mn2$ sits on a centre of inversion of each trinuclear unit. These layers are oriented parallel to (010). Uncoordinated water molecules interact with non-bonded O atoms of the carboxylate groups within the layers through relatively weak O—H···O hydrogen-bonding interactions (Table 2, Fig. 2).

The bipy ligands decorate the top and bottom of each layer. Between parallel pyridine rings of adjacent layers exist π — π interactions with a centroid–centroid distance of 3.473 (5) Å that consolidate the packing of the structure along [010] into a three-dimensional framework (Fig. 3).

Experimental

All reagents were purchased commercially and used without further purification. A mixture of 5-oxyacetato-1,3-benzenebiscarboxylic acid (H_3OABDC ; 0.1205 g, 0.5 mmol), $MnCl_2 \cdot 4H_2O$ (0.0976 g, 0.5 mmol), bipyridine (0.0786 g, 0.5 mmol), was dissolved with NaOH (0.0101 g, 0.25 mmol) in water (15 ml) and loaded in a 25 ml stainless steel reactor with a teflon liner. The autoclave was heated at 433 K for 72 h, and then cooled to room temperature over 3 days with a cooling rate of 5 K per hour. Yellow single crystals of the title compound were obtained by slow evaporation of the filtrate over a few days.

supplementary materials

Refinement

The carbon-bound H-atoms were positioned geometrically and included in the refinement using a riding model [C—H 0.93 for aromatic C atoms and 0.97 Å for methylen C atoms with $U_{\text{iso}}(\text{H}) = 1.2U_{\text{eq}}(\text{C})$]. The water H-atoms were located in a difference Fourier map and refined with an O—H distance restrained to 0.85 (2) Å [$U_{\text{iso}}(\text{H}) = 1.2U_{\text{eq}}(\text{O})$]. The lattice water (O1W) molecule shows half-occupation.

Figures




Fig. 1. The molecular structure of title the compound. Displacement ellipsoids are drawn at the 30% probability level. [Symmetry codes: (1) $x + 1, y, z + 1$; (2) $x, y, z + 1$; (3) $-x, -y, -z - 1$; (4) $-x - 1, -y, -z - 1$; (5) $-x, -y, -z$.]




Fig. 2. two-dimensional layered structure (bipys are omitted for clarity).




Fig. 3. View of $\pi-\pi$ interactions between layers.

Poly[[bis(2,2-bipyridine)bis[μ_6 -5-(carboxylatomoxy)benzene-1,3-dicarboxylato]trimanganese(II)] monohydrate]

Crystal data

$[\text{Mn}_3(\text{C}_{10}\text{H}_5\text{O}_7)_2(\text{C}_{10}\text{H}_8\text{N}_2)_2]\cdot\text{H}_2\text{O}$	$F(000) = 982$
$M_r = 969.48$	$D_x = 1.677 \text{ Mg m}^{-3}$
Monoclinic, $P2_1/n$	Mo $K\alpha$ radiation, $\lambda = 0.71073 \text{ \AA}$
Hall symbol: -P 2yn	Cell parameters from 8045 reflections
$a = 8.5683 (1) \text{ \AA}$	$\theta = 2.4\text{--}27.7^\circ$
$b = 25.2280 (4) \text{ \AA}$	$\mu = 1.05 \text{ mm}^{-1}$
$c = 9.7685 (1) \text{ \AA}$	$T = 296 \text{ K}$
$\beta = 114.633 (1)^\circ$	Block, yellow
$V = 1919.41 (4) \text{ \AA}^3$	$0.37 \times 0.19 \times 0.09 \text{ mm}$
$Z = 2$	

Data collection

Bruker APEXII area-detector diffractometer	3350 independent reflections
Radiation source: fine-focus sealed tube graphite	3012 reflections with $I > 2\sigma(I)$
ω scans	$R_{\text{int}} = 0.023$
Absorption correction: multi-scan (<i>SADABS</i> ; Sheldrick, 1996)	$\theta_{\text{max}} = 25.0^\circ$, $\theta_{\text{min}} = 2.4^\circ$
$T_{\text{min}} = 0.786$, $T_{\text{max}} = 0.907$	$h = -9 \rightarrow 8$
16528 measured reflections	$k = -24 \rightarrow 30$
	$l = -11 \rightarrow 11$

Refinement

Refinement on F^2	Primary atom site location: structure-invariant direct methods
Least-squares matrix: full	Secondary atom site location: difference Fourier map
$R[F^2 > 2\sigma(F^2)] = 0.036$	Hydrogen site location: inferred from neighbouring sites
$wR(F^2) = 0.123$	H atoms treated by a mixture of independent and constrained refinement
$S = 1.10$	$w = 1/[\sigma^2(F_o^2) + (0.0743P)^2 + 1.9128P]$
3350 reflections	where $P = (F_o^2 + 2F_c^2)/3$
292 parameters	$(\Delta/\sigma)_{\text{max}} < 0.001$
6 restraints	$\Delta\rho_{\text{max}} = 1.11 \text{ e \AA}^{-3}$
	$\Delta\rho_{\text{min}} = -0.47 \text{ e \AA}^{-3}$

Special details

Geometry. All e.s.d.'s (except the e.s.d. in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell e.s.d.'s are taken into account individually in the estimation of e.s.d.'s in distances, angles and torsion angles; correlations between e.s.d.'s in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell e.s.d.'s is used for estimating e.s.d.'s involving l.s. planes.

Refinement. Refinement of F^2 against ALL reflections. The weighted R -factor wR and goodness of fit S are based on F^2 , conventional R -factors R are based on F , with F set to zero for negative F^2 . The threshold expression of $F^2 > \sigma(F^2)$ is used only for calculating R -factors(gt) etc. and is not relevant to the choice of reflections for refinement. R -factors based on F^2 are statistically about twice as large as those based on F , and R -factors based on ALL data will be even larger.

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (\AA^2)

	<i>x</i>	<i>y</i>	<i>z</i>	$U_{\text{iso}}^*/U_{\text{eq}}$	Occ. (<1)
Mn1	0.23443 (6)	0.121021 (17)	0.11506 (5)	0.02566 (17)	
Mn2	0.0000	0.0000	0.0000	0.02189 (18)	
O1W	-0.6212 (6)	-0.12729 (17)	-0.9008 (6)	0.0426 (12)	0.50
H1WB	-0.5137 (17)	-0.130 (3)	-0.874 (7)	0.051*	0.50
H1WA	-0.659 (7)	-0.111 (3)	-0.984 (5)	0.051*	0.50
O1	0.0588 (3)	0.08005 (8)	-0.0861 (2)	0.0297 (5)	

supplementary materials

O2	0.2531 (3)	0.12144 (9)	-0.1396 (2)	0.0353 (5)
O3	0.1568 (3)	0.11891 (9)	-0.7023 (3)	0.0416 (6)
O4	0.0038 (3)	0.04991 (9)	-0.8255 (2)	0.0387 (6)
O5	-0.3626 (3)	0.00513 (9)	-0.5459 (2)	0.0337 (5)
O6	-0.7274 (3)	-0.01469 (9)	-0.8866 (2)	0.0337 (5)
O7	-0.5860 (3)	0.06004 (9)	-0.7894 (3)	0.0408 (6)
N1	0.1153 (4)	0.20221 (11)	0.0619 (3)	0.0344 (6)
N2	0.4486 (3)	0.17919 (10)	0.1761 (3)	0.0308 (6)
C1	0.0167 (3)	0.08164 (10)	-0.3438 (3)	0.0207 (6)
C2	0.0778 (4)	0.09280 (10)	-0.4528 (3)	0.0213 (6)
H2	0.1769	0.1128	-0.4285	0.026*
C3	-0.0115 (4)	0.07361 (10)	-0.5978 (3)	0.0217 (6)
C4	-0.1610 (4)	0.04408 (11)	-0.6359 (3)	0.0227 (6)
H4	-0.2187	0.0308	-0.7330	0.027*
C5	-0.2231 (3)	0.03463 (11)	-0.5281 (3)	0.0228 (6)
C6	-0.1349 (4)	0.05419 (11)	-0.3831 (3)	0.0232 (6)
H6	-0.1786	0.0487	-0.3116	0.028*
C7	0.1182 (4)	0.09562 (10)	-0.1816 (3)	0.0237 (6)
C8	0.0547 (4)	0.08148 (12)	-0.7170 (3)	0.0269 (6)
C9	-0.4589 (4)	-0.02186 (12)	-0.6826 (3)	0.0306 (7)
H9A	-0.5073	-0.0537	-0.6605	0.037*
H9B	-0.3821	-0.0326	-0.7277	0.037*
C10	-0.6025 (4)	0.01101 (12)	-0.7951 (3)	0.0264 (6)
C11	-0.0531 (5)	0.21147 (17)	0.0082 (4)	0.0468 (9)
H11	-0.1278	0.1827	-0.0174	0.056*
C12	-0.1217 (6)	0.2624 (2)	-0.0112 (4)	0.0587 (12)
H12	-0.2397	0.2678	-0.0514	0.070*
C13	-0.0103 (6)	0.30430 (18)	0.0307 (5)	0.0613 (12)
H13	-0.0521	0.3388	0.0210	0.074*
C14	0.1621 (6)	0.29541 (15)	0.0868 (4)	0.0514 (10)
H14	0.2382	0.3238	0.1155	0.062*
C15	0.2239 (4)	0.24397 (13)	0.1008 (3)	0.0342 (7)
C16	0.4077 (4)	0.23094 (12)	0.1557 (3)	0.0318 (7)
C17	0.5338 (5)	0.26898 (13)	0.1875 (4)	0.0440 (9)
H17	0.5040	0.3046	0.1724	0.053*
C18	0.7043 (5)	0.25413 (15)	0.2416 (4)	0.0479 (9)
H18	0.7899	0.2796	0.2644	0.057*
C19	0.7455 (5)	0.20131 (15)	0.2613 (4)	0.0456 (9)
H19	0.8589	0.1900	0.2968	0.055*
C20	0.6131 (4)	0.16548 (13)	0.2268 (4)	0.0381 (8)
H20	0.6403	0.1296	0.2397	0.046*

Atomic displacement parameters (\AA^2)

	U^{11}	U^{22}	U^{33}	U^{12}	U^{13}	U^{23}
Mn1	0.0318 (3)	0.0250 (3)	0.0198 (3)	-0.00594 (17)	0.0103 (2)	-0.00228 (16)
Mn2	0.0203 (3)	0.0264 (3)	0.0174 (3)	-0.0036 (2)	0.0062 (2)	-0.0038 (2)
O1W	0.031 (2)	0.022 (2)	0.081 (4)	-0.0176 (19)	0.030 (3)	-0.026 (2)

O1	0.0381 (12)	0.0346 (11)	0.0175 (9)	-0.0066 (9)	0.0128 (9)	-0.0019 (8)
O2	0.0355 (13)	0.0405 (13)	0.0266 (11)	-0.0142 (10)	0.0096 (10)	-0.0066 (9)
O3	0.0529 (15)	0.0437 (14)	0.0397 (13)	-0.0147 (11)	0.0308 (12)	-0.0004 (10)
O4	0.0457 (14)	0.0523 (14)	0.0230 (11)	-0.0026 (11)	0.0191 (10)	-0.0108 (10)
O5	0.0225 (11)	0.0500 (13)	0.0265 (11)	-0.0143 (9)	0.0081 (9)	-0.0015 (9)
O6	0.0240 (11)	0.0355 (12)	0.0331 (11)	-0.0025 (9)	0.0036 (9)	-0.0045 (9)
O7	0.0329 (12)	0.0268 (12)	0.0507 (14)	-0.0042 (9)	0.0056 (11)	0.0036 (10)
N1	0.0399 (16)	0.0386 (15)	0.0291 (13)	0.0006 (12)	0.0188 (12)	0.0027 (11)
N2	0.0389 (15)	0.0249 (13)	0.0306 (13)	-0.0035 (11)	0.0164 (12)	-0.0029 (10)
C1	0.0251 (14)	0.0191 (13)	0.0180 (13)	0.0014 (11)	0.0091 (11)	0.0008 (10)
C2	0.0230 (14)	0.0194 (13)	0.0215 (13)	-0.0018 (11)	0.0093 (11)	-0.0003 (10)
C3	0.0256 (14)	0.0219 (14)	0.0192 (13)	0.0039 (11)	0.0110 (11)	0.0018 (10)
C4	0.0222 (14)	0.0259 (14)	0.0168 (12)	-0.0006 (11)	0.0050 (11)	-0.0017 (10)
C5	0.0181 (13)	0.0262 (14)	0.0225 (13)	0.0007 (11)	0.0069 (11)	0.0039 (11)
C6	0.0233 (14)	0.0288 (14)	0.0181 (13)	0.0016 (11)	0.0091 (11)	0.0034 (11)
C7	0.0302 (15)	0.0199 (13)	0.0200 (13)	0.0019 (12)	0.0096 (12)	-0.0011 (11)
C8	0.0298 (16)	0.0321 (16)	0.0214 (14)	0.0043 (13)	0.0132 (12)	0.0056 (12)
C9	0.0236 (15)	0.0301 (16)	0.0330 (16)	-0.0061 (12)	0.0067 (13)	-0.0023 (13)
C10	0.0220 (15)	0.0307 (16)	0.0273 (15)	-0.0029 (12)	0.0111 (12)	-0.0002 (12)
C11	0.044 (2)	0.062 (2)	0.0393 (19)	0.0037 (18)	0.0220 (16)	0.0068 (17)
C12	0.057 (2)	0.085 (3)	0.042 (2)	0.029 (2)	0.0267 (19)	0.013 (2)
C13	0.081 (3)	0.052 (3)	0.053 (2)	0.029 (2)	0.030 (2)	0.008 (2)
C14	0.068 (3)	0.0360 (19)	0.051 (2)	0.0100 (19)	0.025 (2)	0.0001 (16)
C15	0.050 (2)	0.0301 (16)	0.0258 (15)	0.0028 (14)	0.0192 (14)	0.0017 (12)
C16	0.0471 (19)	0.0265 (15)	0.0246 (15)	-0.0049 (14)	0.0177 (14)	-0.0018 (12)
C17	0.063 (3)	0.0251 (16)	0.048 (2)	-0.0093 (16)	0.0268 (18)	-0.0055 (14)
C18	0.053 (2)	0.046 (2)	0.052 (2)	-0.0217 (18)	0.0289 (18)	-0.0113 (17)
C19	0.041 (2)	0.048 (2)	0.051 (2)	-0.0097 (17)	0.0229 (17)	-0.0070 (17)
C20	0.0378 (19)	0.0321 (17)	0.0450 (19)	-0.0029 (14)	0.0179 (15)	-0.0035 (14)

Geometric parameters (\AA , $^\circ$)

Mn1—O7 ⁱ	2.100 (2)	C1—C2	1.396 (4)
Mn1—O3 ⁱⁱ	2.148 (2)	C1—C7	1.498 (4)
Mn1—O1	2.175 (2)	C2—C3	1.386 (4)
Mn1—N2	2.228 (3)	C2—H2	0.9300
Mn1—N1	2.251 (3)	C3—C4	1.392 (4)
Mn2—O4 ⁱⁱ	2.108 (2)	C3—C8	1.506 (4)
Mn2—O4 ⁱⁱⁱ	2.108 (2)	C4—C5	1.384 (4)
Mn2—O6 ^{iv}	2.159 (2)	C4—H4	0.9300
Mn2—O6 ⁱ	2.159 (2)	C5—C6	1.389 (4)
Mn2—O1 ^v	2.322 (2)	C6—H6	0.9300
Mn2—O1	2.322 (2)	C9—C10	1.511 (4)
O1W—H1WB	0.849 (2)	C9—H9A	0.9700
O1W—H1WA	0.849 (2)	C9—H9B	0.9700
O1—C7	1.296 (3)	C11—C12	1.394 (6)
O2—C7	1.238 (4)	C11—H11	0.9300
O3—C8	1.254 (4)	C12—C13	1.366 (7)

supplementary materials

O3—Mn1 ^{vi}	2.148 (2)	C12—H12	0.9300
O4—C8	1.250 (4)	C13—C14	1.363 (6)
O4—Mn2 ^{vi}	2.108 (2)	C13—H13	0.9300
O5—C5	1.356 (3)	C14—C15	1.387 (5)
O5—C9	1.417 (4)	C14—H14	0.9300
O6—C10	1.252 (4)	C15—C16	1.474 (5)
O6—Mn2 ^{vii}	2.159 (2)	C16—C17	1.380 (5)
O7—C10	1.244 (4)	C17—C18	1.382 (6)
O7—Mn1 ^{vii}	2.100 (2)	C17—H17	0.9300
N1—C11	1.334 (5)	C18—C19	1.371 (5)
N1—C15	1.351 (4)	C18—H18	0.9300
N2—C20	1.330 (4)	C19—C20	1.378 (5)
N2—C16	1.345 (4)	C19—H19	0.9300
C1—C6	1.378 (4)	C20—H20	0.9300
O7 ⁱ —Mn1—O3 ⁱⁱ	92.04 (10)	O5—C5—C4	126.5 (2)
O7 ⁱ —Mn1—O1	99.00 (9)	O5—C5—C6	113.7 (2)
O3 ⁱⁱ —Mn1—O1	113.65 (9)	C4—C5—C6	119.7 (3)
O7 ⁱ —Mn1—N2	89.68 (9)	C1—C6—C5	120.8 (3)
O3 ⁱⁱ —Mn1—N2	107.44 (9)	C1—C6—H6	119.6
O1—Mn1—N2	137.52 (8)	C5—C6—H6	119.6
O7 ⁱ —Mn1—N1	161.31 (10)	O2—C7—O1	120.9 (2)
O3 ⁱⁱ —Mn1—N1	86.93 (9)	O2—C7—C1	121.6 (2)
O1—Mn1—N1	98.49 (9)	O1—C7—C1	117.5 (2)
N2—Mn1—N1	72.92 (10)	O4—C8—O3	123.8 (3)
O4 ⁱⁱ —Mn2—O4 ⁱⁱⁱ	180.00 (12)	O4—C8—C3	117.5 (3)
O4 ⁱⁱ —Mn2—O6 ^{iv}	87.76 (9)	O3—C8—C3	118.7 (3)
O4 ⁱⁱⁱ —Mn2—O6 ^{iv}	92.24 (9)	O5—C9—C10	113.6 (2)
O4 ⁱⁱ —Mn2—O6 ⁱ	92.24 (9)	O5—C9—H9A	108.9
O4 ⁱⁱⁱ —Mn2—O6 ⁱ	87.76 (9)	C10—C9—H9A	108.9
O6 ^{iv} —Mn2—O6 ⁱ	180.00 (11)	O5—C9—H9B	108.9
O4 ⁱⁱ —Mn2—O1 ^v	99.16 (8)	C10—C9—H9B	108.9
O4 ⁱⁱⁱ —Mn2—O1 ^v	80.84 (8)	H9A—C9—H9B	107.7
O6 ^{iv} —Mn2—O1 ^v	89.06 (8)	O7—C10—O6	126.5 (3)
O6 ⁱ —Mn2—O1 ^v	90.94 (8)	O7—C10—C9	118.0 (3)
O4 ⁱⁱ —Mn2—O1	80.84 (8)	O6—C10—C9	115.5 (3)
O4 ⁱⁱⁱ —Mn2—O1	99.16 (8)	N1—C11—C12	122.7 (4)
O6 ^{iv} —Mn2—O1	90.94 (8)	N1—C11—H11	118.6
O6 ⁱ —Mn2—O1	89.06 (8)	C12—C11—H11	118.6
O1 ^v —Mn2—O1	180.00 (5)	C13—C12—C11	118.1 (4)
H1WB—O1W—H1WA	105.3 (4)	C13—C12—H12	121.0
C7—O1—Mn1	100.06 (17)	C11—C12—H12	121.0
C7—O1—Mn2	137.20 (18)	C14—C13—C12	119.8 (4)
Mn1—O1—Mn2	105.02 (8)	C14—C13—H13	120.1

C8—O3—Mn1 ^{vi}	111.51 (19)	C12—C13—H13	120.1
C8—O4—Mn2 ^{vi}	161.5 (2)	C13—C14—C15	119.9 (4)
C5—O5—C9	121.3 (2)	C13—C14—H14	120.0
C10—O6—Mn2 ^{vii}	134.2 (2)	C15—C14—H14	120.0
C10—O7—Mn1 ^{vii}	131.3 (2)	N1—C15—C14	120.8 (3)
C11—N1—C15	118.6 (3)	N1—C15—C16	115.8 (3)
C11—N1—Mn1	124.2 (3)	C14—C15—C16	123.4 (3)
C15—N1—Mn1	116.8 (2)	N2—C16—C17	120.7 (3)
C20—N2—C16	118.6 (3)	N2—C16—C15	116.3 (3)
C20—N2—Mn1	123.7 (2)	C17—C16—C15	123.0 (3)
C16—N2—Mn1	117.7 (2)	C16—C17—C18	120.1 (3)
C6—C1—C2	119.9 (2)	C16—C17—H17	120.0
C6—C1—C7	118.5 (2)	C18—C17—H17	120.0
C2—C1—C7	121.5 (2)	C19—C18—C17	119.1 (3)
C3—C2—C1	119.1 (2)	C19—C18—H18	120.5
C3—C2—H2	120.5	C17—C18—H18	120.5
C1—C2—H2	120.5	C18—C19—C20	117.8 (4)
C2—C3—C4	121.0 (2)	C18—C19—H19	121.1
C2—C3—C8	121.3 (2)	C20—C19—H19	121.1
C4—C3—C8	117.7 (2)	N2—C20—C19	123.8 (3)
C5—C4—C3	119.4 (2)	N2—C20—H20	118.1
C5—C4—H4	120.3	C19—C20—H20	118.1
C3—C4—H4	120.3		
O7 ⁱ —Mn1—O1—C7	−94.71 (18)	Mn1—O1—C7—C1	−172.1 (2)
O3 ⁱⁱ —Mn1—O1—C7	169.05 (17)	Mn2—O1—C7—C1	62.5 (4)
N2—Mn1—O1—C7	4.8 (2)	C6—C1—C7—O2	−178.1 (3)
N1—Mn1—O1—C7	78.69 (18)	C2—C1—C7—O2	6.0 (4)
O7 ⁱ —Mn1—O1—Mn2	50.28 (10)	C6—C1—C7—O1	1.3 (4)
O3 ⁱⁱ —Mn1—O1—Mn2	−45.96 (12)	C2—C1—C7—O1	−174.6 (2)
N2—Mn1—O1—Mn2	149.75 (11)	Mn2 ^{vi} —O4—C8—O3	36.8 (8)
N1—Mn1—O1—Mn2	−136.32 (9)	Mn2 ^{vi} —O4—C8—C3	−143.2 (6)
O4 ⁱⁱ —Mn2—O1—C7	162.4 (3)	Mn1 ^{vi} —O3—C8—O4	−0.7 (4)
O4 ⁱⁱⁱ —Mn2—O1—C7	−17.6 (3)	Mn1 ^{vi} —O3—C8—C3	179.27 (19)
O6 ^{iv} —Mn2—O1—C7	−110.0 (3)	C2—C3—C8—O4	157.0 (3)
O6 ⁱ —Mn2—O1—C7	70.0 (3)	C4—C3—C8—O4	−19.8 (4)
O4 ⁱⁱ —Mn2—O1—Mn1	38.66 (9)	C2—C3—C8—O3	−23.0 (4)
O4 ⁱⁱⁱ —Mn2—O1—Mn1	−141.34 (9)	C4—C3—C8—O3	160.2 (3)
O6 ^{iv} —Mn2—O1—Mn1	126.24 (9)	C5—O5—C9—C10	−89.4 (3)
O6 ⁱ —Mn2—O1—Mn1	−53.76 (9)	Mn1 ^{vii} —O7—C10—O6	15.9 (5)
O7 ⁱ —Mn1—N1—C11	−156.2 (3)	Mn1 ^{vii} —O7—C10—C9	−164.6 (2)
O3 ⁱⁱ —Mn1—N1—C11	−68.9 (3)	Mn2 ^{vii} —O6—C10—O7	−32.8 (5)
O1—Mn1—N1—C11	44.5 (3)	Mn2 ^{vii} —O6—C10—C9	147.6 (2)
N2—Mn1—N1—C11	−178.2 (3)	O5—C9—C10—O7	27.3 (4)
O7 ⁱ —Mn1—N1—C15	17.3 (4)	O5—C9—C10—O6	−153.1 (3)

supplementary materials

O3 ⁱⁱ —Mn1—N1—C15	104.6 (2)	C15—N1—C11—C12	0.8 (5)
O1—Mn1—N1—C15	−141.9 (2)	Mn1—N1—C11—C12	174.2 (3)
N2—Mn1—N1—C15	−4.7 (2)	N1—C11—C12—C13	−1.7 (6)
O7 ⁱ —Mn1—N2—C20	9.9 (3)	C11—C12—C13—C14	1.2 (6)
O3 ⁱⁱ —Mn1—N2—C20	102.0 (3)	C12—C13—C14—C15	0.1 (6)
O1—Mn1—N2—C20	−93.1 (3)	C11—N1—C15—C14	0.6 (4)
N1—Mn1—N2—C20	−177.0 (3)	Mn1—N1—C15—C14	−173.3 (2)
O7 ⁱ —Mn1—N2—C16	−172.2 (2)	C11—N1—C15—C16	−178.5 (3)
O3 ⁱⁱ —Mn1—N2—C16	−80.2 (2)	Mn1—N1—C15—C16	7.6 (3)
O1—Mn1—N2—C16	84.7 (2)	C13—C14—C15—N1	−1.1 (5)
N1—Mn1—N2—C16	0.9 (2)	C13—C14—C15—C16	178.0 (3)
C6—C1—C2—C3	−3.2 (4)	C20—N2—C16—C17	−0.4 (4)
C7—C1—C2—C3	172.6 (2)	Mn1—N2—C16—C17	−178.3 (2)
C1—C2—C3—C4	0.7 (4)	C20—N2—C16—C15	−179.4 (3)
C1—C2—C3—C8	−176.0 (2)	Mn1—N2—C16—C15	2.7 (3)
C2—C3—C4—C5	1.3 (4)	N1—C15—C16—N2	−6.8 (4)
C8—C3—C4—C5	178.1 (2)	C14—C15—C16—N2	174.1 (3)
C9—O5—C5—C4	2.6 (4)	N1—C15—C16—C17	174.2 (3)
C9—O5—C5—C6	−174.3 (2)	C14—C15—C16—C17	−4.9 (5)
C3—C4—C5—O5	−177.4 (3)	N2—C16—C17—C18	−0.3 (5)
C3—C4—C5—C6	−0.7 (4)	C15—C16—C17—C18	178.6 (3)
C2—C1—C6—C5	3.8 (4)	C16—C17—C18—C19	0.8 (5)
C7—C1—C6—C5	−172.1 (2)	C17—C18—C19—C20	−0.5 (5)
O5—C5—C6—C1	175.2 (2)	C16—N2—C20—C19	0.6 (5)
C4—C5—C6—C1	−1.9 (4)	Mn1—N2—C20—C19	178.4 (3)
Mn1—O1—C7—O2	7.3 (3)	C18—C19—C20—N2	−0.2 (6)
Mn2—O1—C7—O2	−118.1 (3)		


Symmetry codes: (i) $x+1, y, z+1$; (ii) $x, y, z+1$; (iii) $-x, -y, -z-1$; (iv) $-x-1, -y, -z-1$; (v) $-x, -y, -z$; (vi) $x, y, z-1$; (vii) $x-1, y, z-1$.

Hydrogen-bond geometry (\AA , $^\circ$)

$D\text{—H}\cdots A$	$D\text{—H}$	$H\cdots A$	$D\cdots A$	$D\text{—H}\cdots A$
O1W—H1WB···O2 ⁱⁱⁱ	0.85 (1)	2.19 (2)	3.015 (5)	164 (6)

Symmetry codes: (iii) $-x, -y, -z-1$.

Fig. 1



supplementary materials

Fig. 2




Fig. 3

