

## Bis( $\mu_2$ -quinoline-2-carboxylato)- $\kappa^3N,O^1:O^1$ ; $\kappa^3O^1:N,O^1$ -bis[(acetato- $\kappa O$ )-(ethanol- $\kappa O$ )]lead(II)

Ezzatollah Najafi,<sup>a</sup> Mostafa M. Amini<sup>a</sup> and Seik Weng Ng<sup>b\*</sup>

<sup>a</sup>Department of Chemistry, General Campus, Shahid Beheshti University, Tehran 1983963113, Iran, and <sup>b</sup>Department of Chemistry, University of Malaya, 50603

Kuala Lumpur, Malaysia

Correspondence e-mail: seikweng@um.edu.my


Received 17 January 2011; accepted 18 January 2011

Key indicators: single-crystal X-ray study;  $T = 100$  K; mean  $\sigma(C-C) = 0.007$  Å;  $R$  factor = 0.029;  $wR$  factor = 0.073; data-to-parameter ratio = 17.3.

In the centrosymmetric dinuclear title compound,  $[Pb_2(C_{10}H_6NO_2)_2(CH_3COO)_2(C_2H_5OH)_2]$ , one O atom of the carboxylate group of the quinoline-2-carboxylate anion connects the two  $Pb^{II}$  atoms. The  $Pb^{II}$  atom is surrounded by four O atoms and one N atom in a  $\Psi$ -octahedral  $PbO_4NE$  geometry ( $E$  is the electron lone pair). Two longer  $Pb \cdots O$  interactions distort the geometry towards a  $\Psi$ -square-antiprism. Intermolecular  $O-H \cdots O$  hydrogen bonds link the molecules.

### Related literature

For the analogous methanol-coordinated compound, see: Mohammadnezhad *et al.* (2010).



### Experimental

#### Crystal data

$[Pb_2(C_{10}H_6NO_2)_2(C_2H_3O_2)_2 \cdot (C_2H_5O)_2]$

$M_r = 968.92$   
Monoclinic,  $P2_1/c$

$a = 7.3419$  (1) Å  
 $b = 8.4004$  (1) Å  
 $c = 23.8008$  (4) Å  
 $\beta = 93.722$  (1)°  
 $V = 1464.82$  (4) Å<sup>3</sup>

$Z = 2$   
Mo  $K\alpha$  radiation  
 $\mu = 11.54$  mm<sup>-1</sup>  
 $T = 100$  K  
 $0.20 \times 0.20 \times 0.05$  mm

#### Data collection

Agilent Technologies SuperNova  
Dual diffractometer with an Atlas  
detector  
Absorption correction: multi-scan  
(*CrysAlis PRO*; Agilent

Technologies, 2010)  
 $T_{min} = 0.206$ ,  $T_{max} = 0.596$   
12668 measured reflections  
3314 independent reflections  
3010 reflections with  $I > 2\sigma(I)$   
 $R_{int} = 0.049$

#### Refinement

$R[F^2 > 2\sigma(F^2)] = 0.029$   
 $wR(F^2) = 0.073$   
 $S = 1.08$   
3314 reflections

192 parameters  
H-atom parameters constrained  
 $\Delta\rho_{max} = 1.21$  e Å<sup>-3</sup>  
 $\Delta\rho_{min} = -1.57$  e Å<sup>-3</sup>

**Table 1**

Selected bond lengths (Å).

Pb1—O1	2.377 (4)	Pb1—O5	2.694 (4)
Pb1—O3	2.384 (4)	Pb1—O4	2.763 (3)
Pb1—O1 <sup>i</sup>	2.500 (3)	Pb1—O2 <sup>ii</sup>	3.096 (4)
Pb1—N1	2.645 (4)		

Symmetry codes: (i)  $-x + 1, -y + 1, -z + 1$ ; (ii)  $x + 1, y, z$ .

**Table 2**

Hydrogen-bond geometry (Å, °).

$D-H \cdots A$	$D-H$	$H \cdots A$	$D \cdots A$	$D-H \cdots A$
$O5-H5 \cdots O3^i$	0.84	2.36	2.710 (5)	106

Symmetry code: (i)  $-x + 1, -y + 1, -z + 1$ .

Data collection: *CrysAlis PRO* (Agilent Technologies, 2010); cell refinement: *CrysAlis PRO*; data reduction: *CrysAlis PRO*; program(s) used to solve structure: *SHELXS97* (Sheldrick, 2008); program(s) used to refine structure: *SHELXL97* (Sheldrick, 2008); molecular graphics: *X-SEED* (Barbour, 2001); software used to prepare material for publication: *publCIF* (Westrip, 2010).

We thank Shahid Beheshti University and the University of Malaya for supporting this study.

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: BT5469).

### References

- Agilent Technologies (2010). *CrysAlis PRO*. Agilent Technologies, Yarnton, England.  
Barbour, L. J. (2001). *J. Supramol. Chem.* **1**, 189–191.  
Mohammadnezhad, G., Ghanbarpour, A. R., Amini, M. M. & Ng, S. W. (2010). *Acta Cryst.* **E66**, m963.  
Sheldrick, G. M. (2008). *Acta Cryst.* **A64**, 112–122.  
Westrip, S. P. (2010). *J. Appl. Cryst.* **43**, 920–925.

**supplementary materials**

*Acta Cryst.* (2011). E67, m264 [ doi:10.1107/S1600536811002509 ]

**Bis( $\mu_2$ -quinoline-2-carboxylato)- $\kappa^3N,O^1:O^1$ ;  $\kappa^3O^1:N,O^1$ -bis[(acetato- $\kappa O$ )(ethanol- $\kappa O$ )lead(II)]**

**E. Najafi, M. M. Amini and S. W. Ng**

**Comment**

A previous study of the methanol adduct of the title dinuclear compound,  $[\text{Pb}(\text{C}_{10}\text{H}_6\text{NO}_2)(\text{C}_2\text{H}_3\text{O}_2)(\text{CH}_3\text{OH})_2]_2$ , has found a  $\Psi$ -octahedral geometry for the lead(II) atom. Two longer  $\text{Pb}\cdots\text{O}$  interactions distort the geometry towards a  $\Psi$ -square-antiprism (Mohammadnezhad *et al.*, 2010). Replacing the methanol solvent system by ethanol leads to the analogous ethanol adduct (Scheme I, Fig. 1), which has a similar structure.

**Experimental**

Lead(II) acetate (1 mmol) and quinoline-2-carboxylic acid (1 mmol) were loaded into a convection tube; the tube was filled with dry ethanol and kept at 333 K. Colorless crystals were collected from the side arm after several days.

**Refinement**

H atoms were placed in calculated positions [C—H 0.95 to 0.98, O—H 0.84 Å,  $U_{\text{iso}}(\text{H})$  1.2 to 1.5  $U_{\text{eq}}(\text{C}, \text{O})$ ] and were included in the refinement in the riding model approximation.

The final difference Fourier map had a peak/hole in the vicinity of Pb1.

**Figures**




Fig. 1. Anisotropic displacement ellipsoid plot (Barbour, 2001) of the title compound at the 70% probability level; hydrogen atoms are drawn as spheres of arbitrary radius. The longer interactions that raise the coordination number are shown as dashed lines.

**Bis( $\mu_2$ -quinoline-2-carboxylato)- $\kappa^3N,O^1:O^1$ ;  $\kappa^3O^1:N,O^1$ -bis[(acetato- $\kappa O$ )(ethanol-  $\kappa O$ )lead(II)]**

*Crystal data*

$[\text{Pb}_2(\text{C}_{10}\text{H}_6\text{NO}_2)_2(\text{C}_2\text{H}_3\text{O}_2)_2(\text{C}_2\text{H}_6\text{O})_2]$

$M_r = 968.92$

Monoclinic,  $P2_1/c$

Hall symbol: -P 2ybc

$a = 7.3419(1)$  Å

$F(000) = 912$

$D_x = 2.197$  Mg m<sup>-3</sup>

Mo  $K\alpha$  radiation,  $\lambda = 0.71073$  Å

Cell parameters from 7627 reflections

$\theta = 2.4\text{--}29.3^\circ$

# supplementary materials

$b = 8.4004 (1) \text{ \AA}$   
 $c = 23.8008 (4) \text{ \AA}$   
 $\beta = 93.722 (1)^\circ$   
 $V = 1464.82 (4) \text{ \AA}^3$   
 $Z = 2$

$\mu = 11.54 \text{ mm}^{-1}$   
 $T = 100 \text{ K}$   
Prim, colorless  
 $0.20 \times 0.20 \times 0.05 \text{ mm}$

## Data collection

Agilent Technologies SuperNova Dual diffractometer with an Atlas detector  
Radiation source: SuperNova (Mo) X-ray Source  
Mirror  
Detector resolution:  $10.4041 \text{ pixels mm}^{-1}$   
 $\omega$  scan  
Absorption correction: multi-scan (*Crys.Alis PRO*; Agilent Technologies, 2010)  
 $T_{\min} = 0.206$ ,  $T_{\max} = 0.596$   
12668 measured reflections

3314 independent reflections  
3010 reflections with  $I > 2\sigma(I)$   
 $R_{\text{int}} = 0.049$   
 $\theta_{\max} = 27.5^\circ$ ,  $\theta_{\min} = 2.6^\circ$   
 $h = -9 \rightarrow 9$   
 $k = -10 \rightarrow 10$   
 $l = -30 \rightarrow 30$

## Refinement

Refinement on  $F^2$   
Least-squares matrix: full  
 $R[F^2 > 2\sigma(F^2)] = 0.029$   
 $wR(F^2) = 0.073$   
 $S = 1.08$   
3314 reflections  
192 parameters  
0 restraints

Primary atom site location: structure-invariant direct methods  
Secondary atom site location: difference Fourier map  
Hydrogen site location: inferred from neighbouring sites  
H-atom parameters constrained  
 $w = 1/[\sigma^2(F_o^2) + (0.035P)^2 + 1.3471P]$   
where  $P = (F_o^2 + 2F_c^2)/3$   
 $(\Delta/\sigma)_{\max} = 0.001$   
 $\Delta\rho_{\max} = 1.21 \text{ e \AA}^{-3}$   
 $\Delta\rho_{\min} = -1.57 \text{ e \AA}^{-3}$

## Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters ( $\text{\AA}^2$ )

	x	y	z	$U_{\text{iso}}^*/U_{\text{eq}}$
Pb1	0.72555 (2)	0.46532 (2)	0.554759 (7)	0.01208 (8)
O1	0.4023 (5)	0.4525 (4)	0.54078 (14)	0.0149 (7)
O2	0.1288 (5)	0.3557 (5)	0.55875 (15)	0.0225 (8)
O3	0.6357 (5)	0.7354 (5)	0.56638 (14)	0.0225 (8)
O4	0.7982 (5)	0.6708 (4)	0.64405 (14)	0.0215 (8)
O5	0.6307 (5)	0.1781 (4)	0.51184 (13)	0.0206 (8)
H5	0.5230	0.1444	0.5119	0.031*
N1	0.5510 (5)	0.3454 (5)	0.63884 (16)	0.0134 (8)
C1	0.2921 (7)	0.3768 (6)	0.57117 (19)	0.0144 (10)
C2	0.3771 (7)	0.3126 (6)	0.62685 (19)	0.0145 (10)
C3	0.2694 (7)	0.2289 (6)	0.66278 (19)	0.0168 (10)
H3	0.1453	0.2066	0.6518	0.020*

C4	0.3433 (7)	0.1795 (7)	0.7138 (2)	0.0199 (11)
H4	0.2712	0.1234	0.7390	0.024*
C5	0.5280 (7)	0.2127 (6)	0.72855 (19)	0.0162 (10)
C6	0.6171 (7)	0.1640 (7)	0.7808 (2)	0.0211 (11)
H6	0.5501	0.1091	0.8076	0.025*
C7	0.7974 (7)	0.1958 (6)	0.7925 (2)	0.0206 (11)
H7	0.8559	0.1611	0.8271	0.025*
C8	0.8983 (7)	0.2801 (6)	0.7536 (2)	0.0208 (11)
H8	1.0239	0.3017	0.7624	0.025*
C9	0.8177 (7)	0.3308 (6)	0.70336 (19)	0.0168 (10)
H9	0.8862	0.3887	0.6777	0.020*
C10	0.6310 (7)	0.2963 (6)	0.68997 (19)	0.0155 (10)
C11	0.7125 (7)	0.7694 (6)	0.61446 (19)	0.0155 (10)
C12	0.6856 (8)	0.9361 (6)	0.6361 (2)	0.0225 (12)
H12A	0.8021	0.9774	0.6524	0.034*
H12B	0.5964	0.9343	0.6650	0.034*
H12C	0.6406	1.0049	0.6050	0.034*
C13	0.7704 (9)	0.0864 (8)	0.4891 (2)	0.0311 (14)
H13A	0.7238	-0.0224	0.4810	0.037*
H13B	0.8746	0.0775	0.5175	0.037*
C14	0.8359 (8)	0.1573 (8)	0.4361 (2)	0.0315 (14)
H14A	0.9244	0.0853	0.4204	0.047*
H14B	0.8941	0.2602	0.4448	0.047*
H14C	0.7319	0.1727	0.4087	0.047*

Atomic displacement parameters ( $\text{\AA}^2$ )

	$U^{11}$	$U^{22}$	$U^{33}$	$U^{12}$	$U^{13}$	$U^{23}$
Pb1	0.00993 (11)	0.01418 (12)	0.01215 (12)	0.00129 (7)	0.00073 (7)	0.00054 (6)
O1	0.0135 (18)	0.0172 (19)	0.0138 (17)	0.0006 (14)	-0.0009 (14)	0.0015 (13)
O2	0.0117 (18)	0.032 (2)	0.0239 (18)	0.0019 (16)	-0.0002 (14)	0.0063 (16)
O3	0.024 (2)	0.022 (2)	0.0209 (18)	0.0037 (17)	-0.0060 (15)	-0.0039 (16)
O4	0.028 (2)	0.0179 (19)	0.0179 (17)	0.0039 (17)	-0.0045 (15)	-0.0008 (15)
O5	0.0188 (19)	0.0208 (19)	0.0218 (18)	0.0037 (16)	-0.0012 (14)	-0.0045 (15)
N1	0.013 (2)	0.014 (2)	0.0129 (19)	0.0040 (17)	-0.0012 (15)	-0.0020 (16)
C1	0.012 (2)	0.016 (3)	0.015 (2)	0.004 (2)	0.0023 (18)	-0.0019 (19)
C2	0.016 (2)	0.013 (2)	0.015 (2)	0.004 (2)	0.0026 (18)	-0.0019 (19)
C3	0.015 (2)	0.016 (3)	0.020 (2)	-0.001 (2)	0.0040 (19)	0.002 (2)
C4	0.020 (3)	0.020 (3)	0.021 (2)	0.000 (2)	0.008 (2)	0.005 (2)
C5	0.018 (3)	0.017 (3)	0.014 (2)	0.006 (2)	0.0017 (19)	-0.0030 (19)
C6	0.028 (3)	0.024 (3)	0.012 (2)	0.002 (2)	0.005 (2)	0.003 (2)
C7	0.026 (3)	0.021 (3)	0.014 (2)	0.006 (2)	-0.002 (2)	0.004 (2)
C8	0.018 (3)	0.025 (3)	0.019 (2)	0.007 (2)	-0.003 (2)	-0.004 (2)
C9	0.018 (3)	0.018 (3)	0.015 (2)	-0.002 (2)	0.0029 (19)	0.000 (2)
C10	0.019 (3)	0.011 (2)	0.017 (2)	0.005 (2)	0.0023 (19)	-0.0042 (19)
C11	0.014 (2)	0.013 (2)	0.020 (2)	-0.004 (2)	0.0050 (19)	0.000 (2)
C12	0.032 (3)	0.012 (3)	0.023 (3)	0.003 (2)	-0.002 (2)	-0.003 (2)
C13	0.030 (3)	0.031 (3)	0.034 (3)	0.019 (3)	0.007 (3)	0.003 (3)

## supplementary materials

---

C14                    0.024 (3)                    0.039 (4)                    0.032 (3)                    0.009 (3)                    0.008 (2)                    0.003 (3)

### *Geometric parameters (Å, °)*

Pb1—O1	2.377 (4)	C4—H4	0.9500
Pb1—O3	2.384 (4)	C5—C10	1.414 (7)
Pb1—O1 <sup>i</sup>	2.500 (3)	C5—C6	1.427 (7)
Pb1—N1	2.645 (4)	C6—C7	1.362 (8)
Pb1—O5	2.694 (4)	C6—H6	0.9500
Pb1—O4	2.763 (3)	C7—C8	1.413 (7)
Pb1—O2 <sup>ii</sup>	3.096 (4)	C7—H7	0.9500
O1—C1	1.288 (6)	C8—C9	1.367 (7)
O1—Pb1 <sup>i</sup>	2.500 (3)	C8—H8	0.9500
O2—C1	1.229 (6)	C9—C10	1.417 (7)
O3—C11	1.275 (6)	C9—H9	0.9500
O4—C11	1.233 (6)	C11—C12	1.510 (7)
O5—C13	1.418 (6)	C12—H12A	0.9800
O5—H5	0.8400	C12—H12B	0.9800
N1—C2	1.319 (6)	C12—H12C	0.9800
N1—C10	1.380 (6)	C13—C14	1.502 (8)
C1—C2	1.526 (7)	C13—H13A	0.9900
C2—C3	1.392 (7)	C13—H13B	0.9900
C3—C4	1.362 (7)	C14—H14A	0.9800
C3—H3	0.9500	C14—H14B	0.9800
C4—C5	1.407 (7)	C14—H14C	0.9800
O1—Pb1—O3	77.18 (12)	C3—C4—H4	120.5
O1—Pb1—O1 <sup>i</sup>	64.70 (14)	C5—C4—H4	120.5
O3—Pb1—O1 <sup>i</sup>	75.67 (11)	C4—C5—C10	119.0 (4)
O1—Pb1—N1	63.92 (12)	C4—C5—C6	122.7 (5)
O3—Pb1—N1	97.09 (12)	C10—C5—C6	118.3 (5)
O1 <sup>i</sup> —Pb1—N1	128.43 (12)	C7—C6—C5	120.4 (5)
O1—Pb1—O5	71.05 (11)	C7—C6—H6	119.8
O3—Pb1—O5	146.07 (12)	C5—C6—H6	119.8
O1 <sup>i</sup> —Pb1—O5	80.20 (11)	C6—C7—C8	120.5 (5)
N1—Pb1—O5	79.62 (11)	C6—C7—H7	119.7
O1—Pb1—O4	106.08 (11)	C8—C7—H7	119.7
O3—Pb1—O4	50.00 (11)	C9—C8—C7	120.9 (5)
O1 <sup>i</sup> —Pb1—O4	124.72 (10)	C9—C8—H8	119.5
N1—Pb1—O4	74.61 (11)	C7—C8—H8	119.5
O5—Pb1—O4	152.06 (10)	C8—C9—C10	119.4 (5)
O1—Pb1—O2 <sup>ii</sup>	159.18 (11)	C8—C9—H9	120.3
O3—Pb1—O2 <sup>ii</sup>	123.42 (12)	C10—C9—H9	120.3
O1 <sup>i</sup> —Pb1—O2 <sup>ii</sup>	114.32 (10)	N1—C10—C5	120.4 (4)
N1—Pb1—O2 <sup>ii</sup>	111.89 (10)	N1—C10—C9	119.2 (4)
O5—Pb1—O2 <sup>ii</sup>	88.18 (11)	C5—C10—C9	120.4 (4)
O4—Pb1—O2 <sup>ii</sup>	91.48 (10)	O4—C11—O3	122.8 (5)

C1—O1—Pb1	127.0 (3)	O4—C11—C12	120.1 (4)
C1—O1—Pb1 <sup>i</sup>	115.8 (3)	O3—C11—C12	117.0 (4)
Pb1—O1—Pb1 <sup>i</sup>	115.30 (14)	C11—C12—H12A	109.5
C11—O3—Pb1	102.0 (3)	C11—C12—H12B	109.5
C11—O4—Pb1	85.1 (3)	H12A—C12—H12B	109.5
C13—O5—Pb1	117.1 (4)	C11—C12—H12C	109.5
C13—O5—H5	121.4	H12A—C12—H12C	109.5
Pb1—O5—H5	121.4	H12B—C12—H12C	109.5
C2—N1—C10	118.6 (4)	O5—C13—C14	112.5 (5)
C2—N1—Pb1	115.2 (3)	O5—C13—H13A	109.1
C10—N1—Pb1	125.6 (3)	C14—C13—H13A	109.1
O2—C1—O1	125.1 (4)	O5—C13—H13B	109.1
O2—C1—C2	119.7 (4)	C14—C13—H13B	109.1
O1—C1—C2	115.2 (4)	H13A—C13—H13B	107.8
N1—C2—C3	123.6 (4)	C13—C14—H14A	109.5
N1—C2—C1	116.7 (4)	C13—C14—H14B	109.5
C3—C2—C1	119.7 (4)	H14A—C14—H14B	109.5
C4—C3—C2	119.5 (5)	C13—C14—H14C	109.5
C4—C3—H3	120.3	H14A—C14—H14C	109.5
C2—C3—H3	120.3	H14B—C14—H14C	109.5
C3—C4—C5	118.9 (5)		
O3—Pb1—O1—C1	-116.7 (4)	O4—Pb1—N1—C10	-60.4 (4)
O1 <sup>i</sup> —Pb1—O1—C1	163.3 (4)	O2 <sup>ii</sup> —Pb1—N1—C10	24.9 (4)
N1—Pb1—O1—C1	-12.1 (3)	Pb1—O1—C1—O2	-170.1 (4)
O5—Pb1—O1—C1	75.4 (4)	Pb1 <sup>i</sup> —O1—C1—O2	-6.8 (6)
O4—Pb1—O1—C1	-75.4 (4)	Pb1—O1—C1—C2	10.8 (6)
O2 <sup>ii</sup> —Pb1—O1—C1	71.1 (5)	Pb1 <sup>i</sup> —O1—C1—C2	174.0 (3)
O3—Pb1—O1—Pb1 <sup>i</sup>	80.02 (15)	C10—N1—C2—C3	-0.8 (7)
O1 <sup>i</sup> —Pb1—O1—Pb1 <sup>i</sup>	0.0	Pb1—N1—C2—C3	170.7 (4)
N1—Pb1—O1—Pb1 <sup>i</sup>	-175.37 (19)	C10—N1—C2—C1	176.9 (4)
O5—Pb1—O1—Pb1 <sup>i</sup>	-87.89 (15)	Pb1—N1—C2—C1	-11.6 (5)
O4—Pb1—O1—Pb1 <sup>i</sup>	121.29 (13)	O2—C1—C2—N1	-177.0 (4)
O2 <sup>ii</sup> —Pb1—O1—Pb1 <sup>i</sup>	-92.2 (3)	O1—C1—C2—N1	2.2 (6)
O1—Pb1—O3—C11	126.0 (3)	O2—C1—C2—C3	0.8 (7)
O1 <sup>i</sup> —Pb1—O3—C11	-167.2 (3)	O1—C1—C2—C3	180.0 (4)
N1—Pb1—O3—C11	64.9 (3)	N1—C2—C3—C4	1.2 (8)
O5—Pb1—O3—C11	146.8 (3)	C1—C2—C3—C4	-176.4 (5)
O4—Pb1—O3—C11	1.8 (3)	C2—C3—C4—C5	-0.8 (7)
O2 <sup>ii</sup> —Pb1—O3—C11	-57.3 (3)	C3—C4—C5—C10	0.0 (7)
O1—Pb1—O4—C11	-59.0 (3)	C3—C4—C5—C6	-179.4 (5)
O3—Pb1—O4—C11	-1.9 (3)	C4—C5—C6—C7	178.4 (5)
O1 <sup>i</sup> —Pb1—O4—C11	11.1 (3)	C10—C5—C6—C7	-0.9 (8)
N1—Pb1—O4—C11	-115.3 (3)	C5—C6—C7—C8	1.2 (8)
O5—Pb1—O4—C11	-138.7 (3)	C6—C7—C8—C9	-0.3 (8)
O2 <sup>ii</sup> —Pb1—O4—C11	132.4 (3)	C7—C8—C9—C10	-0.9 (8)
O1—Pb1—O5—C13	164.3 (3)	C2—N1—C10—C5	-0.1 (7)

## supplementary materials

---

O3—Pb1—O5—C13	142.8 (3)	Pb1—N1—C10—C5	-170.6 (3)
O1 <sup>i</sup> —Pb1—O5—C13	97.8 (3)	C2—N1—C10—C9	180.0 (4)
N1—Pb1—O5—C13	-129.9 (3)	Pb1—N1—C10—C9	9.4 (6)
O4—Pb1—O5—C13	-107.0 (4)	C4—C5—C10—N1	0.4 (7)
O2 <sup>ii</sup> —Pb1—O5—C13	-17.3 (3)	C6—C5—C10—N1	179.8 (4)
O1—Pb1—N1—C2	11.7 (3)	C4—C5—C10—C9	-179.6 (5)
O3—Pb1—N1—C2	83.6 (3)	C6—C5—C10—C9	-0.2 (7)
O1 <sup>i</sup> —Pb1—N1—C2	6.3 (4)	C8—C9—C10—N1	-178.9 (5)
O5—Pb1—N1—C2	-62.2 (3)	C8—C9—C10—C5	1.1 (7)
O4—Pb1—N1—C2	128.7 (3)	Pb1—O4—C11—O3	3.2 (5)
O2 <sup>ii</sup> —Pb1—N1—C2	-146.0 (3)	Pb1—O4—C11—C12	179.6 (4)
O1—Pb1—N1—C10	-177.5 (4)	Pb1—O3—C11—O4	-3.8 (5)
O3—Pb1—N1—C10	-105.5 (4)	Pb1—O3—C11—C12	179.7 (4)
O1 <sup>i</sup> —Pb1—N1—C10	177.2 (3)	Pb1—O5—C13—C14	-68.5 (6)
O5—Pb1—N1—C10	108.7 (4)		

Symmetry codes: (i)  $-x+1, -y+1, -z+1$ ; (ii)  $x+1, y, z$ .

### Hydrogen-bond geometry ( $\text{\AA}$ , $^\circ$ )

$D-H\cdots A$	$D-H$	$H\cdots A$	$D\cdots A$	$D-H\cdots A$
O5—H5 $\cdots$ O3 <sup>i</sup>	0.84	2.36	2.710 (5)	106

Symmetry codes: (i)  $-x+1, -y+1, -z+1$ .



Fig. 1

