

## Bis(2-amino-6-methylpyridinium) *trans*-diaquabis(pyrazine-2,3-dicarboxylato)-cobaltate(II) octahydrate

Hossein Eshtiagh-Hosseini,<sup>a</sup> Nafiseh Alfi,<sup>a</sup> Masoud Mirzaei<sup>a</sup> and Philip E. Fanwick<sup>b\*</sup>

<sup>a</sup>Department of Chemistry, School of Sciences, Ferdowsi University of Mashhad, Mashhad, Iran, and <sup>b</sup>Department of Chemistry, Purdue University, W. Lafayette, IN 47907, USA

Correspondence e-mail: pfanwick@purdue.edu


Received 20 December 2010; accepted 8 January 2011

Key indicators: single-crystal X-ray study;  $T = 150$  K; mean  $\sigma(\text{C}-\text{C}) = 0.003$  Å;  $R$  factor = 0.030;  $wR$  factor = 0.087; data-to-parameter ratio = 11.1.

The title compound,  $(\text{C}_6\text{H}_9\text{N}_2)_2[\text{Co}(\text{C}_6\text{H}_2\text{N}_2\text{O}_4)_2(\text{H}_2\text{O})_2] \cdot 8\text{H}_2\text{O}$ , was obtained by the reaction of  $\text{CoCl}_2 \cdot 6\text{H}_2\text{O}$  with 1,4-pyrazine-2,3-dicarboxylic acid and 2-amino-6-methylpyridine in aqueous solution (molar ratio 1:2:2). The  $\text{Co}^{\text{II}}$  ion is situated on an inversion centre and is coordinated by two O and two N atoms of two symmetry-related 1,4-pyrazine-2,3-dicarboxylate ligands and two water molecules and has a disorted octahedral coordination environment. The asymmetric unit also contains four water molecules. In the crystal, extensive intermolecular classical  $\text{N}-\text{H} \cdots \text{O}$ ,  $\text{O}-\text{H} \cdots \text{O}$  and  $\text{O}-\text{H} \cdots \text{N}$  hydrogen bonds and  $\pi-\pi$  stacking interactions [centroid-centroid distance =  $3.490(1)$  Å] connect the various components, forming a three-dimensional network.

### Related literature

For related structures based on 1,4-pyrazine-2,3-dicarboxylate ligands, see: Eshtiagh-Hosseini, Alfi *et al.* (2010). Eshtiagh-Hosseini, Gschwind *et al.* (2010). Eshtiagh-Hosseini, Necas *et al.* (2010).



### Experimental

#### Crystal data

$(\text{C}_6\text{H}_9\text{N}_2)_2[\text{Co}(\text{C}_6\text{H}_2\text{N}_2\text{O}_4)_2(\text{H}_2\text{O})_2] \cdot 8\text{H}_2\text{O}$   
 $M_r = 789.58$   
 Triclinic,  $P\bar{1}$   
 $a = 6.8570(4)$  Å  
 $b = 10.2348(5)$  Å  
 $c = 13.6403(10)$  Å  
 $\alpha = 109.604(4)^\circ$

$\beta = 90.424(5)^\circ$   
 $\gamma = 105.524(4)^\circ$   
 $V = 863.89(9)$  Å<sup>3</sup>  
 $Z = 1$   
 Cu  $K\alpha$  radiation  
 $\mu = 4.68$  mm<sup>-1</sup>  
 $T = 150$  K  
 $0.20 \times 0.18 \times 0.14$  mm

#### Data collection

Rigaku RAPID II diffractometer  
 Absorption correction: multi-scan (SCALEPACK; Otwinowski & Minor, 1997)  
 $T_{\text{min}} = 0.280$ ,  $T_{\text{max}} = 0.508$

19137 measured reflections  
 3152 independent reflections  
 3151 reflections with  $> 2.0\sigma(I)$   
 $R_{\text{int}} = 0.036$

#### Refinement

$R[F^2 > 2\sigma(F^2)] = 0.030$   
 $wR(F^2) = 0.087$   
 $S = 1.04$   
 3152 reflections  
 285 parameters

H atoms treated by a mixture of independent and constrained refinement  
 $\Delta\rho_{\text{max}} = 0.23$  e Å<sup>-3</sup>  
 $\Delta\rho_{\text{min}} = -0.40$  e Å<sup>-3</sup>

**Table 1**

Hydrogen-bond geometry (Å, °).

$D-\text{H} \cdots A$	$D-\text{H}$	$\text{H} \cdots A$	$D \cdots A$	$D-\text{H} \cdots A$
$\text{N11}-\text{H11} \cdots \text{O31}$	0.82 (2)	1.98 (2)	2.794 (3)	179 (2)
$\text{N12}-\text{H121} \cdots \text{O2W}^{\text{i}}$	0.86 (3)	2.05 (2)	2.900 (2)	170 (2)
$\text{N12}-\text{H122} \cdots \text{O32}$	0.84 (3)	1.97 (3)	2.804 (3)	176 (2)
$\text{O1W}-\text{H1W1} \cdots \text{O2W}^{\text{ii}}$	0.78 (3)	1.99 (3)	2.770 (3)	178 (3)
$\text{O1W}-\text{H1W2} \cdots \text{O5W}^{\text{iii}}$	0.85 (3)	1.85 (3)	2.697 (2)	172 (3)
$\text{O2W}-\text{H2W1} \cdots \text{O21}$	0.85 (3)	2.10 (3)	2.942 (3)	168 (3)
$\text{O2W}-\text{H2W2} \cdots \text{O4W}$	0.72 (3)	2.07 (3)	2.784 (3)	176 (2)
$\text{O3W}-\text{H3W2} \cdots \text{O4W}$	0.83 (3)	1.99 (3)	2.811 (3)	167 (3)
$\text{O4W}-\text{H4W1} \cdots \text{O3W}^{\text{iii}}$	0.80 (3)	1.98 (2)	2.755 (2)	165 (3)
$\text{O4W}-\text{H4W2} \cdots \text{O31}^{\text{iv}}$	0.80 (2)	1.94 (2)	2.738 (2)	178 (3)
$\text{O5W}-\text{H5W1} \cdots \text{O32}^{\text{v}}$	0.79 (2)	2.00 (3)	2.767 (2)	161 (3)
$\text{O5W}-\text{H5W2} \cdots \text{N4}^{\text{vi}}$	0.77 (3)	2.11 (3)	2.871 (3)	167 (3)

Symmetry codes: (i)  $x, y-1, z$ ; (ii)  $-x+1, -y+1, -z$ ; (iii)  $-x+1, -y+1, -z+1$ ; (iv)  $x-1, y, z$ ; (v)  $x, y+1, z$ ; (vi)  $x-1, y+1, z$ .

Data collection: *CrystalClear* (Molecular Structure Corporation & Rigaku, 2001); cell refinement: *DENZO/SCALEPACK* (Otwinowski & Minor, 1997); data reduction: *DENZO/SCALEPACK*; program(s) used to solve structure: *SIR2004* (Burla *et al.*, 2005); program(s) used to refine structure: *SHELXL97* (Sheldrick, 2008); molecular graphics: *ORTEP* (Johnson, 1976) and *PLATON* (Spek, 2009); software used to prepare material for publication: *SHELXL97* and local programs.

The authors express their appreciation to the Ferdowsi University of Mashhad for financial support of this research (grant No. P/355).

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: BT5444).

### References

- Burla, M. C., Caliendo, R., Camalli, M., Carrozzini, B., Cascarano, G. L., De Caro, L., Giacovazzo, C., Polidori, G. & Spagna, R. (2005). *J. Appl. Cryst.* **38**, 381–388.  
 Eshtiagh-Hosseini, H., Alfi, N., Mirzaei, M. & Necas, M. (2010). *Acta Cryst.* **E66**, o2810–o2811.

- Eshtiagh-Hosseini, H., Gschwind, F., Alfi, N. & Mirzaei, M. (2010). *Acta Cryst. E66*, m826–m827.
- Eshtiagh-Hosseini, H., Necas, M., Alfi, N. & Mirzaei, M. (2010). *Acta Cryst. E66*, m1320–m1321.
- Johnson, C. K. (1976). *ORTEP*. Report ORNL-5138. Oak Ridge National Laboratory, Tennessee, USA.
- Molecular Structure Corporation & Rigaku (2001). *CrystalClear*. MSC, The Woodlands, Texas, USA, and Rigaku Corporation, Tokyo, Japan.
- Otwinowski, Z. & Minor, W. (1997). *Methods in Enzymology*, Vol. 276, *Macromolecular Crystallography*, Part A, edited by C. W. Carter Jr & R. M. Sweet, pp. 307–326. New York: Academic Press.
- Sheldrick, G. M. (2008). *Acta Cryst. A64*, 112–122.
- Spek, A. L. (2009). *Acta Cryst. D65*, 148–155.

**supplementary materials**

*Acta Cryst.* (2011). E67, m266-m267 [ doi:10.1107/S1600536811001127 ]

## Bis(2-amino-6-methylpyridinium) *trans*-diaquabis(pyrazine-2,3-dicarboxylato)cobaltate(II) octahydrate

H. Eshtiagh-Hosseini, N. Alfi, M. Mirzaei and P. E. Fanwick

### Comment

In the recent years, our research group has been interested in the synthesis of proton transfer compounds and study of their behavior with metal ions. We have focused on the proton delivery from polycarboxylic acids, which are considered as very good donors and amines as acceptors. Among polycarboxylic acids, 1,4-pyrazine-2,3-dicarboxylic acid (pyzdcH<sub>2</sub>) as a very important carboxylate derivative has attracted much interest in coordination chemistry and it is the one that we utilized widely in our studies (Eshtiagh-Hosseini, Alfi *et al.*, 2010). In order to develop novel systems, we wish to report the first complex of Co<sup>II</sup> ion with pyzdcH<sub>2</sub> as proton donor and 2a-6 m as proton acceptor. PyzdcH<sub>2</sub> has proved to be well suited for the construction of multidimensional frameworks due to the presence of two adjacent carboxylate groups (O donor atoms) as substituents on the N-heterocyclic pyrazine ring (N donor atoms).

The asymmetric unit of the title compound (Fig. 1), contains half a [Co(pyzdc)<sub>2</sub>(H<sub>2</sub>O)<sub>2</sub>]<sup>2-</sup> anion, a (2a-6mpyH)<sup>+</sup> cation, and four uncoordinated water molecules. In the anions, Co<sup>II</sup> ion has a N<sub>2</sub>O<sub>4</sub> donor set bond with normal distances and angles. The title compound can be compared with the mono-nuclear coordination compound of Co<sup>II</sup> ion which has recently been synthesized and characterized by our research group (Eshtiagh-Hosseini, Necas *et al.*, 2010). There are some hydrogen bonding interactions such as O–H...O and N–H...O between cations, anions and uncoordinated water molecules (Table 2). The water molecules act also as bridging agents and link anions and cationic fragments together *via* hydrogen bonds which resulted in the creation of six supramolecular synthons as R<sup>2</sup><sub>2</sub>(8), R<sup>3</sup><sub>4</sub>(10), R<sup>3</sup><sub>5</sub>(10), R<sup>4</sup><sub>5</sub>(15), R<sup>4</sup><sub>4</sub>(18), R<sup>4</sup><sub>4</sub>(26) (Figs. 2, 3). As it is seen in Fig. 4, there are also  $\pi$ - $\pi$  stacking interactions between the aromatic rings of the coordinated (pyzdc)<sup>2-</sup> and carboxylate functional group anions and (2a-6mpyH)<sup>+</sup> cation. Ion pairing, hydrogen bonds,  $\pi$ - $\pi$  stacking, and van der Waals interactions stabilize the crystal structure. These interactions lead to formation of a three-dimensional structure. By the help of hydrogen bond interactions between uncoordinated water molecules, the related crystalline network bears (H<sub>2</sub>O)<sub>6</sub> cluster in the form of two branched-cyclohexane (Eshtiagh-Hosseini, Gschwind *et al.*, 2010).

### Experimental

A solution of pyzdcH<sub>2</sub> (0.6 mmol, 0.1 g) and 2a-6mpy (1.2 mmol, 0.13 g) in water (10 ml) was refluxed for an hour, then a solution of CoCl<sub>2</sub>.6H<sub>2</sub>O (0.02 mmol, 0.05 g) was added dropwise and continued refluxing for 6 hrs at 293 K. The obtained orange solution gave orange block like crystals of title compound after slow evaporation of solvent at R.T.

### Refinement

Carbon bound hydrogen atoms were positioned geometrically and refined as riding using standard *SHELXTL* constraints, with their  $U_{\text{iso}}$  set to either 1.2 $U_{\text{eq}}(\text{C})$  or 1.5 $U_{\text{eq}}(\text{C}_{\text{methyl}})$  of their parent atoms. The C–H distances were set to 0.93 and

## supplementary materials

0.96 Å for aromatic and methyl H atoms, respectively. Hydrogen atoms bonded to N and O were located in a difference Fourier map and refined isotropically.

### Figures




Fig. 1. An *ORTEP* drawing of the title compound showing 50% ellipsoid probability. Only the symmetry independent atoms are labelled.




Fig. 2. Water molecules connecting anions and cations.




Fig. 3. Schematic representation of different graph set motifs in the crystalline network of **1**. Dashed lines indicate the hydrogen bonds.



Fig. 4. Perspective views of the  $\pi$ - $\pi$  stacking interactions.

### Bis(2-amino-6-methylpyridinium) *trans*-diaquabis(pyrazine-2,3-dicarboxylato)cobaltate(II) octahydrate

#### Crystal data

(C<sub>6</sub>H<sub>9</sub>N<sub>2</sub>)<sub>2</sub>[Co(C<sub>6</sub>H<sub>2</sub>N<sub>2</sub>O<sub>4</sub>)<sub>2</sub>(H<sub>2</sub>O)<sub>2</sub>]·8H<sub>2</sub>O

$M_r = 789.58$

Triclinic,  $P\bar{1}$

Hall symbol: -P 1

$a = 6.8570$  (4) Å

$b = 10.2348$  (5) Å

$c = 13.6403$  (10) Å

$\alpha = 109.604$  (4)°

$\beta = 90.424$  (5)°

$\gamma = 105.524$  (4)°

$V = 863.89$  (9) Å<sup>3</sup>

$Z = 1$

$F(000) = 413$

$D_x = 1.518$  Mg m<sup>-3</sup>

Cu -  $K\alpha$  radiation,  $\lambda = 1.54184$  Å

Cell parameters from 3181 reflections

$\theta = 3$ –71°

$\mu = 4.68$  mm<sup>-1</sup>

$T = 150$  K

Chunk, brown

$0.20 \times 0.18 \times 0.14$  mm

#### Data collection

Rigaku RAPID II  
diffractometer

confocal optics

$\omega$  scans

Absorption correction: multi-scan  
(*SCALEPACK*; Otwinowski & Minor, 1997)

3151 reflections with  $> 2.0\sigma(I)$

$R_{int} = 0.036$

$\theta_{max} = 71.9^\circ$ ,  $\theta_{min} = 3.5^\circ$

$h = 0 \rightarrow 8$

$T_{\min} = 0.280$ ,  $T_{\max} = 0.508$   
 19137 measured reflections  
 3152 independent reflections

$k = -12 \rightarrow 12$   
 $l = -16 \rightarrow 16$

### Refinement

Refinement on $F^2$	0 restraints
Least-squares matrix: full	H atoms treated by a mixture of independent and constrained refinement
$R[F^2 > 2\sigma(F^2)] = 0.030$	$w = 1/[\sigma^2(F_o^2) + (0.057P)^2 + 0.2735P]$
$wR(F^2) = 0.087$	where $P = (F_o^2 + 2F_c^2)/3$
$S = 1.04$	$(\Delta/\sigma)_{\max} = 0.001$
3152 reflections	$\Delta\rho_{\max} = 0.23 \text{ e } \text{\AA}^{-3}$
285 parameters	$\Delta\rho_{\min} = -0.40 \text{ e } \text{\AA}^{-3}$

### Special details

**Geometry.** All e.s.d.'s (except the e.s.d. in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell e.s.d.'s are taken into account individually in the estimation of e.s.d.'s in distances, angles and torsion angles; correlations between e.s.d.'s in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell e.s.d.'s is used for estimating e.s.d.'s involving l.s. planes.

**Refinement.** Refinement on  $F^2$  for ALL reflections except for 0 with very negative  $F^2$  or flagged by the user for potential systematic errors. Weighted  $R$ -factors  $wR$  and all goodnesses of fit  $S$  are based on  $F^2$ , conventional  $R$ -factors  $R$  are based on  $F$ , with  $F$  set to zero for negative  $F^2$ . The observed criterion of  $F^2 > \sigma(F^2)$  is used only for calculating  $R_{\text{factor\_obs}}$  etc. and is not relevant to the choice of reflections for refinement.  $R$ -factors based on  $F^2$  are statistically about twice as large as those based on  $F$ , and  $R$ -factors based on ALL data will be even larger.

### Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters ( $\text{\AA}^2$ )

	$x$	$y$	$z$	$U_{\text{iso}}^*/U_{\text{eq}}$
Co1	1.0000	0.5000	0.0000	0.01861 (12)
O1W	0.78751 (19)	0.34893 (13)	-0.12293 (10)	0.0281 (3)
O21	0.79587 (17)	0.45953 (12)	0.10552 (10)	0.0244 (3)
O22	0.70494 (18)	0.32395 (14)	0.20508 (10)	0.0309 (3)
O2W	0.4756 (2)	0.56148 (13)	0.22356 (12)	0.0277 (3)
O31	0.97825 (17)	0.20966 (12)	0.31826 (9)	0.0257 (3)
O32	0.77764 (18)	0.02378 (12)	0.18707 (10)	0.0278 (3)
O3W	0.6610 (2)	0.37101 (15)	0.42401 (11)	0.0318 (3)
O4W	0.3166 (2)	0.44434 (16)	0.37295 (12)	0.0307 (3)
O5W	0.3645 (2)	0.92255 (14)	0.12233 (11)	0.0301 (3)
N1	1.08935 (19)	0.33770 (13)	0.03478 (10)	0.0185 (3)
N4	1.1850 (2)	0.14113 (14)	0.11111 (11)	0.0212 (3)
N11	0.82673 (19)	0.07875 (14)	0.46263 (11)	0.0201 (3)
N12	0.6818 (2)	-0.13230 (16)	0.32292 (12)	0.0243 (3)
C2	0.9819 (2)	0.29538 (15)	0.10633 (12)	0.0179 (3)

## supplementary materials

---

C3	1.0303 (2)	0.19613 (16)	0.14422 (13)	0.0188 (3)
C5	1.2920 (2)	0.18647 (17)	0.04140 (13)	0.0219 (3)
C6	1.2436 (2)	0.28428 (17)	0.00249 (13)	0.0213 (3)
C12	0.7163 (2)	-0.06269 (17)	0.42549 (13)	0.0200 (3)
C13	0.6442 (2)	-0.12847 (18)	0.49910 (14)	0.0240 (4)
C14	0.6901 (3)	-0.0497 (2)	0.60314 (15)	0.0286 (4)
C15	0.8071 (3)	0.0960 (2)	0.63821 (14)	0.0281 (4)
C16	0.8729 (2)	0.15970 (18)	0.56645 (14)	0.0239 (3)
C17	0.9948 (3)	0.31426 (18)	0.59367 (16)	0.0307 (4)
C21	0.8125 (2)	0.36383 (17)	0.14298 (13)	0.0215 (3)
C31	0.9158 (2)	0.13968 (17)	0.22335 (13)	0.0211 (3)
H5	1.4018	0.1516	0.0183	0.026*
H6	1.3200	0.3128	-0.0467	0.026*
H11	0.869 (3)	0.117 (2)	0.4202 (17)	0.023 (5)*
H13	0.5660	-0.2248	0.4768	0.029*
H14	0.6432	-0.0930	0.6518	0.034*
H15	0.8394	0.1485	0.7095	0.034*
H121	0.618 (3)	-0.223 (3)	0.3014 (17)	0.032 (5)*
H122	0.716 (3)	-0.086 (3)	0.2827 (19)	0.035 (6)*
H17A	1.1225	0.3185	0.5646	0.046*
H17B	1.0194	0.3600	0.6684	0.046*
H17C	0.9210	0.3636	0.5655	0.046*
H1W1	0.712 (4)	0.374 (3)	-0.150 (2)	0.044 (7)*
H1W2	0.729 (4)	0.263 (3)	-0.1249 (19)	0.045 (7)*
H2W1	0.561 (4)	0.520 (3)	0.191 (2)	0.052 (7)*
H2W2	0.431 (4)	0.528 (3)	0.260 (2)	0.037 (7)*
H3W1	0.728 (4)	0.345 (3)	0.378 (2)	0.046 (7)*
H3W2	0.564 (4)	0.389 (3)	0.399 (2)	0.050 (7)*
H4W1	0.300 (4)	0.493 (3)	0.430 (2)	0.039 (7)*
H4W2	0.219 (4)	0.375 (3)	0.356 (2)	0.051 (7)*
H5W1	0.478 (4)	0.970 (3)	0.145 (2)	0.049 (7)*
H5W2	0.304 (4)	0.977 (3)	0.125 (2)	0.042 (7)*

### Atomic displacement parameters ( $\text{\AA}^2$ )

	$U^{11}$	$U^{22}$	$U^{33}$	$U^{12}$	$U^{13}$	$U^{23}$
Co1	0.02261 (19)	0.01812 (18)	0.0174 (2)	0.00806 (13)	0.00137 (13)	0.00745 (14)
O1W	0.0329 (6)	0.0208 (6)	0.0290 (8)	0.0054 (5)	-0.0089 (5)	0.0086 (5)
O21	0.0262 (6)	0.0268 (6)	0.0269 (7)	0.0142 (5)	0.0064 (5)	0.0126 (5)
O22	0.0295 (6)	0.0455 (7)	0.0302 (8)	0.0197 (5)	0.0137 (5)	0.0219 (6)
O2W	0.0279 (6)	0.0239 (6)	0.0322 (8)	0.0108 (5)	0.0006 (5)	0.0083 (6)
O31	0.0292 (6)	0.0268 (6)	0.0191 (7)	0.0024 (5)	-0.0001 (5)	0.0100 (5)
O32	0.0292 (6)	0.0258 (6)	0.0242 (7)	-0.0011 (5)	-0.0008 (5)	0.0105 (5)
O3W	0.0355 (7)	0.0367 (7)	0.0249 (8)	0.0124 (6)	0.0078 (6)	0.0113 (6)
O4W	0.0306 (7)	0.0281 (6)	0.0273 (9)	0.0037 (6)	0.0009 (5)	0.0058 (6)
O5W	0.0247 (6)	0.0248 (6)	0.0455 (9)	0.0092 (5)	0.0019 (5)	0.0166 (6)
N1	0.0216 (6)	0.0175 (6)	0.0158 (7)	0.0058 (5)	0.0009 (5)	0.0051 (5)
N4	0.0232 (6)	0.0203 (6)	0.0211 (8)	0.0084 (5)	0.0008 (5)	0.0069 (5)

N11	0.0193 (6)	0.0221 (6)	0.0213 (8)	0.0071 (5)	0.0034 (5)	0.0098 (6)
N12	0.0257 (7)	0.0217 (7)	0.0235 (9)	0.0014 (6)	0.0001 (6)	0.0098 (6)
C2	0.0194 (7)	0.0173 (7)	0.0152 (9)	0.0045 (5)	0.0001 (6)	0.0040 (6)
C3	0.0193 (7)	0.0178 (7)	0.0167 (9)	0.0035 (5)	-0.0015 (6)	0.0042 (6)
C5	0.0224 (7)	0.0232 (7)	0.0205 (9)	0.0101 (6)	0.0038 (6)	0.0053 (6)
C6	0.0233 (7)	0.0223 (7)	0.0192 (9)	0.0081 (6)	0.0061 (6)	0.0070 (6)
C12	0.0157 (7)	0.0239 (7)	0.0232 (10)	0.0081 (6)	0.0024 (6)	0.0099 (7)
C13	0.0206 (7)	0.0264 (8)	0.0288 (11)	0.0072 (6)	0.0056 (6)	0.0139 (7)
C14	0.0259 (8)	0.0404 (10)	0.0286 (11)	0.0145 (7)	0.0109 (7)	0.0194 (8)
C15	0.0276 (8)	0.0383 (9)	0.0201 (10)	0.0159 (7)	0.0057 (7)	0.0072 (8)
C16	0.0200 (7)	0.0268 (8)	0.0260 (10)	0.0123 (6)	0.0021 (6)	0.0062 (7)
C17	0.0326 (9)	0.0248 (8)	0.0312 (11)	0.0111 (7)	-0.0016 (7)	0.0031 (7)
C21	0.0211 (7)	0.0243 (8)	0.0196 (9)	0.0086 (6)	0.0009 (6)	0.0066 (7)
C31	0.0214 (7)	0.0220 (7)	0.0230 (10)	0.0079 (6)	0.0019 (6)	0.0104 (7)

*Geometric parameters (Å, °)*

Co1—O21 <sup>i</sup>	2.0790 (12)	N11—C12	1.357 (2)
Co1—O21	2.0790 (11)	N11—C16	1.363 (2)
Co1—O1W	2.0841 (12)	N11—H11	0.82 (2)
Co1—O1W <sup>i</sup>	2.0841 (12)	N12—C12	1.325 (2)
Co1—N1	2.1045 (12)	N12—H121	0.86 (2)
Co1—N1 <sup>i</sup>	2.1045 (12)	N12—H122	0.84 (2)
O1W—H1W1	0.77 (3)	C2—C3	1.393 (2)
O1W—H1W2	0.86 (3)	C2—C21	1.516 (2)
O21—C21	1.2760 (19)	C3—C31	1.518 (2)
O22—C21	1.228 (2)	C5—C6	1.388 (2)
O2W—H2W1	0.85 (3)	C5—H5	0.9300
O2W—H2W2	0.72 (3)	C6—H6	0.9300
O31—C31	1.256 (2)	C12—C13	1.410 (2)
O32—C31	1.244 (2)	C13—C14	1.361 (3)
O3W—H3W1	0.80 (3)	C13—H13	0.9300
O3W—H3W2	0.84 (3)	C14—C15	1.404 (3)
O4W—H4W1	0.80 (3)	C14—H14	0.9300
O4W—H4W2	0.80 (3)	C15—C16	1.365 (3)
O5W—H5W1	0.79 (3)	C15—H15	0.9300
O5W—H5W2	0.77 (3)	C16—C17	1.494 (2)
N1—C6	1.329 (2)	C17—H17A	0.9600
N1—C2	1.344 (2)	C17—H17B	0.9600
N4—C5	1.335 (2)	C17—H17C	0.9600
N4—C3	1.342 (2)		
O21 <sup>i</sup> —Co1—O21	180.00 (7)	N4—C3—C2	121.24 (14)
O21 <sup>i</sup> —Co1—O1W	90.50 (5)	N4—C3—C31	114.49 (13)
O21—Co1—O1W	89.50 (5)	C2—C3—C31	124.27 (14)
O21 <sup>i</sup> —Co1—O1W <sup>i</sup>	89.50 (5)	N4—C5—C6	121.69 (14)
O21—Co1—O1W <sup>i</sup>	90.50 (5)	N4—C5—H5	119.20
O1W—Co1—O1W <sup>i</sup>	180.00 (6)	C6—C5—H5	119.20



## supplementary materials

---

O21 <sup>i</sup> —Co1—N1	100.88 (5)	N1—C6—C5	120.75 (14)
O21—Co1—N1	79.12 (5)	N1—C6—H6	119.60
O1W—Co1—N1	92.48 (5)	C5—C6—H6	119.60
O1W <sup>i</sup> —Co1—N1	87.52 (5)	N12—C12—N11	119.07 (15)
O21 <sup>i</sup> —Co1—N1 <sup>i</sup>	79.12 (5)	N12—C12—C13	123.23 (15)
O21—Co1—N1 <sup>i</sup>	100.88 (5)	N11—C12—C13	117.70 (15)
O1W—Co1—N1 <sup>i</sup>	87.52 (5)	C14—C13—C12	119.36 (15)
O1W <sup>i</sup> —Co1—N1 <sup>i</sup>	92.48 (5)	C14—C13—H13	120.30
N1—Co1—N1 <sup>i</sup>	180.00 (6)	C12—C13—H13	120.30
Co1—O1W—H1W1	119.7 (19)	C13—C14—C15	121.07 (16)
Co1—O1W—H1W2	122.8 (17)	C13—C14—H14	119.50
H1W1—O1W—H1W2	109 (2)	C15—C14—H14	119.50
C21—O21—Co1	116.29 (10)	C16—C15—C14	119.19 (17)
H2W1—O2W—H2W2	110 (3)	C16—C15—H15	120.40
H3W1—O3W—H3W2	108 (3)	C14—C15—H15	120.40
H4W1—O4W—H4W2	104 (3)	N11—C16—C15	118.88 (15)
H5W1—O5W—H5W2	106 (3)	N11—C16—C17	116.76 (16)
C6—N1—C2	118.50 (13)	C15—C16—C17	124.35 (17)
C6—N1—Co1	128.65 (11)	C16—C17—H17A	109.50
C2—N1—Co1	112.57 (10)	C16—C17—H17B	109.50
C5—N4—C3	117.44 (13)	H17A—C17—H17B	109.50
C12—N11—C16	123.77 (15)	C16—C17—H17C	109.50
C12—N11—H11	117.9 (15)	H17A—C17—H17C	109.50
C16—N11—H11	118.3 (15)	H17B—C17—H17C	109.50
C12—N12—H121	117.3 (15)	O22—C21—O21	125.85 (15)
C12—N12—H122	119.2 (16)	O22—C21—C2	118.38 (14)
H121—N12—H122	123 (2)	O21—C21—C2	115.77 (13)
N1—C2—C3	120.37 (14)	O32—C31—O31	126.69 (15)
N1—C2—C21	116.14 (13)	O32—C31—C3	116.34 (15)
C3—C2—C21	123.47 (14)	O31—C31—C3	116.80 (13)
O21 <sup>i</sup> —Co1—O21—C21	113 (47)	C3—N4—C5—C6	-1.4 (2)
O1W—Co1—O21—C21	93.19 (12)	C2—N1—C6—C5	0.3 (2)
O1W <sup>i</sup> —Co1—O21—C21	-86.81 (12)	Co1—N1—C6—C5	173.81 (11)
N1—Co1—O21—C21	0.57 (11)	N4—C5—C6—N1	0.9 (2)
N1 <sup>i</sup> —Co1—O21—C21	-179.43 (11)	C16—N11—C12—N12	-179.22 (14)
O21 <sup>i</sup> —Co1—N1—C6	3.94 (14)	C16—N11—C12—C13	0.6 (2)
O21—Co1—N1—C6	-176.06 (14)	N12—C12—C13—C14	178.72 (15)
O1W—Co1—N1—C6	94.93 (14)	N11—C12—C13—C14	-1.1 (2)
O1W <sup>i</sup> —Co1—N1—C6	-85.07 (14)	C12—C13—C14—C15	0.3 (2)
N1 <sup>i</sup> —Co1—N1—C6	85 (62)	C13—C14—C15—C16	1.0 (2)
O21 <sup>i</sup> —Co1—N1—C2	177.76 (10)	C12—N11—C16—C15	0.7 (2)
O21—Co1—N1—C2	-2.24 (10)	C12—N11—C16—C17	-179.59 (14)
O1W—Co1—N1—C2	-91.26 (11)	C14—C15—C16—N11	-1.5 (2)
O1W <sup>i</sup> —Co1—N1—C2	88.74 (11)	C14—C15—C16—C17	178.83 (15)
N1 <sup>i</sup> —Co1—N1—C2	-102 (62)	Co1—O21—C21—O22	-179.10 (14)

C6—N1—C2—C3	-0.9 (2)	Co1—O21—C21—C2	1.06 (18)
Co1—N1—C2—C3	-175.43 (11)	N1—C2—C21—O22	177.04 (14)
C6—N1—C2—C21	177.95 (13)	C3—C2—C21—O22	-4.1 (2)
Co1—N1—C2—C21	3.44 (16)	N1—C2—C21—O21	-3.1 (2)
C5—N4—C3—C2	0.7 (2)	C3—C2—C21—O21	175.73 (14)
C5—N4—C3—C31	-179.98 (13)	N4—C3—C31—O32	-85.16 (17)
N1—C2—C3—N4	0.4 (2)	C2—C3—C31—O32	94.11 (19)
C21—C2—C3—N4	-178.37 (14)	N4—C3—C31—O31	90.54 (17)
N1—C2—C3—C31	-178.80 (14)	C2—C3—C31—O31	-90.20 (19)
C21—C2—C3—C31	2.4 (2)		

Symmetry codes: (i)  $-x+2, -y+1, -z$ .

*Hydrogen-bond geometry (Å, °)*

<i>D</i> —H... <i>A</i>	<i>D</i> —H	H... <i>A</i>	<i>D</i> ... <i>A</i>	<i>D</i> —H... <i>A</i>
N11—H11...O31	0.82 (2)	1.98 (2)	2.794 (3)	179 (2)
N12—H121...O2W <sup>ii</sup>	0.86 (3)	2.05 (2)	2.900 (2)	170 (2)
N12—H122...O32	0.84 (3)	1.97 (3)	2.804 (3)	176 (2)
O1W—H1W1...O2W <sup>iii</sup>	0.78 (3)	1.99 (3)	2.770 (3)	178 (3)
O1W—H1W2...O5W <sup>iii</sup>	0.85 (3)	1.85 (3)	2.697 (2)	172 (3)
O2W—H2W1...O21	0.85 (3)	2.10 (3)	2.942 (3)	168 (3)
O2W—H2W2...O4W	0.72 (3)	2.07 (3)	2.784 (3)	176 (2)
O3W—H3W2...O4W	0.83 (3)	1.99 (3)	2.811 (3)	167 (3)
O4W—H4W1...O3W <sup>iv</sup>	0.80 (3)	1.98 (2)	2.755 (2)	165 (3)
O4W—H4W2...O31 <sup>v</sup>	0.80 (2)	1.94 (2)	2.738 (2)	178 (3)
O5W—H5W1...O32 <sup>vi</sup>	0.79 (2)	2.00 (3)	2.767 (2)	161 (3)
O5W—H5W2...N4 <sup>vii</sup>	0.77 (3)	2.11 (3)	2.871 (3)	167 (3)

Symmetry codes: (ii)  $x, y-1, z$ ; (iii)  $-x+1, -y+1, -z$ ; (iv)  $-x+1, -y+1, -z+1$ ; (v)  $x-1, y, z$ ; (vi)  $x, y+1, z$ ; (vii)  $x-1, y+1, z$ .

Fig. 1




Fig. 2




Fig. 3




Fig. 4

