

2-(Furan-2-yl)-3-hydroxy-4H-chromen-4-one

Michał Wera,^a Vasyl G. Pivovarenko,^b Artur Sikorski,^a Tadeusz Lis^c and Jerzy Błażejowski^{a*}

^aFaculty of Chemistry, University of Gdańsk, J. Sobieskiego 18, 80-952 Gdańsk, Poland, ^bFaculty of Chemistry, Kyiv Taras Shevchenko National University, Volodymyrska 64, 01033 Kyiv, Ukraine, and ^cFaculty of Chemistry, University of Wrocław, F. Joliot-Curie 14, 50-383 Wrocław, Poland

Correspondence e-mail: bla@chem.univ.gda.pl


Received 14 December 2010; accepted 21 December 2010

Key indicators: single-crystal X-ray study; $T = 100\text{ K}$; mean $\sigma(\text{C}-\text{C}) = 0.003\text{ \AA}$; R factor = 0.037; wR factor = 0.092; data-to-parameter ratio = 11.3.

In the crystal structure of the title compound, $\text{C}_{13}\text{H}_8\text{O}_4$, the inversely oriented molecules form inversion dimers through pairs of $\text{O}-\text{H}\cdots\text{O}$ hydrogen-bonding interactions. An intramolecular $\text{O}-\text{H}\cdots\text{O}$ hydrogen bond occurs. In the packing of the molecules, the nearly planar 2-(furan-2-yl)-4H-chromene units [dihedral angle between the chromene and furan rings = $3.8(1)^\circ$] are either parallel or inclined at an angle of $80.7(1)^\circ$.

Related literature

For general features of flavonols (derivatives of 3-hydroxy-2-phenyl-4H-chromen-4-one), see: Klymchenko *et al.* (2003); Sengupta & Kasha (1979). For related structures, see: Etter *et al.* (1986); Waller *et al.* (2003). For intermolecular interactions, see: Aakeröy *et al.* (1992); Novoa *et al.* (2006). For the synthesis, see: Klymchenko *et al.* (2003).



Experimental

Crystal data

$\text{C}_{13}\text{H}_8\text{O}_4$

$M_r = 228.19$

Monoclinic, $P2_1/c$
 $a = 14.365(8)\text{ \AA}$
 $b = 4.421(3)\text{ \AA}$
 $c = 17.086(10)\text{ \AA}$
 $\beta = 110.91(5)^\circ$
 $V = 1013.6(11)\text{ \AA}^3$

$Z = 4$
 Mo $K\alpha$ radiation
 $\mu = 0.11\text{ mm}^{-1}$
 $T = 100\text{ K}$
 $0.40 \times 0.40 \times 0.14\text{ mm}$

Data collection

Kuma KM4 CCD κ -geometry diffractometer
 7132 measured reflections
 1779 independent reflections
 1436 reflections with $I > 2\sigma(I)$
 $R_{\text{int}} = 0.030$

Refinement

$R[F^2 > 2\sigma(F^2)] = 0.037$
 $wR(F^2) = 0.092$
 $S = 1.10$
 1779 reflections
 158 parameters
 H atoms treated by a mixture of independent and constrained refinement
 $\Delta\rho_{\text{max}} = 0.21\text{ e \AA}^{-3}$
 $\Delta\rho_{\text{min}} = -0.21\text{ e \AA}^{-3}$

Table 1
 Hydrogen-bond geometry (\AA , $^\circ$).

$D-\text{H}\cdots A$	$D-\text{H}$	$\text{H}\cdots A$	$D\cdots A$	$D-\text{H}\cdots A$
C6—H6···O12 ⁱ	0.97 (2)	2.53 (2)	3.352 (3)	142 (2)
O11—H11···O12	0.89 (2)	2.37 (2)	2.776 (3)	108 (2)
O11—H11···O12 ⁱⁱ	0.89 (2)	1.87 (2)	2.683 (3)	152 (2)

Symmetry codes: (i) $-x, y - \frac{1}{2}, -z + \frac{1}{2}$; (ii) $-x, -y + 1, -z + 1$.

Data collection: *CrysAlis CCD* (Oxford Diffraction, 2003); cell refinement: *CrysAlis RED* (Oxford Diffraction, 2003); data reduction: *CrysAlis RED*; program(s) used to solve structure: *SHELXS97* (Sheldrick, 2008); program(s) used to refine structure: *SHELXL97* (Sheldrick, 2008); molecular graphics: *ORTEP-3* (Farrugia, 1997); software used to prepare material for publication: *SHELXL97* and *PLATON* (Spek, 2009).

This study was financed by the State Funds for Scientific Research (grant DS/8220-4-0087-9).

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: NG5090).

References

- Aakeröy, C. B., Seddon, K. R. & Leslie, M. (1992). *Struct. Chem.* **3**, 63–65.
- Etter, M. C., Urbańczyk-Lipkowska, Z., Baer, S. & Barbara, P. F. (1986). *J. Mol. Struct.* **144**, 155–167.
- Farrugia, L. J. (1997). *J. Appl. Cryst.* **30**, 565.
- Klymchenko, A. S., Pivovarenko, V. G. & Demchenko, A. P. (2003). *Spectrochim. Acta Part A*, **59**, 787–792.
- Novoa, J. J., Mota, F. & D’Oria, E. (2006). *Hydrogen Bonding – New Insights*, edited by S. Grabowski, pp. 193–244. The Netherlands: Springer.
- Oxford Diffraction (2003). *KM-4-CCD Software*. Oxford Diffraction Poland, Wrocław, Poland.
- Sengupta, P. K. & Kasha, M. (1979). *Chem. Phys. Lett.* **68**, 382–385.
- Sheldrick, G. M. (2008). *Acta Cryst. A* **64**, 112–122.
- Spek, A. L. (2009). *Acta Cryst. D* **65**, 148–155.
- Waller, M. P., Hibbs, D. E., Overgaard, J., Hanrahan, J. R. & Hambley, T. W. (2003). *Acta Cryst. E* **59**, o767–o768.

supplementary materials

Acta Cryst. (2011). E67, o266 [doi:10.1107/S1600536810053596]

2-(Furan-2-yl)-3-hydroxy-4H-chromen-4-one

M. Wera, V. G. Pivovarenko, A. Sikorski, T. Lis and J. Blazejowski

Comment

The structure of 2-(furan-2-yl)-3-hydroxy-4H-chromen-4-one is presented. This compound, in which Excited State Intramolecular Proton Transfer (ESIPT) takes place (Sengupta & Kasha, 1979), is a good candidate for a fluorescent probe sensitive to the properties of a medium (Klymchenko *et al.*, 2003).

In the title compound (Fig. 1), the bond lengths and angles characterizing the geometry of the 4H-chromen-4-one moiety are similar to those in 2-phenyl-4H-chromen-4-one (Waller *et al.*, 2003) and 3-hydroxy-2-phenyl-4H-chromen-4-one (Etter *et al.*, 1986). The average deviations from planarity of the phenyl, 4H-chromene and 2-(furan-2-yl)-4H-chromene cores are 0.0024 (2), 0.0046 (2) and 0.0298 (2), respectively, which implies that the molecule is practically planar (the dihedral angle between the planes of the 4H-chromene and furanyl fragments is only 3.8 (1) $^{\circ}$). Intramolecular O–H \cdots O and C–H \cdots O interactions (Table 1, Figs. 1 and 2) undoubtedly make the molecule more rigid and contribute to its planarity, the former being the one involved in the ESIPT characteristic of flavonols (Sengupta & Kasha, 1979). The mean planes of adjacent 2-(furan-2-yl)-4H-chromene moieties are either parallel (remain at an angle 0.0 (1) $^{\circ}$) in the crystal lattice or are inclined at an angle of 80.7 (1) $^{\circ}$.

In the crystal structure, the inversely oriented molecules form dimers through a pair of O–H \cdots O (Aakeröy *et al.*, 1992) interactions (Table 1, Fig. 2). Adjacent dimers are linked by C–H \cdots O (Novoa *et al.*, 2006) interactions (Table 1, Fig. 2). The crystal structure is stabilized by these specific interactions, as well as by non-specific dispersive interactions.

Experimental

The title compound was obtained by means of the oxidative heterocyclization of 3-(furan-2-yl)-1-(2-hydroxyphenyl)prop-2-en-1-one, synthesized by the condensation of 1-(2-hydroxyphenyl)ethanone with furan-2-carbaldehyde in methanol/50% aqueous NaOH (1/1 v/v), in alkaline methanol/H₂O₂ (Klymchenko *et al.*, 2003). The product was separated by filtration, and greenish-yellow crystals suitable for X-ray investigations were grown from ethanol (m.p. = 445 – 446 K).

Refinement

H atoms involved in C–H \cdots O and O–H \cdots O interactions were located on a difference Fourier map and refined isotropically with $U_{\text{iso}}(\text{H}) = 1.2U_{\text{eq}}(\text{C})$ and $U_{\text{iso}}(\text{H}) = 1.5U_{\text{eq}}(\text{O})$, respectively. H atoms of other C–H bonds were positioned geometrically, with C–H = 0.95 Å, and constrained to ride on their parent atoms with $U_{\text{iso}}(\text{H}) = 1.2U_{\text{eq}}(\text{C})$.

supplementary materials

Figures




Fig. 1. The molecular structure of the title compound showing the atom-labeling scheme. Displacement ellipsoids are drawn at the 25% probability level, and H atoms are shown as small spheres of arbitrary radius. The O—H···O and C—H···O hydrogen bonds are represented by dashed lines.




Fig. 2. The arrangement of the molecules in the crystal structure. The O—H···O and C—H···O interactions are represented by dashed lines. H atoms not involved in interactions have been omitted. [Symmetry codes: (i) $-x, y - 1/2, -z + 1/2$; (ii) $-x, -y + 1, -z + 1$.]

2-(Furan-2-yl)-3-hydroxy-4*H*-chromen-4-one

Crystal data

C ₁₃ H ₈ O ₄	F(000) = 472
M _r = 228.19	D _x = 1.495 Mg m ⁻³
Monoclinic, P2 ₁ /c	Mo K α radiation, λ = 0.71073 Å
Hall symbol: -P 2ybc	Cell parameters from 1436 reflections
a = 14.365 (8) Å	θ = 3.4–25.0°
b = 4.421 (3) Å	μ = 0.11 mm ⁻¹
c = 17.086 (10) Å	T = 100 K
β = 110.91 (5)°	Plate, greenish-yellow
V = 1013.6 (11) Å ³	0.40 × 0.40 × 0.14 mm
Z = 4	

Data collection

Kuma KM4 CCD κ -geometry diffractometer	1436 reflections with $I > 2\sigma(I)$
Radiation source: fine-focus sealed tube	$R_{\text{int}} = 0.030$
graphite	$\theta_{\text{max}} = 25.0^\circ$, $\theta_{\text{min}} = 3.4^\circ$
ω scans	$h = -17 \rightarrow 17$
7132 measured reflections	$k = -4 \rightarrow 5$
1779 independent reflections	$l = -20 \rightarrow 18$

Refinement

Refinement on F^2	Primary atom site location: structure-invariant direct methods
Least-squares matrix: full	Secondary atom site location: difference Fourier map
$R[F^2 > 2\sigma(F^2)] = 0.037$	Hydrogen site location: inferred from neighbouring sites

$wR(F^2) = 0.092$	H atoms treated by a mixture of independent and constrained refinement
$S = 1.10$	$w = 1/[\sigma^2(F_o^2) + (0.0517P)^2 + 0.0439P]$
1779 reflections	where $P = (F_o^2 + 2F_c^2)/3$
158 parameters	$(\Delta/\sigma)_{\max} < 0.001$
0 restraints	$\Delta\rho_{\max} = 0.21 \text{ e } \text{\AA}^{-3}$
	$\Delta\rho_{\min} = -0.21 \text{ e } \text{\AA}^{-3}$

Special details

Geometry. All e.s.d.'s (except the e.s.d. in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell e.s.d.'s are taken into account individually in the estimation of e.s.d.'s in distances, angles and torsion angles; correlations between e.s.d.'s in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell e.s.d.'s is used for estimating e.s.d.'s involving l.s. planes.

Refinement. Refinement of F^2 against ALL reflections. The weighted R -factor wR and goodness of fit S are based on F^2 , conventional R -factors R are based on F , with F set to zero for negative F^2 . The threshold expression of $F^2 > \sigma(F^2)$ is used only for calculating R -factors(gt) etc. and is not relevant to the choice of reflections for refinement. R -factors based on F^2 are statistically about twice as large as those based on F , and R -factors based on ALL data will be even larger.

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (\AA^2)

	x	y	z	$U_{\text{iso}}^*/U_{\text{eq}}$
O1	0.30889 (8)	0.5549 (2)	0.43752 (6)	0.0209 (3)
C2	0.27113 (12)	0.6670 (4)	0.49500 (9)	0.0189 (4)
C3	0.17955 (12)	0.5867 (4)	0.49522 (9)	0.0192 (4)
C4	0.11644 (12)	0.3802 (4)	0.43368 (9)	0.0200 (4)
C5	0.10589 (12)	0.0603 (4)	0.30910 (9)	0.0212 (4)
H5	0.0416	-0.0076	0.3051	0.025*
C6	0.14678 (13)	-0.0412 (4)	0.25248 (10)	0.0232 (4)
H6	0.1102 (13)	-0.177 (4)	0.2072 (11)	0.028*
C7	0.24244 (13)	0.0564 (4)	0.25881 (10)	0.0248 (4)
H7	0.2707	-0.0148	0.2197	0.030*
C8	0.29541 (13)	0.2533 (4)	0.32077 (10)	0.0229 (4)
H8	0.3601	0.3182	0.3250	0.027*
C9	0.15832 (12)	0.2640 (4)	0.37305 (9)	0.0193 (4)
C10	0.25266 (12)	0.3566 (4)	0.37755 (9)	0.0189 (4)
O11	0.14832 (9)	0.7044 (3)	0.55453 (7)	0.0266 (3)
H11	0.0840 (16)	0.665 (5)	0.5432 (12)	0.040*
O12	0.03228 (8)	0.3089 (3)	0.43240 (7)	0.0277 (3)
C13	0.33852 (12)	0.8757 (4)	0.55296 (9)	0.0198 (4)
O14	0.42459 (8)	0.9419 (3)	0.53893 (7)	0.0245 (3)
C15	0.47607 (12)	1.1419 (4)	0.59996 (10)	0.0264 (4)
H15	0.5387	1.2273	0.6053	0.032*
C16	0.42634 (12)	1.2018 (4)	0.65152 (10)	0.0243 (4)
H16	0.4471	1.3323	0.6988	0.029*
C17	0.33603 (13)	1.0302 (4)	0.62119 (10)	0.0224 (4)
H17	0.2820 (13)	1.024 (4)	0.6427 (10)	0.027*

supplementary materials

Atomic displacement parameters (\AA^2)

	U^{11}	U^{22}	U^{33}	U^{12}	U^{13}	U^{23}
O1	0.0202 (6)	0.0231 (6)	0.0215 (6)	-0.0020 (5)	0.0099 (5)	-0.0031 (5)
C2	0.0205 (9)	0.0197 (8)	0.0168 (8)	0.0034 (7)	0.0072 (7)	0.0038 (7)
C3	0.0188 (9)	0.0231 (9)	0.0162 (8)	0.0023 (7)	0.0069 (7)	0.0040 (7)
C4	0.0192 (9)	0.0211 (9)	0.0195 (8)	0.0019 (7)	0.0068 (7)	0.0053 (7)
C5	0.0186 (9)	0.0220 (9)	0.0212 (8)	0.0007 (7)	0.0050 (7)	0.0036 (7)
C6	0.0256 (10)	0.0219 (9)	0.0196 (9)	-0.0008 (7)	0.0049 (7)	0.0010 (7)
C7	0.0315 (10)	0.0231 (9)	0.0232 (9)	0.0034 (8)	0.0141 (8)	0.0014 (7)
C8	0.0208 (9)	0.0247 (9)	0.0257 (9)	-0.0013 (7)	0.0115 (7)	0.0014 (7)
C9	0.0198 (9)	0.0200 (9)	0.0170 (8)	0.0020 (7)	0.0051 (7)	0.0044 (6)
C10	0.0192 (8)	0.0180 (9)	0.0187 (8)	0.0010 (7)	0.0055 (7)	0.0028 (7)
O11	0.0201 (7)	0.0390 (8)	0.0226 (6)	-0.0047 (6)	0.0100 (5)	-0.0072 (5)
O12	0.0202 (7)	0.0377 (7)	0.0265 (6)	-0.0055 (5)	0.0099 (5)	-0.0040 (5)
C13	0.0164 (8)	0.0223 (9)	0.0207 (8)	0.0020 (7)	0.0066 (7)	0.0059 (7)
O14	0.0203 (6)	0.0280 (7)	0.0265 (6)	-0.0054 (5)	0.0100 (5)	-0.0047 (5)
C15	0.0201 (9)	0.0258 (10)	0.0290 (9)	-0.0055 (7)	0.0037 (7)	-0.0052 (8)
C16	0.0259 (9)	0.0228 (9)	0.0219 (8)	0.0000 (7)	0.0056 (7)	-0.0004 (7)
C17	0.0221 (9)	0.0238 (9)	0.0211 (8)	0.0022 (7)	0.0075 (7)	0.0021 (7)

Geometric parameters (\AA , $^\circ$)

O1—C10	1.371 (2)	C7—H7	0.9500
O1—C2	1.3729 (19)	C8—C10	1.397 (2)
C2—C3	1.364 (2)	C8—H8	0.9500
C2—C13	1.444 (2)	C9—C10	1.391 (2)
C3—O11	1.3503 (19)	O11—H11	0.89 (2)
C3—C4	1.443 (2)	C13—C17	1.362 (2)
C4—O12	1.2419 (19)	C13—O14	1.372 (2)
C4—C9	1.464 (2)	O14—C15	1.366 (2)
C5—C6	1.374 (2)	C15—C16	1.344 (2)
C5—C9	1.409 (2)	C15—H15	0.9500
C5—H5	0.9500	C16—C17	1.431 (3)
C6—C7	1.407 (2)	C16—H16	0.9500
C6—H6	0.971 (18)	C17—H17	0.971 (17)
C7—C8	1.372 (2)		
C10—O1—C2	119.19 (13)	C10—C8—H8	120.6
C3—C2—O1	122.39 (15)	C10—C9—C5	118.13 (15)
C3—C2—C13	125.15 (14)	C10—C9—C4	119.72 (15)
O1—C2—C13	112.46 (14)	C5—C9—C4	122.15 (15)
O11—C3—C2	118.75 (15)	O1—C10—C9	122.13 (14)
O11—C3—C4	120.01 (14)	O1—C10—C8	116.11 (14)
C2—C3—C4	121.24 (14)	C9—C10—C8	121.76 (15)
O12—C4—C3	121.87 (15)	C3—O11—H11	110.9 (13)
O12—C4—C9	122.81 (15)	C17—C13—O14	110.06 (15)
C3—C4—C9	115.33 (14)	C17—C13—C2	133.72 (16)

C6—C5—C9	120.69 (16)	O14—C13—C2	116.21 (14)
C6—C5—H5	119.7	C15—O14—C13	106.30 (13)
C9—C5—H5	119.7	C16—C15—O14	111.01 (15)
C5—C6—C7	119.78 (16)	C16—C15—H15	124.5
C5—C6—H6	121.1 (10)	O14—C15—H15	124.5
C7—C6—H6	119.1 (10)	C15—C16—C17	106.47 (15)
C8—C7—C6	120.76 (15)	C15—C16—H16	126.8
C8—C7—H7	119.6	C17—C16—H16	126.8
C6—C7—H7	119.6	C13—C17—C16	106.17 (16)
C7—C8—C10	118.87 (16)	C13—C17—H17	125.5 (10)
C7—C8—H8	120.6	C16—C17—H17	128.4 (10)
C10—O1—C2—C3	0.5 (2)	C2—O1—C10—C9	-0.2 (2)
C10—O1—C2—C13	-179.14 (13)	C2—O1—C10—C8	179.56 (13)
O1—C2—C3—O11	179.20 (13)	C5—C9—C10—O1	179.80 (14)
C13—C2—C3—O11	-1.2 (2)	C4—C9—C10—O1	0.1 (2)
O1—C2—C3—C4	-0.8 (2)	C5—C9—C10—C8	0.1 (2)
C13—C2—C3—C4	178.82 (14)	C4—C9—C10—C8	-179.59 (14)
O11—C3—C4—O12	1.2 (2)	C7—C8—C10—O1	-179.42 (14)
C2—C3—C4—O12	-178.86 (15)	C7—C8—C10—C9	0.3 (2)
O11—C3—C4—C9	-179.30 (13)	C3—C2—C13—C17	3.7 (3)
C2—C3—C4—C9	0.7 (2)	O1—C2—C13—C17	-176.67 (16)
C9—C5—C6—C7	0.7 (2)	C3—C2—C13—O14	-175.90 (14)
C5—C6—C7—C8	-0.3 (2)	O1—C2—C13—O14	3.73 (19)
C6—C7—C8—C10	-0.2 (2)	C17—C13—O14—C15	-0.06 (17)
C6—C5—C9—C10	-0.6 (2)	C2—C13—O14—C15	179.64 (13)
C6—C5—C9—C4	179.08 (14)	C13—O14—C15—C16	0.35 (18)
O12—C4—C9—C10	179.17 (14)	O14—C15—C16—C17	-0.49 (19)
C3—C4—C9—C10	-0.4 (2)	O14—C13—C17—C16	-0.23 (18)
O12—C4—C9—C5	-0.5 (2)	C2—C13—C17—C16	-179.85 (17)
C3—C4—C9—C5	179.98 (14)	C15—C16—C17—C13	0.44 (19)

Hydrogen-bond geometry (Å, °)

D—H···A	D—H	H···A	D···A	D—H···A
C6—H6···O12 ⁱ	0.97 (2)	2.53 (2)	3.352 (3)	142 (2)
O11—H11···O12	0.89 (2)	2.37 (2)	2.776 (3)	108 (2)
O11—H11···O12 ⁱⁱ	0.89 (2)	1.87 (2)	2.683 (3)	152 (2)
C17—H17···O11	0.97 (2)	2.43 (2)	2.907 (3)	110 (2)

Symmetry codes: (i) $-x, y-1/2, -z+1/2$; (ii) $-x, -y+1, -z+1$.

supplementary materials

Fig. 1




Fig. 2

