

1-(3-Chlorophenyl)-4,4,6-trimethyl-3,4-dihydropyrimidine-2(1H)-thione

Bohari M. Yamin* and Halima Farag Salem

School of Chemical Sciences and Food Technology, Universiti Kebangsaan Malaysia, UKM 43600 Bangi Selangor, Malaysia

Correspondence e-mail: bohari@ukm.my


Received 19 December 2010; accepted 25 December 2010

Key indicators: single-crystal X-ray study; $T = 298$ K; mean $\sigma(\text{C}-\text{C}) = 0.004$ Å; R factor = 0.048; wR factor = 0.133; data-to-parameter ratio = 16.5.

In the title compound, $\text{C}_{13}\text{H}_{15}\text{ClN}_2\text{S}$, the dihydropyrimidine ring is essentially planar, with a maximum deviation from the least-squares plane of 0.122 (3) Å for the unsubstituted olefinic C atom. The dihedral angle between the dihydropyrimidine and benzene rings is 86.62 (13)°. The crystal structure is stabilized by intermolecular $\text{N}-\text{H}\cdots\text{S}$ hydrogen bonds, which form centrosymmetric dimers arranged along the c axis.

Related literature

For related structures, see: Yamin *et al.* (2005); Ismail *et al.* (2007); Saeed & Bolte, (2010). For the biological activity of dihydropyrimidinone/thione derivatives, see: Alam *et al.* (2005); Kappe (2000); Sriram *et al.* (2006); Leite *et al.* (2006). For graph-set theory, see: Etter *et al.* (1990); Bernstein *et al.* (1995).



Experimental

Crystal data

$\text{C}_{13}\text{H}_{15}\text{ClN}_2\text{S}$
 $M_r = 266.78$
 Monoclinic, $P2_1/c$
 $a = 8.398$ (2) Å
 $b = 14.930$ (4) Å
 $c = 11.468$ (3) Å
 $\beta = 103.909$ (4)°

$V = 1395.7$ (6) Å³
 $Z = 4$
 Mo $K\alpha$ radiation
 $\mu = 0.40$ mm⁻¹
 $T = 298$ K
 $0.40 \times 0.19 \times 0.17$ mm

Data collection

Bruker SMART APEX CCD area-detector diffractometer
 Absorption correction: multi-scan (SADABS; Bruker, 2000)
 $T_{\min} = 0.855$, $T_{\max} = 0.934$
 8215 measured reflections
 2598 independent reflections
 2212 reflections with $I > 2\sigma(I)$
 $R_{\text{int}} = 0.021$

Refinement

$R[F^2 > 2\sigma(F^2)] = 0.048$
 $wR(F^2) = 0.133$
 $S = 1.10$
 2598 reflections
 157 parameters
 H-atom parameters constrained
 $\Delta\rho_{\max} = 0.29$ e Å⁻³
 $\Delta\rho_{\min} = -0.14$ e Å⁻³

Table 1

Hydrogen-bond geometry (Å, °).

$D-\text{H}\cdots A$	$D-\text{H}$	$\text{H}\cdots A$	$D\cdots A$	$D-\text{H}\cdots A$
$\text{N1}-\text{H1A}\cdots\text{S1}^i$	0.85	2.58	3.404 (2)	162

Symmetry code: (i) $-x + 2, -y + 2, -z + 1$.

Data collection: SMART (Bruker, 2000); cell refinement: SAINT (Bruker, 2000); data reduction: SAINT; program(s) used to solve structure: SHELXS97 (Sheldrick, 2008); program(s) used to refine structure: SHELXL97 (Sheldrick, 2008); molecular graphics: ORTEPIII (Burnett & Johnson, 1996), ORTEP-3 for Windows (Farrugia, 1997) and PLATON (Spek, 2009); software used to prepare material for publication: SHELXL97, PARST (Nardelli, 1995) and PLATON.

The authors thank the Ministry of Higher Education of Malaysia and Universiti Kebangsaan Malaysia for the research grant UKM-ST-06-FRGS0114–2009. A scholarship from the Libyan Government to SFH is greatly appreciated.

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: DN2640).

References

- Alam, O., Imran, M. & Khan, S. A. (2005). *Indian J. Heterocycl. Chem.* **14**, 293–296.
- Bernstein, J., Davis, R. E., Shimon, L. & Chang, N.-L. (1995). *Angew. Chem. Int. Ed. Engl.* **34**, 1555–1573.
- Bruker (2000). *SADABS, SMART and SAINT*. Bruker AXS Inc., Madison, Wisconsin, USA.
- Burnett, M. N. & Johnson, C. K. (1996). *ORTEPIII*. Report ORNL-6895. Oak Ridge National Laboratory, Tennessee, USA.
- Etter, M. C., MacDonald, J. C. & Bernstein, J. (1990). *Acta Cryst.* **B46**, 256–262.
- Farrugia, L. J. (1997). *J. Appl. Cryst.* **30**, 565.
- Ismail, N. L., Othman, E. & Yamin, B. M. (2007). *Acta Cryst.* **E63**, o2442–o2443.
- Kappe, C. O. (2000). *Acc. Chem. Res.* **33**, 879–888.
- Leite, A. C. L., Lima, R. S., Moreira, D. R. M., Cardoso, M. V. O., Brito, A. C. G., Santos, L. M. F., Hernandez, M. Z., Kiperstok, A. C., Lima, R. S. & Soares, M. B. P. (2006). *Bioorg. Med. Chem.* **14**, 3749–3757.
- Nardelli, M. (1995). *J. Appl. Cryst.* **28**, 659.
- Saeed, A. & Bolte, M. (2010). *Acta Cryst.* **E66**, o440.
- Sheldrick, G. M. (2008). *Acta Cryst.* **A64**, 112–122.
- Spek, A. L. (2009). *Acta Cryst.* **D65**, 148–155.
- Sriram, D., Yogeewari, P. & Devakaram, R. V. (2006). *Bioorg. Med. Chem.* **14**, 3113–3118.
- Yamin, B. M., Kasim, N. A. M. & Hamzah, N. (2005). *Acta Cryst.* **E61**, o55–o57.

supplementary materials

Acta Cryst. (2011). E67, o282 [doi:10.1107/S1600536810054292]

1-(3-Chlorophenyl)-4,4,6-trimethyl-3,4-dihydropyrimidine-2(1H)-thione

B. M. Yamin and H. F. Salem

Comment

The dihydropyrimidinone/thione derivatives are medicinally important due to their therapeutic and pharmacological properties (Kappe, 2000; Alam *et al.*, 2005; Sriram *et al.*, 2006; Leite *et al.*, 2006).

The title compound (I) is a *meta* isomer of the previously reported 1-(4-Chlorophenyl)-4,4,6-trimethyl-3,4-dihydropyrimidine-2(1H)-thione (Saeed & Bolte, 2010). The dihydropyrimidine N1/C1/N2/C2/C3/C4 ring is essentially planar with maximum deviation of 0.122 (3) Å for the unsubstituted olefinic carbon C3 atom compare to that in the *para* isomer where the C4 atom bearing the two methyl substituents deviated by 0.44 (2)%A from the other five almost coplanar atoms (Saeed & Bolte, 2010). The dihedral angle between the dihydropyrimidine and benzene ring is 86.62 (13)° (Fig. 1), smaller than that in the *para* isomer of 89.59 (5) Å. The bond lengths and bond angles agree with closely related structures (Ismail *et al.*, 2007; Yamin *et al.*, 2005).

The molecular packing is also characterized by centrosymmetric dimers connected by the N—H...S intermolecular hydrogen bond forming a $R_2^2(8)$ ring (Etter *et al.*, 1990, Bernstein *et al.*, 1995) and are arranged parallel to the *c* axis (Table 1, Fig 2).

Experimental

The title compound was prepared by the reaction of thiocynic acid (5.4 mmol) and 3-chloroaniline (5.4 mmol) in acetone. The reaction mixture was stirred for 2–3 h. Then the clear was left for slow evaporation at room temperature. Colourless crystals of 1-(3-Chlorophenyl)-4,4,6-trimethyl -3,4-dihydropyrimidine-2 (1H)-thione were obtained after three days with 80% yield. Anal. Calcd for C₁₃ H₁₅ Cl N₂ S: C, 58.53; H, 5.67; N, 10.50; S, 12.02%; found: C, 58.49; H, 5.72; N, 10.61; S, 12.14. IR(KBr), ν (cm⁻¹) 1535 (C=S), 1591 (C=C), 3184 (N—H).

Refinement

All H atoms attached to C atoms were fixed geometrically and treated as riding with C—H = 0.93 Å (aromatic) or 0.96 Å (methyl) with $U_{\text{iso}}(\text{H}) = 1.2U_{\text{eq}}(\text{C}_{\text{aromatic}})$ and $1.5U_{\text{eq}}(\text{C}_{\text{methyl}})$. The amino hydrogen atom was located from the difference map and refined freely with $U_{\text{iso}}(\text{H}) = 1.2U_{\text{eq}}(\text{N})$. In the last cycles of refinement, it was treated as riding on the parent N atom. Both methyl groups attached to C3 display rather elongated ellipsoids however no correct disordered model could be defined and these large ellipsoids may be related to dynamic motion.

Figures




Fig. 1. Molecular view of (I) with the atom-labelling scheme. Displacement ellipsoids are drawn at the 30% probability level. H atoms are represented as small spheres of arbitrary radii.




Fig. 2. The molecular packing of (I) viewed down the *a* axis. H atoms not involved in hydrogen bondings have been omitted for clarity. H bonds are represented as dashed lines. [Symmetry code: (i) $-x+2, -y+2, -z+1$]

1-(3-Chlorophenyl)-4,4,6-trimethyl-3,4-dihydropyrimidine-2(1*H*)-thione

Crystal data

$C_{13}H_{15}ClN_2S$

$M_r = 266.78$

Monoclinic, $P2_1/c$

Hall symbol: $-P\ 2ybc$

$a = 8.398\ (2)\ \text{\AA}$

$b = 14.930\ (4)\ \text{\AA}$

$c = 11.468\ (3)\ \text{\AA}$

$\beta = 103.909\ (4)^\circ$

$V = 1395.7\ (6)\ \text{\AA}^3$

$Z = 4$

$F(000) = 560$

$D_x = 1.270\ \text{Mg m}^{-3}$

Melting point = $427.6\text{--}429.6\ \text{K}$

Mo $K\alpha$ radiation, $\lambda = 0.71073\ \text{\AA}$

Cell parameters from 2921 reflections

$\theta = 2.2\text{--}25.5^\circ$

$\mu = 0.40\ \text{mm}^{-1}$

$T = 298\ \text{K}$

Block, colourless

$0.40 \times 0.19 \times 0.17\ \text{mm}$

Data collection

Bruker SMART APEX CCD area-detector diffractometer

Radiation source: fine-focus sealed tube graphite

Detector resolution: $83.66\ \text{pixels mm}^{-1}$

ω scans

Absorption correction: multi-scan (*SADABS*; Bruker, 2000)

$T_{\min} = 0.855, T_{\max} = 0.934$

8215 measured reflections

2598 independent reflections

2212 reflections with $I > 2\sigma(I)$

$R_{\text{int}} = 0.021$

$\theta_{\max} = 25.5^\circ, \theta_{\min} = 2.2^\circ$

$h = -10 \rightarrow 10$

$k = -13 \rightarrow 18$

$l = -12 \rightarrow 13$

Refinement

Refinement on F^2

Least-squares matrix: full

Primary atom site location: structure-invariant direct methods

Secondary atom site location: difference Fourier map

$R[F^2 > 2\sigma(F^2)] = 0.048$	Hydrogen site location: inferred from neighbouring sites
$wR(F^2) = 0.133$	H-atom parameters constrained
$S = 1.10$	$w = 1/[\sigma^2(F_o^2) + (0.0696P)^2 + 0.4092P]$
2598 reflections	where $P = (F_o^2 + 2F_c^2)/3$
157 parameters	$(\Delta/\sigma)_{\max} < 0.001$
0 restraints	$\Delta\rho_{\max} = 0.29 \text{ e } \text{\AA}^{-3}$
	$\Delta\rho_{\min} = -0.14 \text{ e } \text{\AA}^{-3}$

Special details

Geometry. All esds (except the esd in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell esds are taken into account individually in the estimation of esds in distances, angles and torsion angles; correlations between esds in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell esds is used for estimating esds involving l.s. planes.

Refinement. Refinement of F^2 against ALL reflections. The weighted R -factor wR and goodness of fit S are based on F^2 , conventional R -factors R are based on F , with F set to zero for negative F^2 . The threshold expression of $F^2 > \sigma(F^2)$ is used only for calculating R -factors(gt) *etc.* and is not relevant to the choice of reflections for refinement. R -factors based on F^2 are statistically about twice as large as those based on F , and R -factors based on ALL data will be even larger.

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (\AA^2)

	x	y	z	$U_{\text{iso}}^*/U_{\text{eq}}$
C11	0.26790 (8)	0.97009 (5)	-0.03961 (6)	0.0647 (2)
S1	0.77220 (7)	1.01839 (4)	0.36163 (5)	0.0496 (2)
N1	0.8934 (2)	0.87127 (13)	0.47479 (18)	0.0508 (5)
H1A	0.9700	0.9086	0.5030	0.061*
N2	0.6306 (2)	0.85790 (12)	0.35950 (17)	0.0467 (5)
C1	0.7657 (3)	0.90982 (15)	0.40192 (19)	0.0408 (5)
C2	0.6333 (3)	0.76397 (16)	0.3814 (3)	0.0598 (7)
C3	0.7636 (4)	0.72845 (18)	0.4552 (3)	0.0693 (8)
H3	0.7690	0.6664	0.4621	0.083*
C4	0.9012 (3)	0.78148 (16)	0.5276 (2)	0.0602 (7)
C5	0.4926 (4)	0.7110 (2)	0.3124 (4)	0.1029 (13)
H5A	0.5133	0.6484	0.3283	0.154*
H5B	0.4787	0.7222	0.2281	0.154*
H5C	0.3947	0.7281	0.3360	0.154*
C7	1.0658 (5)	0.7413 (2)	0.5222 (5)	0.1179 (16)
H7A	1.0740	0.7392	0.4401	0.177*
H7B	1.0745	0.6818	0.5547	0.177*
H7C	1.1528	0.7777	0.5682	0.177*
C8	0.4808 (3)	0.90016 (15)	0.2952 (2)	0.0450 (5)
C9	0.4507 (3)	0.91237 (15)	0.1734 (2)	0.0448 (5)
H9	0.5261	0.8933	0.1311	0.054*
C10	0.3064 (3)	0.95355 (15)	0.1145 (2)	0.0468 (5)
C11	0.1924 (3)	0.98094 (17)	0.1755 (3)	0.0571 (7)
H11	0.0959	1.0088	0.1351	0.069*

supplementary materials

C12	0.2238 (3)	0.96636 (19)	0.2971 (3)	0.0644 (7)
H12	0.1467	0.9837	0.3389	0.077*
C13	0.3679 (3)	0.92637 (18)	0.3583 (2)	0.0578 (6)
H13	0.3887	0.9173	0.4408	0.069*
C6	0.8890 (7)	0.7904 (3)	0.6570 (3)	0.137 (2)
H6A	0.9756	0.8284	0.7001	0.206*
H6B	0.8989	0.7323	0.6940	0.206*
H6C	0.7849	0.8161	0.6590	0.206*

Atomic displacement parameters (\AA^2)

	U^{11}	U^{22}	U^{33}	U^{12}	U^{13}	U^{23}
C11	0.0586 (4)	0.0726 (5)	0.0561 (4)	-0.0055 (3)	0.0005 (3)	0.0109 (3)
S1	0.0455 (4)	0.0420 (3)	0.0541 (4)	-0.0027 (2)	-0.0022 (3)	0.0110 (3)
N1	0.0452 (11)	0.0431 (11)	0.0568 (11)	0.0034 (8)	-0.0022 (9)	0.0078 (9)
N2	0.0430 (10)	0.0391 (10)	0.0546 (11)	-0.0009 (8)	0.0050 (8)	0.0046 (8)
C1	0.0399 (11)	0.0419 (12)	0.0397 (11)	0.0035 (9)	0.0077 (9)	0.0014 (9)
C2	0.0618 (16)	0.0402 (13)	0.0743 (17)	-0.0030 (11)	0.0104 (13)	0.0033 (12)
C3	0.082 (2)	0.0373 (13)	0.081 (2)	0.0015 (13)	0.0059 (16)	0.0088 (13)
C4	0.0760 (18)	0.0425 (13)	0.0546 (14)	0.0126 (12)	0.0010 (12)	0.0076 (11)
C5	0.082 (2)	0.0519 (17)	0.158 (4)	-0.0206 (16)	-0.004 (2)	0.008 (2)
C7	0.080 (2)	0.064 (2)	0.185 (5)	0.0305 (18)	-0.015 (2)	-0.003 (3)
C8	0.0378 (11)	0.0412 (12)	0.0534 (13)	-0.0050 (9)	0.0056 (9)	-0.0017 (10)
C9	0.0393 (11)	0.0438 (12)	0.0520 (13)	-0.0049 (9)	0.0123 (10)	-0.0022 (10)
C10	0.0412 (12)	0.0414 (11)	0.0533 (13)	-0.0084 (10)	0.0025 (10)	0.0012 (10)
C11	0.0412 (13)	0.0535 (15)	0.0714 (17)	0.0033 (11)	0.0035 (12)	-0.0057 (12)
C12	0.0506 (15)	0.0716 (19)	0.0732 (19)	0.0066 (12)	0.0195 (14)	-0.0122 (14)
C13	0.0538 (15)	0.0655 (17)	0.0546 (14)	0.0006 (12)	0.0142 (12)	-0.0055 (12)
C6	0.270 (6)	0.082 (3)	0.058 (2)	-0.033 (3)	0.035 (3)	0.0098 (19)

Geometric parameters (\AA , $^\circ$)

C11—C10	1.736 (3)	C7—H7A	0.9600
S1—C1	1.690 (2)	C7—H7B	0.9600
N1—C1	1.323 (3)	C7—H7C	0.9600
N1—C4	1.466 (3)	C8—C9	1.372 (3)
N1—H1A	0.8541	C8—C13	1.380 (3)
N2—C1	1.364 (3)	C9—C10	1.382 (3)
N2—C2	1.424 (3)	C9—H9	0.9300
N2—C8	1.441 (3)	C10—C11	1.376 (4)
C2—C3	1.323 (4)	C11—C12	1.372 (4)
C2—C5	1.483 (4)	C11—H11	0.9300
C3—C4	1.480 (4)	C12—C13	1.381 (4)
C3—H3	0.9300	C12—H12	0.9300
C4—C6	1.518 (5)	C13—H13	0.9300
C4—C7	1.522 (4)	C6—H6A	0.9600
C5—H5A	0.9600	C6—H6B	0.9600
C5—H5B	0.9600	C6—H6C	0.9600
C5—H5C	0.9600		

C1—N1—C4	127.4 (2)	H7A—C7—H7B	109.5
C1—N1—H1A	112.2	C4—C7—H7C	109.5
C4—N1—H1A	119.1	H7A—C7—H7C	109.5
C1—N2—C2	121.35 (19)	H7B—C7—H7C	109.5
C1—N2—C8	118.85 (18)	C9—C8—C13	121.0 (2)
C2—N2—C8	119.75 (18)	C9—C8—N2	120.2 (2)
N1—C1—N2	117.2 (2)	C13—C8—N2	118.8 (2)
N1—C1—S1	121.01 (17)	C8—C9—C10	118.8 (2)
N2—C1—S1	121.78 (15)	C8—C9—H9	120.6
C3—C2—N2	118.9 (2)	C10—C9—H9	120.6
C3—C2—C5	123.9 (2)	C11—C10—C9	121.3 (2)
N2—C2—C5	117.0 (2)	C11—C10—C11	119.56 (19)
C2—C3—C4	124.0 (2)	C9—C10—C11	119.16 (19)
C2—C3—H3	118.0	C12—C11—C10	118.8 (2)
C4—C3—H3	118.0	C12—C11—H11	120.6
N1—C4—C3	107.8 (2)	C10—C11—H11	120.6
N1—C4—C6	108.5 (2)	C11—C12—C13	121.0 (2)
C3—C4—C6	111.6 (3)	C11—C12—H12	119.5
N1—C4—C7	107.1 (3)	C13—C12—H12	119.5
C3—C4—C7	111.2 (3)	C8—C13—C12	119.0 (2)
C6—C4—C7	110.5 (3)	C8—C13—H13	120.5
C2—C5—H5A	109.5	C12—C13—H13	120.5
C2—C5—H5B	109.5	C4—C6—H6A	109.5
H5A—C5—H5B	109.5	C4—C6—H6B	109.5
C2—C5—H5C	109.5	H6A—C6—H6B	109.5
H5A—C5—H5C	109.5	C4—C6—H6C	109.5
H5B—C5—H5C	109.5	H6A—C6—H6C	109.5
C4—C7—H7A	109.5	H6B—C6—H6C	109.5
C4—C7—H7B	109.5		
C4—N1—C1—N2	-8.0 (4)	C2—C3—C4—C6	100.9 (4)
C4—N1—C1—S1	173.1 (2)	C2—C3—C4—C7	-135.3 (3)
C2—N2—C1—N1	-7.0 (3)	C1—N2—C8—C9	88.4 (3)
C8—N2—C1—N1	170.6 (2)	C2—N2—C8—C9	-94.0 (3)
C2—N2—C1—S1	171.87 (19)	C1—N2—C8—C13	-92.8 (3)
C8—N2—C1—S1	-10.5 (3)	C2—N2—C8—C13	84.9 (3)
C1—N2—C2—C3	7.4 (4)	C13—C8—C9—C10	1.5 (3)
C8—N2—C2—C3	-170.1 (3)	N2—C8—C9—C10	-179.65 (19)
C1—N2—C2—C5	-168.4 (3)	C8—C9—C10—C11	-1.0 (3)
C8—N2—C2—C5	14.1 (4)	C8—C9—C10—C11	179.51 (17)
N2—C2—C3—C4	6.8 (4)	C9—C10—C11—C12	-0.3 (4)
C5—C2—C3—C4	-177.7 (3)	C11—C10—C11—C12	179.2 (2)
C1—N1—C4—C3	19.4 (4)	C10—C11—C12—C13	1.1 (4)
C1—N1—C4—C6	-101.6 (3)	C9—C8—C13—C12	-0.7 (4)
C1—N1—C4—C7	139.1 (3)	N2—C8—C13—C12	-179.6 (2)
C2—C3—C4—N1	-18.2 (4)	C11—C12—C13—C8	-0.6 (4)

Hydrogen-bond geometry (Å, °)

<i>D</i> —H··· <i>A</i>	<i>D</i> —H	H··· <i>A</i>	<i>D</i> ··· <i>A</i>	<i>D</i> —H··· <i>A</i>
-------------------------	-------------	---------------	-----------------------	-------------------------

supplementary materials

N1—H1A···S1ⁱ

0.85

2.58

3.404 (2)

162

Symmetry codes: (i) $-x+2, -y+2, -z+1$.

Fig. 1




Fig. 2

