

Acta Crystallographica Section E

Structure Reports

Online

ISSN 1600-5368

3-Chloro-*N'*-(2-methoxynaphthalen-1-yl)methylidene]benzohydrazide

Tian-Yi Li* and Yan-Xia Ge

School of Chemical Engineering, Changchun University of Technology, Changchun 130012, People's Republic of China

Correspondence e-mail: cooperationwell@126.com


Received 31 December 2010; accepted 6 January 2011

Key indicators: single-crystal X-ray study; $T = 298$ K; mean $\sigma(\text{C}-\text{C}) = 0.005$ Å; R factor = 0.060; wR factor = 0.147; data-to-parameter ratio = 16.2.

The title compound, $\text{C}_{19}\text{H}_{15}\text{ClN}_2\text{O}_2$, was prepared by the reaction of 2-methoxy-1-naphthaldehyde with 3-chlorobenzohydrazide in methanol. The dihedral angle between the benzene ring and the naphthyl ring system is $69.0(3)^\circ$. In the crystal, intermolecular $\text{N}-\text{H}\cdots\text{O}$ hydrogen bonds link the molecules into chains along the c axis. The crystal packing exhibits $\pi-\pi$ interactions, as indicated by distances of $3.768(3)$ Å between the centroids of the naphthyl rings of neighbouring molecules.

Related literature

For a related structure, see: Li & Li (2011). For reference bond lengths, see: Allen *et al.* (1987). For details of the synthesis, see: Zhu (2010).



Experimental

Crystal data

$\text{C}_{19}\text{H}_{15}\text{ClN}_2\text{O}_2$

$M_r = 338.78$

Monoclinic, $P2_1/c$
 $a = 12.181(2)$ Å
 $b = 16.953(4)$ Å
 $c = 8.5482(15)$ Å
 $\beta = 109.446(2)^\circ$
 $V = 1664.5(6)$ Å³

$Z = 4$
 Mo $K\alpha$ radiation
 $\mu = 0.24$ mm⁻¹
 $T = 298$ K
 $0.18 \times 0.18 \times 0.16$ mm

Data collection

Bruker APEXII CCD area-detector diffractometer
 Absorption correction: multi-scan (SADABS; Bruker, 2005)
 $T_{\min} = 0.958$, $T_{\max} = 0.962$

8964 measured reflections
 3586 independent reflections
 1624 reflections with $I > 2\sigma(I)$
 $R_{\text{int}} = 0.067$

Refinement

$R[F^2 > 2\sigma(F^2)] = 0.060$
 $wR(F^2) = 0.147$
 $S = 1.00$
 3586 reflections
 221 parameters
 1 restraint

H atoms treated by a mixture of independent and constrained refinement
 $\Delta\rho_{\text{max}} = 0.19$ e Å⁻³
 $\Delta\rho_{\text{min}} = -0.24$ e Å⁻³

Table 1

Hydrogen-bond geometry (Å, °).

$D-H\cdots A$	$D-H$	$H\cdots A$	$D\cdots A$	$D-H\cdots A$
$\text{N2}-\text{H2}\cdots\text{O2}^i$	0.91 (3)	2.04 (3)	2.937 (3)	170 (3)

Symmetry code: (i) $x, -y + \frac{1}{2}, z + \frac{1}{2}$.

Data collection: APEX2 (Bruker, 2005); cell refinement: SAINT (Bruker, 2005); data reduction: SAINT; program(s) used to solve structure: SHELXTL (Sheldrick, 2008); program(s) used to refine structure: SHELXTL; molecular graphics: SHELXTL; software used to prepare material for publication: SHELXTL.

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: CV5034).

References

- Allen, F. H., Kennard, O., Watson, D. G., Brammer, L., Orpen, A. G. & Taylor, R. (1987). *J. Chem. Soc. Perkin Trans. 2*, pp. S1–19.
 Bruker (2005). APEX2, SAINT and SADABS. Bruker AXS Inc., Madison, Wisconsin, USA.
 Li, T.-Y. & Li, W. (2011). *Acta Cryst. E67*, o373.
 Sheldrick, G. M. (2008). *Acta Cryst. A64*, 112–122.
 Zhu, H.-Y. (2010). *Acta Cryst. E66*, o2562.

supplementary materials

Acta Cryst. (2011). E67, o374 [doi:10.1107/S1600536811000924]

3-Chloro-*N'*-(2-methoxynaphthalen-1-yl)methylidene]benzohydrazide

T.-Y. Li and Y.-X. Ge

Comment

In continuation of our structural study of naphthylidenebenzohydrazide derivatives (Li & Li, 2011) we present here the title compound (I).

In (I) (Fig. 1), the dihedral angle between the benzene ring and the naphthyl bicycle is 69.0 (3)°. All the bond lengths are within normal values (Allen *et al.*, 1987). Intermolecular N—H···O hydrogen bonds (Table 1) link the molecules into chains along the *c* axis (Fig. 2). The crystal packing exhibits π - π interactions proved by short distances of 3.768 (3) Å between the centroids of naphthyl rings from the neighbouring molecules.

Experimental

The compound was prepared and crystallized according to the literature method (Zhu, 2010). 2-Methoxy-1-naphthaldehyde (0.186 g, 1 mmol) and 3-chlorobenzohydrazide (0.171 g, 1 mmol) were dissolved in 30 ml absolute methanol. The mixture was stirred at reflux for 10 min, and cooled to room temperature. The clear colorless solution was left to slow evaporation in air for eight days, yielding colorless block-shaped crystals, which were collected by filtration and washed with methanol.

Refinement

The amino H atom was located from a difference Fourier map and refined isotropically, with the N—H distance restrained to 0.90 (1) Å. The other H atoms were positioned geometrically and refined using the riding-model approximation, with C—H = 0.93–0.96 Å, and $U_{\text{iso}}(\text{H}) = 1.2U_{\text{eq}}(\text{C})$ or $U_{\text{iso}}(\text{H}) = 1.5U_{\text{eq}}(\text{C19})$.

Figures




Fig. 1. The molecular structure of the title compound with 30% probability displacement ellipsoids for non-hydrogen atoms.




Fig. 2. A portion of the crystal packing showing hydrogen bonds as dashed lines.

3-Chloro-*N*'-[(2-methoxynaphthalen-1-yl)methylidene]benzohydrazide

Crystal data

$C_{19}H_{15}ClN_2O_2$	$F(000) = 704$
$M_r = 338.78$	$D_x = 1.352 \text{ Mg m}^{-3}$
Monoclinic, $P2_1/c$	Mo $K\alpha$ radiation, $\lambda = 0.71073 \text{ \AA}$
$a = 12.181 (2) \text{ \AA}$	Cell parameters from 762 reflections
$b = 16.953 (4) \text{ \AA}$	$\theta = 2.5\text{--}24.3^\circ$
$c = 8.5482 (15) \text{ \AA}$	$\mu = 0.24 \text{ mm}^{-1}$
$\beta = 109.446 (2)^\circ$	$T = 298 \text{ K}$
$V = 1664.5 (6) \text{ \AA}^3$	Block, colourless
$Z = 4$	$0.18 \times 0.18 \times 0.16 \text{ mm}$

Data collection

Bruker APEXII CCD area-detector diffractometer	3586 independent reflections
Radiation source: fine-focus sealed tube graphite	1624 reflections with $I > 2\sigma(I)$
ω scans	$R_{\text{int}} = 0.067$
Absorption correction: multi-scan (<i>SADABS</i> ; Bruker, 2005)	$\theta_{\text{max}} = 27.0^\circ$, $\theta_{\text{min}} = 2.4^\circ$
$T_{\text{min}} = 0.958$, $T_{\text{max}} = 0.962$	$h = -14 \rightarrow 15$
8964 measured reflections	$k = -19 \rightarrow 21$
	$l = -10 \rightarrow 10$

Refinement

Refinement on F^2	Primary atom site location: structure-invariant direct methods
Least-squares matrix: full	Secondary atom site location: difference Fourier map
$R[F^2 > 2\sigma(F^2)] = 0.060$	Hydrogen site location: inferred from neighbouring sites
$wR(F^2) = 0.147$	H atoms treated by a mixture of independent and constrained refinement
$S = 1.00$	$w = 1/[\sigma^2(F_o^2) + (0.0386P)^2]$
3586 reflections	where $P = (F_o^2 + 2F_c^2)/3$
221 parameters	$(\Delta/\sigma)_{\text{max}} < 0.001$
1 restraint	$\Delta\rho_{\text{max}} = 0.19 \text{ e \AA}^{-3}$
	$\Delta\rho_{\text{min}} = -0.24 \text{ e \AA}^{-3}$

Special details

Geometry. All e.s.d.'s (except the e.s.d. in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell e.s.d.'s are taken into account individually in the estimation of e.s.d.'s in distances, angles and torsion angles; correlations between e.s.d.'s in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell e.s.d.'s is used for estimating e.s.d.'s involving l.s. planes.

Refinement. Refinement of F^2 against ALL reflections. The weighted R -factor wR and goodness of fit S are based on F^2 , conventional R -factors R are based on F , with F set to zero for negative F^2 . The threshold expression of $F^2 > \sigma(F^2)$ is used only for calculating R -factors(gt) *etc.* and is not relevant to the choice of reflections for refinement. R -factors based on F^2 are statistically about twice as large as those based on F , and R -factors based on ALL data will be even larger.

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (\AA^2)

	x	y	z	$U_{\text{iso}}^*/U_{\text{eq}}$
C11	0.57105 (8)	0.10288 (6)	1.03193 (12)	0.0793 (4)
N1	0.8646 (2)	0.28050 (15)	0.6055 (3)	0.0469 (7)
N2	0.7753 (2)	0.24546 (15)	0.6487 (3)	0.0467 (7)
O1	0.9252 (2)	0.49223 (13)	0.7763 (3)	0.0709 (7)
O2	0.71701 (19)	0.17210 (12)	0.4129 (3)	0.0619 (7)
C1	1.0045 (3)	0.38640 (17)	0.6734 (4)	0.0453 (8)
C2	1.0122 (3)	0.46507 (19)	0.7236 (4)	0.0540 (9)
C3	1.1046 (3)	0.5131 (2)	0.7182 (4)	0.0707 (11)
H3	1.1088	0.5653	0.7527	0.085*
C4	1.1874 (3)	0.4831 (2)	0.6626 (5)	0.0753 (11)
H4	1.2479	0.5157	0.6589	0.090*
C5	1.1857 (3)	0.4049 (2)	0.6107 (4)	0.0563 (9)
C6	1.2739 (3)	0.3739 (2)	0.5573 (5)	0.0737 (11)
H6	1.3347	0.4067	0.5556	0.088*
C7	1.2739 (3)	0.2977 (3)	0.5082 (5)	0.0751 (11)
H7	1.3337	0.2785	0.4739	0.090*
C8	1.1819 (3)	0.2489 (2)	0.5103 (4)	0.0676 (10)
H8	1.1807	0.1965	0.4773	0.081*
C9	1.0942 (3)	0.27685 (19)	0.5598 (4)	0.0549 (9)
H9	1.0337	0.2431	0.5582	0.066*
C10	1.0917 (3)	0.35562 (17)	0.6136 (4)	0.0458 (8)
C11	0.9075 (3)	0.34138 (18)	0.6916 (4)	0.0488 (8)
H11	0.8752	0.3580	0.7703	0.059*
C12	0.7072 (3)	0.19108 (18)	0.5452 (4)	0.0463 (8)
C13	0.6157 (3)	0.15628 (16)	0.6049 (4)	0.0418 (7)
C14	0.6300 (2)	0.14888 (16)	0.7713 (4)	0.0433 (8)
H14	0.6972	0.1682	0.8504	0.052*
C15	0.5463 (3)	0.11327 (18)	0.8211 (4)	0.0506 (8)
C16	0.4458 (3)	0.0859 (2)	0.7070 (5)	0.0752 (11)
H16	0.3886	0.0626	0.7417	0.090*
C17	0.4298 (3)	0.0930 (2)	0.5414 (5)	0.0840 (12)
H17	0.3618	0.0743	0.4633	0.101*
C18	0.5144 (3)	0.1278 (2)	0.4901 (4)	0.0670 (10)
H18	0.5031	0.1322	0.3774	0.080*
C19	0.9307 (4)	0.5719 (2)	0.8359 (5)	0.0905 (13)
H19A	1.0029	0.5798	0.9245	0.136*
H19B	0.8669	0.5812	0.8758	0.136*
H19C	0.9260	0.6079	0.7471	0.136*
H2	0.757 (3)	0.265 (2)	0.736 (3)	0.109*

supplementary materials

Atomic displacement parameters (\AA^2)

	U^{11}	U^{22}	U^{33}	U^{12}	U^{13}	U^{23}
C11	0.0826 (7)	0.1061 (8)	0.0635 (7)	-0.0240 (6)	0.0433 (6)	0.0039 (5)
N1	0.0513 (16)	0.0548 (16)	0.0406 (16)	-0.0094 (13)	0.0231 (14)	-0.0007 (13)
N2	0.0505 (16)	0.0514 (16)	0.0461 (18)	-0.0066 (13)	0.0267 (15)	-0.0026 (13)
O1	0.0942 (18)	0.0526 (15)	0.0704 (18)	0.0061 (13)	0.0333 (16)	-0.0040 (12)
O2	0.0833 (17)	0.0664 (15)	0.0490 (15)	-0.0153 (12)	0.0392 (14)	-0.0103 (11)
C1	0.055 (2)	0.0434 (19)	0.0354 (19)	-0.0063 (15)	0.0130 (17)	0.0031 (14)
C2	0.066 (2)	0.050 (2)	0.041 (2)	-0.0017 (18)	0.0120 (18)	0.0071 (16)
C3	0.087 (3)	0.047 (2)	0.067 (3)	-0.015 (2)	0.010 (2)	-0.0055 (18)
C4	0.070 (3)	0.067 (3)	0.081 (3)	-0.030 (2)	0.015 (2)	0.002 (2)
C5	0.054 (2)	0.065 (2)	0.047 (2)	-0.0132 (17)	0.0121 (18)	0.0083 (18)
C6	0.055 (2)	0.091 (3)	0.080 (3)	-0.014 (2)	0.029 (2)	0.008 (2)
C7	0.056 (2)	0.094 (3)	0.085 (3)	0.004 (2)	0.037 (2)	0.006 (2)
C8	0.068 (2)	0.075 (2)	0.068 (3)	-0.001 (2)	0.034 (2)	-0.001 (2)
C9	0.051 (2)	0.064 (2)	0.054 (2)	-0.0092 (17)	0.0235 (19)	0.0010 (17)
C10	0.053 (2)	0.0466 (19)	0.0367 (19)	-0.0077 (15)	0.0138 (17)	0.0048 (15)
C11	0.052 (2)	0.055 (2)	0.042 (2)	-0.0037 (16)	0.0195 (17)	0.0020 (17)
C12	0.053 (2)	0.0482 (19)	0.044 (2)	0.0026 (16)	0.0239 (18)	0.0015 (16)
C13	0.0424 (19)	0.0426 (17)	0.042 (2)	0.0016 (14)	0.0169 (17)	-0.0018 (15)
C14	0.0440 (18)	0.0436 (17)	0.047 (2)	-0.0031 (14)	0.0221 (17)	-0.0027 (15)
C15	0.048 (2)	0.058 (2)	0.055 (2)	-0.0020 (16)	0.0298 (19)	0.0024 (17)
C16	0.049 (2)	0.098 (3)	0.086 (3)	-0.019 (2)	0.032 (2)	0.003 (2)
C17	0.055 (2)	0.126 (4)	0.065 (3)	-0.029 (2)	0.012 (2)	-0.007 (3)
C18	0.064 (2)	0.087 (3)	0.049 (2)	-0.010 (2)	0.018 (2)	0.0005 (19)
C19	0.141 (4)	0.059 (2)	0.065 (3)	0.024 (2)	0.025 (3)	-0.007 (2)

Geometric parameters (\AA , $^\circ$)

C11—C15	1.735 (3)	C7—H7	0.9300
N1—C11	1.274 (3)	C8—C9	1.358 (4)
N1—N2	1.392 (3)	C8—H8	0.9300
N2—C12	1.354 (4)	C9—C10	1.416 (4)
N2—H2	0.91 (3)	C9—H9	0.9300
O1—C2	1.362 (4)	C11—H11	0.9300
O1—C19	1.437 (4)	C12—C13	1.494 (4)
O2—C12	1.219 (3)	C13—C14	1.380 (4)
C1—C2	1.394 (4)	C13—C18	1.383 (4)
C1—C10	1.422 (4)	C14—C15	1.370 (4)
C1—C11	1.458 (4)	C14—H14	0.9300
C2—C3	1.402 (4)	C15—C16	1.367 (4)
C3—C4	1.349 (4)	C16—C17	1.368 (5)
C3—H3	0.9300	C16—H16	0.9300
C4—C5	1.395 (5)	C17—C18	1.379 (5)
C4—H4	0.9300	C17—H17	0.9300
C5—C6	1.402 (5)	C18—H18	0.9300
C5—C10	1.425 (4)	C19—H19A	0.9600

C6—C7	1.358 (5)	C19—H19B	0.9600
C6—H6	0.9300	C19—H19C	0.9600
C7—C8	1.399 (5)		
C11—N1—N2	113.5 (2)	C9—C10—C5	116.3 (3)
C12—N2—N1	118.7 (2)	C1—C10—C5	119.5 (3)
C12—N2—H2	121 (2)	N1—C11—C1	123.7 (3)
N1—N2—H2	119 (2)	N1—C11—H11	118.2
C2—O1—C19	119.1 (3)	C1—C11—H11	118.2
C2—C1—C10	118.7 (3)	O2—C12—N2	123.9 (3)
C2—C1—C11	116.2 (3)	O2—C12—C13	121.8 (3)
C10—C1—C11	125.1 (3)	N2—C12—C13	114.2 (3)
O1—C2—C1	116.4 (3)	C14—C13—C18	118.5 (3)
O1—C2—C3	122.4 (3)	C14—C13—C12	122.4 (3)
C1—C2—C3	121.1 (3)	C18—C13—C12	119.1 (3)
C4—C3—C2	119.7 (3)	C15—C14—C13	120.6 (3)
C4—C3—H3	120.1	C15—C14—H14	119.7
C2—C3—H3	120.1	C13—C14—H14	119.7
C3—C4—C5	122.4 (3)	C16—C15—C14	120.7 (3)
C3—C4—H4	118.8	C16—C15—Cl1	120.8 (2)
C5—C4—H4	118.8	C14—C15—Cl1	118.5 (3)
C4—C5—C6	121.9 (3)	C15—C16—C17	119.5 (3)
C4—C5—C10	118.6 (3)	C15—C16—H16	120.2
C6—C5—C10	119.5 (3)	C17—C16—H16	120.2
C7—C6—C5	122.4 (3)	C16—C17—C18	120.2 (3)
C7—C6—H6	118.8	C16—C17—H17	119.9
C5—C6—H6	118.8	C18—C17—H17	119.9
C6—C7—C8	118.4 (3)	C17—C18—C13	120.5 (3)
C6—C7—H7	120.8	C17—C18—H18	119.7
C8—C7—H7	120.8	C13—C18—H18	119.7
C9—C8—C7	121.0 (3)	O1—C19—H19A	109.5
C9—C8—H8	119.5	O1—C19—H19B	109.5
C7—C8—H8	119.5	H19A—C19—H19B	109.5
C8—C9—C10	122.3 (3)	O1—C19—H19C	109.5
C8—C9—H9	118.9	H19A—C19—H19C	109.5
C10—C9—H9	118.9	H19B—C19—H19C	109.5
C9—C10—C1	124.2 (3)		

Hydrogen-bond geometry (Å, °)

<i>D</i> —H \cdots <i>A</i>	<i>D</i> —H	H \cdots <i>A</i>	<i>D</i> \cdots <i>A</i>	<i>D</i> —H \cdots <i>A</i>
N2—H2 \cdots O2 ⁱ	0.91 (3)	2.04 (3)	2.937 (3)	170 (3)

Symmetry codes: (i) *x*, $-y+1/2$, $z+1/2$.

Fig. 1




Fig. 2

