

2-[1-(4-Chlorobenzoyl)pyrrolidin-2-yl]-4,4,5,5-tetramethyl-4,5-dihydro-imidazole-1-oxyl-3-oxide

Min Tian,^a Zhuo Xiang,^a Si-Yuan Zhou,^b Lin-Lin Jing,^c
Hai-Bo Wang^a and Xiao-Li Sun^{a*}

^aDepartment of Chemistry, School of Pharmacy, Fourth Military Medical University, Changle West Road 17, 710032 Xi-An, People's Republic of China, ^bDepartment of Pharmaceutics, School of Pharmacy, Fourth Military Medical University, Changle West Road 17, 710032 Xi-An, People's Republic of China, and ^cDepartment of Pharmacy, Lanzhou General Hospital, Lanzhou Command, Lanzhou 730050, People's Republic of China

Correspondence e-mail: xiaoli_sun@yahoo.cn


Received 17 December 2010; accepted 11 January 2011

Key indicators: single-crystal X-ray study; $T = 296\text{ K}$; mean $\sigma(\text{C}-\text{C}) = 0.005\text{ \AA}$; R factor = 0.051; wR factor = 0.152; data-to-parameter ratio = 14.5.

In the title compound, $\text{C}_{18}\text{H}_{23}\text{ClN}_3\text{O}_3$, the imidazole ring system has an envelope conformation, whereas the nitronyl nitroxide unit displays a half-chair or twisted conformation. In the crystal, $\text{C}-\text{H}\cdots\text{O}$ hydrogen bonds build up a three-dimensional network.

Related literature

For the biological activity of nitronyl nitroxides, see: Soule *et al.* (2007) and for their coordination properties, see: Masuda *et al.* (2009). For related structures, see: Iqbal *et al.* (2009); Wang *et al.* (2009). For puckering parameters, see: Cremer & Pople (1975).



Experimental

Crystal data

$\text{C}_{18}\text{H}_{23}\text{ClN}_3\text{O}_3$
 $M_r = 364.84$

Monoclinic, $P2_1/n$
 $a = 11.6202 (19)\text{ \AA}$

$b = 8.2694 (13)\text{ \AA}$
 $c = 20.315 (3)\text{ \AA}$
 $\beta = 105.636 (2)^\circ$
 $V = 1879.8 (5)\text{ \AA}^3$
 $Z = 4$

Mo $K\alpha$ radiation
 $\mu = 0.23\text{ mm}^{-1}$
 $T = 296\text{ K}$
 $0.34 \times 0.29 \times 0.18\text{ mm}$

Data collection

Bruker APEXII CCD
diffractometer
Absorption correction: multi-scan
(*SADABS*; Sheldrick, 2008a)
 $R_{\text{min}} = 0.928$, $T_{\text{max}} = 0.960$

9089 measured reflections
3342 independent reflections
1791 reflections with $I > 2\sigma(I)$
 $R_{\text{int}} = 0.051$

Refinement

$R[F^2 > 2\sigma(F^2)] = 0.051$
 $wR(F^2) = 0.152$
 $S = 1.03$
3342 reflections

230 parameters
H-atom parameters constrained
 $\Delta\rho_{\text{max}} = 0.19\text{ e \AA}^{-3}$
 $\Delta\rho_{\text{min}} = -0.18\text{ e \AA}^{-3}$

Table 1
Hydrogen-bond geometry (\AA , $^\circ$).

$D-\text{H}\cdots A$	$D-\text{H}$	$\text{H}\cdots A$	$D\cdots A$	$D-\text{H}\cdots A$
C2—H2A \cdots O2 ⁱ	0.93	2.54	3.458 (4)	170
C3—H3A \cdots O1 ⁱ	0.93	2.45	3.314 (4)	154
C8—H8A \cdots O3 ⁱⁱ	0.97	2.44	3.101 (4)	125

Symmetry codes: (i) $-x + 1, -y, -z$; (ii) $-x + \frac{3}{2}, y - \frac{1}{2}, -z + \frac{1}{2}$.

Data collection: *APEX2* (Bruker, 2007); cell refinement: *SAINT* (Bruker, 2007); data reduction: *SAINT*; program(s) used to solve structure: *SHELXS97* (Sheldrick, 2008b); program(s) used to refine structure: *SHELXL97* (Sheldrick, 2008b); molecular graphics: *ORTEPIII* (Burnett & Johnson, 1996) and *ORTEP-3 for Windows* (Farrugia, 1997); software used to prepare material for publication: *SHELXTL* (Sheldrick, 2008b) and *PLATON* (Spek, 2009).

We thank the Natural Science Foundation of China (grant Nos. 81001398, 30901883, 20802091) for financial support.

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: DN2639).

References

- Bruker (2007). *APEX2* and *SAINT*. Bruker AXS Inc., Madison, Wisconsin, USA.
- Burnett, M. N. & Johnson, C. K. (1996). *ORTEPIII*. Report ORNL-6895. Oak Ridge National Laboratory, Tennessee, USA.
- Cremer, D. & Pople, J. A. (1975). *J. Am. Chem. Soc.* **97**, 1354–1358.
- Farrugia, L. J. (1997). *J. Appl. Cryst.* **30**, 565.
- Iqbal, A. L., Anirban, P. & Sambhu, N. D. (2009). *J. Phys. Chem. A*, **113**, 1595–4673.
- Masuda, Y., Kurats, M., Suzuki, S., Kozaki, M., Shiomi, D., Sato, K., Takui, T., Hosokoshi, Y., Miyazaki, Y., Inada, A. & Okada, K. (2009). *J. Am. Chem. Soc.* **131**, 4670–4673.
- Sheldrick, G. M. (2008a). *SADABS*. University of Göttingen, Germany.
- Sheldrick, G. M. (2008b). *Acta Cryst. A* **64**, 112–122.
- Soule, B. P., Hyodo, F., Matsumoto, K., Simone, N. L., Cook, J. A., Krishna, M. C. & Mitchell, J. B. (2007). *Free Radic. Biol. Med.* **42**, 1632–1650.
- Spek, A. L. (2009). *Acta Cryst. D* **65**, 148–155.
- Wang, H.-B., Jing, L.-L., Gao, P. & Sun, X.-L. (2009). *Acta Cryst. E* **65**, o2090.

supplementary materials

Acta Cryst. (2011). E67, o425 [doi:10.1107/S1600536811001462]

2-[1-(4-Chlorobenzoyl)pyrrolidin-2-yl]-4,4,5,5-tetramethyl-4,5-dihydroimidazole-1-oxyl-3-oxide

M. Tian, Z. Xiang, S.-Y. Zhou, L.-L. Jing, H.-B. Wang and X.-L. Sun

Comment

Nitronyl nitroxides, stable organic radicals, display interesting properties in many fields as magnetism, anticancer, antiradiation and antioxidation *etc* (Soule *et al.*, 2007). Nitronyl nitroxides have received considerable attention recently (Iqbal *et al.*, 2009; Wang *et al.*, 2009). The title compound can also be used for coordination with many metal cations, such as Mn²⁺, Cu²⁺ and Ni²⁺ leading to some molecule based magnetic materials (Masuda, *et al.*, 2009).

In the title compound, the indole ring system has an envelope conformation with puckering parameters Q(2)= 0.370 (4) Å and φ= 78.0 (5)° (Cremer & Pople, 1975) whereas the nitronyl nitroxide moiety displays a half-chair or twisted conformation with Q(2)= 0.228 (3) Å and φ= 308.6 (7)°. Occurrence of C-H···O hydrogen bonds build up a three dimensional network (Table 1).

Experimental

2,3-Dimethyl-2,3-bis(hydroxylamino)butane (1.48 g, 10.0 mmol) and 1-(4-chlorobenzoyl)pyrrolidine-2-carbaldehyde (2.38 g, 10.0 mmol) were dissolved in methanol. The reaction was stirred for 10 h at reflux temperature, then cooled to room temperature and filtered. The cake was suspended in dichloromethane (150.0 ml) and cooled at ice bath for 10 min, Then the reaction mixture was added to an aqueous solution of NaIO₄ and stirred for 15 min to give a amaranthine solution. The aqueous phase was extracted with CH₂Cl₂ and the organic layer was combined and dried over MgSO₄. Then the solvent was removed to give a amaranthine residue which was purified by a flash column chromatography with the elution of *n*-hexane/ethyl acetate (1:1) to yield the title compound (I) as a dark amaranthine powder. Single crystals of compound (I) were obtained from the 1/1 mixed solution of *n*-heptane and dichloromethane.

Refinement

All H atoms attached to C atoms were fixed geometrically and treated as riding with C—H = 0.96 Å (methyl), 0.97 Å (methylene) and 0.93 Å (aromatic) with U_{iso}(H) = 1.2U_{eq}(C) or U_{iso}(H) = 1.5U_{eq}(Cmethyl).

Figures




Fig. 1. Molecular structure of the title compound (I), showing the atom labeling scheme. Displacement ellipsoids are drawn at the 30% probability level. H atoms are represented as small spheres of arbitrary radii.

supplementary materials

2-[1-(4-Chlorobenzoyl)pyrrolidin-2-yl]-4,4,5-tetramethyl-4,5-dihydroimidazole-1-oxyl-3-oxide

Crystal data

C ₁₈ H ₂₃ ClN ₃ O ₃	<i>F</i> (000) = 772
<i>M_r</i> = 364.84	<i>D_x</i> = 1.289 Mg m ⁻³
Monoclinic, <i>P2₁/n</i>	Mo <i>Kα</i> radiation, λ = 0.71073 Å
Hall symbol: -P 2yn	Cell parameters from 1006 reflections
<i>a</i> = 11.6202 (19) Å	θ = 2.3–19.1°
<i>b</i> = 8.2694 (13) Å	μ = 0.23 mm ⁻¹
<i>c</i> = 20.315 (3) Å	<i>T</i> = 296 K
β = 105.636 (2)°	Block, purple
<i>V</i> = 1879.8 (5) Å ³	0.34 × 0.29 × 0.18 mm
<i>Z</i> = 4	

Data collection

Bruker APEXII CCD diffractometer	3342 independent reflections
Radiation source: fine-focus sealed tube graphite	1791 reflections with $I > 2\sigma(I)$
φ and ω scans	$R_{\text{int}} = 0.051$
Absorption correction: multi-scan (<i>SADABS</i> ; Sheldrick, 2008a)	$\theta_{\text{max}} = 25.1^\circ$, $\theta_{\text{min}} = 1.8^\circ$
$T_{\text{min}} = 0.928$, $T_{\text{max}} = 0.960$	$h = -13 \rightarrow 13$
9089 measured reflections	$k = -9 \rightarrow 5$
	$l = -24 \rightarrow 22$

Refinement

Refinement on F^2	Primary atom site location: structure-invariant direct methods
Least-squares matrix: full	Secondary atom site location: difference Fourier map
$R[F^2 > 2\sigma(F^2)] = 0.051$	Hydrogen site location: inferred from neighbouring sites
$wR(F^2) = 0.152$	H-atom parameters constrained
$S = 1.03$	$w = 1/[\sigma^2(F_o^2) + (0.0548P)^2]$ where $P = (F_o^2 + 2F_c^2)/3$
3342 reflections	$(\Delta/\sigma)_{\text{max}} < 0.001$
230 parameters	$\Delta\rho_{\text{max}} = 0.19 \text{ e \AA}^{-3}$
0 restraints	$\Delta\rho_{\text{min}} = -0.18 \text{ e \AA}^{-3}$

Special details

Geometry. All esds (except the esd in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell esds are taken into account individually in the estimation of esds in distances, angles and torsion angles; correlations between esds

in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell esds is used for estimating esds involving l.s. planes.

Refinement. Refinement of F^2 against ALL reflections. The weighted R -factor wR and goodness of fit S are based on F^2 , conventional R -factors R are based on F , with F set to zero for negative F^2 . The threshold expression of $F^2 > \sigma(F^2)$ is used only for calculating R -factors(gt) etc. and is not relevant to the choice of reflections for refinement. R -factors based on F^2 are statistically about twice as large as those based on F , and R -factors based on ALL data will be even larger.

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (\AA^2)

	<i>x</i>	<i>y</i>	<i>z</i>	$U_{\text{iso}}^*/U_{\text{eq}}$
Cl1	0.24418 (8)	-0.02934 (14)	0.22799 (5)	0.0934 (4)
O1	0.61663 (17)	0.1756 (3)	0.04529 (10)	0.0602 (6)
O2	0.84803 (18)	0.2887 (2)	-0.03089 (10)	0.0613 (6)
O3	0.7771 (2)	0.5304 (3)	0.15800 (11)	0.0793 (8)
N1	0.7572 (2)	0.1723 (3)	0.14502 (11)	0.0496 (6)
N2	0.81945 (18)	0.3955 (3)	0.00771 (11)	0.0462 (6)
N3	0.7868 (2)	0.5102 (3)	0.09729 (12)	0.0522 (7)
C1	0.3608 (3)	0.0241 (4)	0.19376 (17)	0.0586 (9)
C2	0.3684 (3)	-0.0441 (4)	0.13384 (17)	0.0601 (9)
H2A	0.3113	-0.1182	0.1111	0.072*
C3	0.4613 (3)	-0.0021 (3)	0.10737 (15)	0.0537 (8)
H3A	0.4661	-0.0471	0.0662	0.064*
C4	0.5484 (2)	0.1070 (3)	0.14140 (14)	0.0472 (7)
C5	0.5373 (3)	0.1743 (4)	0.20168 (15)	0.0561 (8)
H5	0.5942	0.2480	0.2250	0.067*
C6	0.4434 (3)	0.1341 (4)	0.22772 (16)	0.0609 (9)
H6	0.4362	0.1812	0.2680	0.073*
C7	0.6433 (3)	0.1542 (3)	0.10742 (15)	0.0479 (7)
C8	0.8104 (3)	0.1429 (4)	0.21806 (14)	0.0617 (9)
H8A	0.7748	0.0491	0.2335	0.074*
H8B	0.8008	0.2360	0.2451	0.074*
C9	0.9400 (3)	0.1137 (5)	0.22247 (16)	0.0697 (10)
H9A	0.9537	0.0020	0.2121	0.084*
H9B	0.9904	0.1401	0.2676	0.084*
C10	0.9636 (3)	0.2265 (4)	0.16896 (16)	0.0635 (9)
H10A	1.0315	0.1900	0.1537	0.076*
H10B	0.9787	0.3357	0.1866	0.076*
C11	0.8480 (2)	0.2188 (4)	0.11055 (14)	0.0486 (8)
H11	0.8553	0.1332	0.0786	0.058*
C12	0.8187 (2)	0.3736 (4)	0.07277 (13)	0.0438 (7)
C13	0.7659 (3)	0.5550 (4)	-0.01959 (15)	0.0510 (8)
C14	0.7768 (3)	0.6477 (4)	0.04813 (15)	0.0531 (8)
C15	0.8360 (3)	0.6285 (4)	-0.06555 (17)	0.0779 (11)
H15A	0.9181	0.6412	-0.0402	0.117*
H15B	0.8029	0.7322	-0.0816	0.117*
H15C	0.8311	0.5585	-0.1039	0.117*
C16	0.6380 (3)	0.5167 (4)	-0.05963 (18)	0.0816 (12)
H16A	0.6388	0.4451	-0.0967	0.122*

supplementary materials

H16B	0.5976	0.6150	-0.0775	0.122*
H16C	0.5971	0.4657	-0.0300	0.122*
C17	0.8923 (3)	0.7448 (4)	0.07346 (18)	0.0801 (11)
H17A	0.8999	0.7809	0.1193	0.120*
H17B	0.8901	0.8367	0.0443	0.120*
H17C	0.9593	0.6777	0.0727	0.120*
C18	0.6703 (3)	0.7508 (5)	0.05121 (19)	0.0856 (12)
H18A	0.5987	0.6872	0.0375	0.128*
H18B	0.6643	0.8415	0.0210	0.128*
H18C	0.6807	0.7887	0.0971	0.128*

Atomic displacement parameters (\AA^2)

	U^{11}	U^{22}	U^{33}	U^{12}	U^{13}	U^{23}
Cl1	0.0622 (6)	0.1224 (10)	0.1045 (8)	-0.0041 (5)	0.0375 (6)	0.0165 (6)
O1	0.0588 (13)	0.0744 (16)	0.0446 (13)	-0.0067 (11)	0.0091 (10)	0.0026 (10)
O2	0.0730 (14)	0.0614 (16)	0.0543 (13)	0.0069 (11)	0.0252 (11)	-0.0072 (11)
O3	0.114 (2)	0.0787 (18)	0.0516 (14)	0.0164 (14)	0.0341 (13)	-0.0040 (12)
N1	0.0472 (15)	0.0576 (17)	0.0428 (14)	-0.0007 (12)	0.0099 (12)	0.0103 (11)
N2	0.0456 (13)	0.0504 (16)	0.0447 (15)	0.0033 (12)	0.0155 (11)	-0.0006 (12)
N3	0.0622 (16)	0.0504 (17)	0.0464 (16)	0.0085 (12)	0.0185 (13)	0.0005 (13)
C1	0.0430 (18)	0.068 (2)	0.065 (2)	0.0065 (16)	0.0152 (16)	0.0131 (18)
C2	0.0481 (19)	0.052 (2)	0.075 (2)	-0.0038 (15)	0.0087 (17)	0.0014 (17)
C3	0.0515 (19)	0.053 (2)	0.0521 (19)	0.0012 (16)	0.0069 (16)	-0.0061 (15)
C4	0.0467 (17)	0.0432 (19)	0.0498 (18)	0.0038 (14)	0.0100 (14)	0.0042 (14)
C5	0.0522 (19)	0.059 (2)	0.055 (2)	-0.0026 (15)	0.0107 (16)	-0.0082 (16)
C6	0.055 (2)	0.075 (3)	0.0533 (19)	0.0059 (18)	0.0165 (16)	-0.0048 (17)
C7	0.0521 (19)	0.0430 (19)	0.0470 (18)	0.0001 (14)	0.0107 (15)	-0.0006 (14)
C8	0.058 (2)	0.077 (3)	0.0484 (19)	0.0023 (17)	0.0103 (15)	0.0142 (16)
C9	0.056 (2)	0.088 (3)	0.062 (2)	0.0021 (18)	0.0107 (16)	0.0231 (19)
C10	0.0503 (18)	0.070 (2)	0.067 (2)	0.0038 (16)	0.0088 (16)	0.0188 (17)
C11	0.0472 (17)	0.049 (2)	0.0495 (18)	0.0049 (14)	0.0131 (14)	0.0086 (14)
C12	0.0428 (16)	0.049 (2)	0.0401 (17)	0.0037 (14)	0.0116 (13)	0.0050 (15)
C13	0.0522 (18)	0.048 (2)	0.0488 (18)	0.0073 (15)	0.0074 (14)	0.0112 (14)
C14	0.0546 (19)	0.047 (2)	0.058 (2)	0.0111 (15)	0.0157 (15)	0.0076 (15)
C15	0.104 (3)	0.070 (3)	0.068 (2)	0.001 (2)	0.039 (2)	0.0161 (18)
C16	0.067 (2)	0.084 (3)	0.076 (2)	0.0130 (19)	-0.011 (2)	0.0042 (19)
C17	0.087 (3)	0.068 (3)	0.086 (3)	-0.015 (2)	0.024 (2)	-0.0091 (19)
C18	0.087 (3)	0.082 (3)	0.089 (3)	0.040 (2)	0.027 (2)	0.008 (2)

Geometric parameters (\AA , $^\circ$)

Cl1—C1	1.738 (3)	C9—C10	1.513 (4)
O1—C7	1.229 (3)	C9—H9A	0.9700
O2—N2	1.283 (3)	C9—H9B	0.9700
O3—N3	1.279 (3)	C10—C11	1.535 (4)
N1—C7	1.346 (3)	C10—H10A	0.9700
N1—C8	1.466 (3)	C10—H10B	0.9700
N1—C11	1.467 (3)	C11—C12	1.484 (4)

N2—C12	1.336 (3)	C11—H11	0.9800
N2—C13	1.500 (4)	C13—C15	1.521 (4)
N3—C12	1.328 (3)	C13—C16	1.523 (4)
N3—C14	1.497 (4)	C13—C14	1.550 (4)
C1—C2	1.366 (4)	C14—C18	1.517 (4)
C1—C6	1.367 (4)	C14—C17	1.529 (4)
C2—C3	1.374 (4)	C15—H15A	0.9600
C2—H2A	0.9300	C15—H15B	0.9600
C3—C4	1.392 (4)	C15—H15C	0.9600
C3—H3A	0.9300	C16—H16A	0.9600
C4—C5	1.382 (4)	C16—H16B	0.9600
C4—C7	1.502 (4)	C16—H16C	0.9600
C5—C6	1.376 (4)	C17—H17A	0.9600
C5—H5	0.9300	C17—H17B	0.9600
C6—H6	0.9300	C17—H17C	0.9600
C8—C9	1.503 (4)	C18—H18A	0.9600
C8—H8A	0.9700	C18—H18B	0.9600
C8—H8B	0.9700	C18—H18C	0.9600
C7—N1—C8	129.7 (2)	H10A—C10—H10B	109.0
C7—N1—C11	118.8 (2)	N1—C11—C12	112.2 (2)
C8—N1—C11	111.4 (2)	N1—C11—C10	103.5 (2)
O2—N2—C12	125.7 (2)	C12—C11—C10	113.3 (2)
O2—N2—C13	122.0 (2)	N1—C11—H11	109.2
C12—N2—C13	111.9 (2)	C12—C11—H11	109.2
O3—N3—C12	125.4 (2)	C10—C11—H11	109.2
O3—N3—C14	122.2 (2)	N3—C12—N2	109.4 (2)
C12—N3—C14	112.1 (2)	N3—C12—C11	126.1 (3)
C2—C1—C6	121.2 (3)	N2—C12—C11	124.5 (3)
C2—C1—Cl1	119.8 (3)	N2—C13—C15	110.0 (2)
C6—C1—Cl1	119.0 (3)	N2—C13—C16	105.2 (2)
C1—C2—C3	119.4 (3)	C15—C13—C16	111.2 (3)
C1—C2—H2A	120.3	N2—C13—C14	100.4 (2)
C3—C2—H2A	120.3	C15—C13—C14	114.7 (3)
C2—C3—C4	120.8 (3)	C16—C13—C14	114.3 (3)
C2—C3—H3A	119.6	N3—C14—C18	108.5 (3)
C4—C3—H3A	119.6	N3—C14—C17	105.7 (2)
C5—C4—C3	118.2 (3)	C18—C14—C17	110.0 (3)
C5—C4—C7	123.9 (3)	N3—C14—C13	100.9 (2)
C3—C4—C7	117.7 (3)	C18—C14—C13	116.2 (3)
C6—C5—C4	121.1 (3)	C17—C14—C13	114.3 (3)
C6—C5—H5	119.5	C13—C15—H15A	109.5
C4—C5—H5	119.5	C13—C15—H15B	109.5
C1—C6—C5	119.3 (3)	H15A—C15—H15B	109.5
C1—C6—H6	120.4	C13—C15—H15C	109.5
C5—C6—H6	120.4	H15A—C15—H15C	109.5
O1—C7—N1	120.3 (3)	H15B—C15—H15C	109.5
O1—C7—C4	119.8 (3)	C13—C16—H16A	109.5
N1—C7—C4	120.0 (2)	C13—C16—H16B	109.5
N1—C8—C9	103.3 (2)	H16A—C16—H16B	109.5

supplementary materials

N1—C8—H8A	111.1	C13—C16—H16C	109.5
C9—C8—H8A	111.1	H16A—C16—H16C	109.5
N1—C8—H8B	111.1	H16B—C16—H16C	109.5
C9—C8—H8B	111.1	C14—C17—H17A	109.5
H8A—C8—H8B	109.1	C14—C17—H17B	109.5
C8—C9—C10	103.4 (2)	H17A—C17—H17B	109.5
C8—C9—H9A	111.1	C14—C17—H17C	109.5
C10—C9—H9A	111.1	H17A—C17—H17C	109.5
C8—C9—H9B	111.1	H17B—C17—H17C	109.5
C10—C9—H9B	111.1	C14—C18—H18A	109.5
H9A—C9—H9B	109.1	C14—C18—H18B	109.5
C9—C10—C11	103.9 (2)	H18A—C18—H18B	109.5
C9—C10—H10A	111.0	C14—C18—H18C	109.5
C11—C10—H10A	111.0	H18A—C18—H18C	109.5
C9—C10—H10B	111.0	H18B—C18—H18C	109.5
C11—C10—H10B	111.0		
C6—C1—C2—C3	0.4 (5)	C14—N3—C12—C11	173.7 (2)
C11—C1—C2—C3	−179.5 (2)	O2—N2—C12—N3	178.9 (2)
C1—C2—C3—C4	0.9 (4)	C13—N2—C12—N3	−9.2 (3)
C2—C3—C4—C5	−1.3 (4)	O2—N2—C12—C11	−1.8 (4)
C2—C3—C4—C7	−176.6 (3)	C13—N2—C12—C11	170.2 (2)
C3—C4—C5—C6	0.4 (4)	N1—C11—C12—N3	49.2 (4)
C7—C4—C5—C6	175.4 (3)	C10—C11—C12—N3	−67.6 (4)
C2—C1—C6—C5	−1.3 (5)	N1—C11—C12—N2	−130.0 (3)
C11—C1—C6—C5	178.6 (2)	C10—C11—C12—N2	113.2 (3)
C4—C5—C6—C1	0.9 (5)	O2—N2—C13—C15	−46.3 (3)
C8—N1—C7—O1	−175.4 (3)	C12—N2—C13—C15	141.3 (2)
C11—N1—C7—O1	0.4 (4)	O2—N2—C13—C16	73.5 (3)
C8—N1—C7—C4	4.3 (4)	C12—N2—C13—C16	−98.8 (3)
C11—N1—C7—C4	−179.9 (2)	O2—N2—C13—C14	−167.6 (2)
C5—C4—C7—O1	−135.9 (3)	C12—N2—C13—C14	20.1 (3)
C3—C4—C7—O1	39.1 (4)	O3—N3—C14—C18	−45.1 (4)
C5—C4—C7—N1	44.4 (4)	C12—N3—C14—C18	141.5 (3)
C3—C4—C7—N1	−140.7 (3)	O3—N3—C14—C17	72.9 (3)
C7—N1—C8—C9	156.1 (3)	C12—N3—C14—C17	−100.5 (3)
C11—N1—C8—C9	−20.0 (3)	O3—N3—C14—C13	−167.8 (3)
N1—C8—C9—C10	35.2 (3)	C12—N3—C14—C13	18.9 (3)
C8—C9—C10—C11	−37.6 (3)	N2—C13—C14—N3	−21.4 (3)
C7—N1—C11—C12	57.7 (3)	C15—C13—C14—N3	−139.2 (3)
C8—N1—C11—C12	−125.8 (3)	C16—C13—C14—N3	90.6 (3)
C7—N1—C11—C10	−179.8 (2)	N2—C13—C14—C18	−138.5 (3)
C8—N1—C11—C10	−3.3 (3)	C15—C13—C14—C18	103.7 (3)
C9—C10—C11—N1	25.2 (3)	C16—C13—C14—C18	−26.5 (4)
C9—C10—C11—C12	147.0 (3)	N2—C13—C14—C17	91.5 (3)
O3—N3—C12—N2	179.9 (3)	C15—C13—C14—C17	−26.3 (4)
C14—N3—C12—N2	−7.0 (3)	C16—C13—C14—C17	−156.4 (3)
O3—N3—C12—C11	0.6 (5)		

Hydrogen-bond geometry (Å, °)

	D—H	H···A	D···A	D—H···A
D—H···A				
C2—H2A···O2 ⁱ	0.93	2.54	3.458 (4)	170
C3—H3A···O1 ⁱ	0.93	2.45	3.314 (4)	154
C8—H8A···O3 ⁱⁱ	0.97	2.44	3.101 (4)	125

Symmetry codes: (i) $-x+1, -y, -z$; (ii) $-x+3/2, y-1/2, -z+1/2$.

supplementary materials

Fig. 1

