

Acta Crystallographica Section E

Structure Reports

Online

ISSN 1600-5368

(E)-N'-(2,4-Dihydroxybenzylidene)-4-nitrobenzohydrazide

Nooraziah Mohd Lair, Hamid Khaleedi* and Hapipah Mohd Ali

Department of Chemistry, University of Malaya, 50603 Kuala Lumpur, Malaysia
Correspondence e-mail: khaleedi@siswa.um.edu.my


Received 10 January 2011; accepted 14 January 2011

Key indicators: single-crystal X-ray study; $T = 100$ K; mean $\sigma(\text{C}-\text{C}) = 0.002$ Å; R factor = 0.034; wR factor = 0.099; data-to-parameter ratio = 11.5.

The title compound, $\text{C}_{14}\text{H}_{11}\text{N}_3\text{O}_5$, is essentially planar, with an r.m.s. deviation for the non-H atoms of 0.0832 (3) Å. In the crystal, $\text{O}-\text{H}\cdots\text{O}$ and $\text{N}-\text{H}\cdots\text{O}$ hydrogen bonds link adjacent molecules into layers parallel to (101). These layers are further connected into a three-dimensional network *via* $\text{C}-\text{H}\cdots\text{O}$ interactions. In addition, a $\pi-\pi$ interaction occurs between the aromatic rings [centroid-centroid distance = 3.5425 (8) Å]. An intramolecular $\text{O}-\text{H}\cdots\text{N}$ hydrogen bond is also observed.

Related literature

For related structures, see: Han & Zhao (2010); Mohd Lair *et al.* (2009); Raj *et al.* (2008).



Experimental

Crystal data

$\text{C}_{14}\text{H}_{11}\text{N}_3\text{O}_5$
 $M_r = 301.26$
 Monoclinic, $P2_1/c$
 $a = 8.0248$ (1) Å
 $b = 12.5674$ (2) Å

$c = 12.8770$ (2) Å
 $\beta = 96.732$ (1)°
 $V = 1289.70$ (3) Å³
 $Z = 4$
 Mo $K\alpha$ radiation

$\mu = 0.12$ mm⁻¹
 $T = 100$ K

0.21 × 0.15 × 0.08 mm

Data collection

Bruker APEXII CCD diffractometer
 Absorption correction: multi-scan (SADABS; Sheldrick, 1996)
 $T_{\min} = 0.975$, $T_{\max} = 0.990$

10373 measured reflections
 2398 independent reflections
 2004 reflections with $I > 2\sigma(I)$
 $R_{\text{int}} = 0.027$

Refinement

$R[F^2 > 2\sigma(F^2)] = 0.034$
 $wR(F^2) = 0.099$
 $S = 1.05$
 2398 reflections
 208 parameters
 3 restraints

H atoms treated by a mixture of independent and constrained refinement
 $\Delta\rho_{\text{max}} = 0.22$ e Å⁻³
 $\Delta\rho_{\text{min}} = -0.27$ e Å⁻³

Table 1
 Hydrogen-bond geometry (Å, °).

$D-H\cdots A$	$D-H$	$H\cdots A$	$D\cdots A$	$D-H\cdots A$
$\text{O1}-\text{H1}\cdots\text{N1}$	0.86 (2)	1.93 (2)	2.6818 (15)	146 (2)
$\text{O2}-\text{H2A}\cdots\text{O3}^i$	0.84 (1)	1.84 (2)	2.6759 (14)	173 (2)
$\text{N2}-\text{H2B}\cdots\text{O5}^{ii}$	0.87 (1)	2.28 (1)	3.0606 (16)	150 (1)
$\text{C2}-\text{H2}\cdots\text{O3}^i$	0.95	2.47	3.1730 (17)	131
$\text{C4}-\text{H4}\cdots\text{O4}^{iii}$	0.95	2.54	3.3428 (17)	143
$\text{C7}-\text{H7}\cdots\text{O5}^{ii}$	0.95	2.40	3.2082 (17)	143
$\text{C10}-\text{H10}\cdots\text{O4}^{ii}$	0.95	2.52	3.3627 (18)	147

Symmetry codes: (i) $-x, y + \frac{1}{2}, -z + \frac{5}{2}$; (ii) $-x + 1, y + \frac{1}{2}, -z + \frac{3}{2}$; (iii) $x - 1, y + 1, z$.

Data collection: APEX2 (Bruker, 2007); cell refinement: SAINT (Bruker, 2007); data reduction: SAINT; program(s) used to solve structure: SHELXS97 (Sheldrick, 2008); program(s) used to refine structure: SHELXL97 (Sheldrick, 2008); molecular graphics: X-SEED (Barbour, 2001); software used to prepare material for publication: SHELXL97 and publCIF (Westrip, 2010).

The authors thank the University of Malaya for funding this study (FRGS grant No. FP004/2010B).

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: IS2667).

References

- Barbour, L. J. (2001). *J. Supramol. Chem.* **1**, 189–191.
 Bruker (2007). APEX2 and SAINT. Bruker AXS Inc., Madison, Wisconsin, USA.
 Han, Y.-Y. & Zhao, Q.-R. (2010). *Acta Cryst.* **E66**, o1041.
 Mohd Lair, N., Mohd Ali, H. & Ng, S. W. (2009). *Acta Cryst.* **E65**, o189.
 Raj, B. N. B., Kurup, M. R. P. & Suresh, E. (2008). *Spectrochim. Acta Part A*, **71**, 1253–1260.
 Sheldrick, G. M. (1996). SADABS. University of Göttingen, Germany.
 Sheldrick, G. M. (2008). *Acta Cryst.* **A64**, 112–122.
 Westrip, S. P. (2010). *J. Appl. Cryst.* **43**, 920–925.

supplementary materials

Acta Cryst. (2011). E67, o459 [doi:10.1107/S1600536811002224]

(*E*)-*N'*-(2,4-Dihydroxybenzylidene)-4-nitrobenzohydrazide

N. Mohd Lair, H. Khaledi and H. Mohd Ali

Comment

The title compound is the product of the condensation reaction of 4-nitrobenzohydrazide and 4-hydroxysalicylaldehyde. In agreement with the structures of similar benzoylhydrazones (Han & Zhao, 2010; Mohd Lair *et al.*, 2009; Raj *et al.*, 2008), the molecular structure of the present molecule is almost planar, the r.m.s. deviation for the non-H atoms being 0.0832 Å. The crystal structure is stabilized by O—H···O, N—H···O and C—H···O intermolecular and also O—H···N intramolecular hydrogen bonding. Moreover, a π – π interaction occurs between the aromatic rings of pairs of molecules related by symmetry $-x, -y + 1, -z + 2$ with centroid-centroid separation of 3.5425 (8) Å.

Experimental

A mixture of 4-nitrobenzohydrazide (0.54 g, 3 mmol) and 4-hydroxysalicylaldehyde (0.39 g, 3 mmol) in ethanol (50 ml) and in the presence of a few drops of acetic acid was refluxed for 5 hr. The solution was then left at room temperature. The crystals of the title compound were obtained in a few days.

Refinement

The carbon-bound H atoms were placed at calculated positions (C—H = 0.95 Å) and treated as riding on their parent carbon atoms. The nitrogen- and oxygen-bound H atoms were located in a difference map and refined as free atoms, with N—H and O—H distances restrained to 0.88 (2) and 0.84 (2) Å, respectively. $U_{\text{iso}}(\text{H})$ values were set to 1.2–1.5 $U_{\text{eq}}(\text{carrier atom})$.

Figures




Fig. 1. Displacement ellipsoid plot of the title compound at the 50% probability level. Hydrogen atoms are drawn as spheres of arbitrary radius

(*E*)-*N'*-(2,4-Dihydroxybenzylidene)-4-nitrobenzohydrazide

Crystal data

C₁₄H₁₁N₃O₅

$M_r = 301.26$

Monoclinic, $P2_1/c$

Hall symbol: $-P\ 2ybc$

$a = 8.0248$ (1) Å

$b = 12.5674$ (2) Å

$c = 12.8770$ (2) Å

$F(000) = 624$

$D_x = 1.552$ Mg m⁻³

Mo $K\alpha$ radiation, $\lambda = 0.71073$ Å

Cell parameters from 3471 reflections

$\theta = 3.2$ – 30.4°

$\mu = 0.12$ mm⁻¹

$T = 100$ K

supplementary materials

$\beta = 96.732 (1)^\circ$ Block, orange
 $V = 1289.70 (3) \text{ \AA}^3$ $0.21 \times 0.15 \times 0.08 \text{ mm}$
 $Z = 4$

Data collection

Bruker APEXII CCD diffractometer	2398 independent reflections
Radiation source: fine-focus sealed tube graphite	2004 reflections with $I > 2\sigma(I)$
φ and ω scans	$R_{\text{int}} = 0.027$
Absorption correction: multi-scan (SADABS; Sheldrick, 1996)	$\theta_{\text{max}} = 25.5^\circ$, $\theta_{\text{min}} = 2.3^\circ$
$T_{\text{min}} = 0.975$, $T_{\text{max}} = 0.990$	$h = -9 \rightarrow 9$
10373 measured reflections	$k = -15 \rightarrow 12$
	$l = -15 \rightarrow 15$

Refinement

Refinement on F^2	Primary atom site location: structure-invariant direct methods
Least-squares matrix: full	Secondary atom site location: difference Fourier map
$R[F^2 > 2\sigma(F^2)] = 0.034$	Hydrogen site location: inferred from neighbouring sites
$wR(F^2) = 0.099$	H atoms treated by a mixture of independent and constrained refinement
$S = 1.05$	$w = 1/[\sigma^2(F_o^2) + (0.0584P)^2 + 0.2741P]$
2398 reflections	where $P = (F_o^2 + 2F_c^2)/3$
208 parameters	$(\Delta/\sigma)_{\text{max}} = 0.001$
3 restraints	$\Delta\rho_{\text{max}} = 0.22 \text{ e \AA}^{-3}$
	$\Delta\rho_{\text{min}} = -0.27 \text{ e \AA}^{-3}$

Special details

Geometry. All e.s.d.'s (except the e.s.d. in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell e.s.d.'s are taken into account individually in the estimation of e.s.d.'s in distances, angles and torsion angles; correlations between e.s.d.'s in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell e.s.d.'s is used for estimating e.s.d.'s involving l.s. planes.

Refinement. Refinement of F^2 against ALL reflections. The weighted R -factor wR and goodness of fit S are based on F^2 , conventional R -factors R are based on F , with F set to zero for negative F^2 . The threshold expression of $F^2 > \sigma(F^2)$ is used only for calculating R -factors(gt) etc. and is not relevant to the choice of reflections for refinement. R -factors based on F^2 are statistically about twice as large as those based on F , and R -factors based on ALL data will be even larger.

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (\AA^2)

	x	y	z	$U_{\text{iso}}^*/U_{\text{eq}}$
O1	0.01817 (13)	0.64009 (9)	1.21087 (8)	0.0250 (3)
H1	0.073 (2)	0.5985 (14)	1.1740 (14)	0.038*

O2	-0.25926 (12)	0.97321 (8)	1.19160 (8)	0.0200 (2)
H2A	-0.271 (2)	0.9547 (14)	1.2529 (12)	0.030*
O3	0.31162 (13)	0.40161 (8)	1.11922 (8)	0.0232 (3)
O4	0.70514 (13)	0.04460 (8)	0.83143 (8)	0.0287 (3)
O5	0.69197 (12)	0.15345 (9)	0.70111 (8)	0.0248 (3)
N1	0.15626 (13)	0.57805 (9)	1.04095 (9)	0.0180 (3)
N2	0.23932 (14)	0.50897 (9)	0.98041 (9)	0.0173 (3)
H2B	0.2412 (19)	0.5289 (12)	0.9163 (11)	0.021*
N3	0.66607 (14)	0.13011 (10)	0.79089 (9)	0.0193 (3)
C1	-0.02829 (16)	0.72791 (11)	1.15383 (11)	0.0170 (3)
C2	-0.11597 (16)	0.80504 (11)	1.20227 (11)	0.0171 (3)
H2	-0.1393	0.7950	1.2722	0.021*
C3	-0.16953 (16)	0.89687 (11)	1.14835 (11)	0.0168 (3)
C4	-0.13465 (16)	0.91253 (11)	1.04548 (11)	0.0182 (3)
H4	-0.1708	0.9756	1.0088	0.022*
C5	-0.04773 (16)	0.83605 (11)	0.99795 (11)	0.0178 (3)
H5	-0.0243	0.8472	0.9282	0.021*
C6	0.00759 (16)	0.74153 (11)	1.04988 (11)	0.0165 (3)
C7	0.09681 (16)	0.66335 (11)	0.99533 (11)	0.0178 (3)
H7	0.1119	0.6752	0.9242	0.021*
C8	0.31525 (16)	0.42271 (11)	1.02595 (11)	0.0162 (3)
C9	0.40734 (15)	0.35067 (11)	0.95936 (10)	0.0156 (3)
C10	0.44714 (16)	0.37683 (11)	0.85999 (11)	0.0171 (3)
H10	0.4134	0.4436	0.8299	0.020*
C11	0.53565 (16)	0.30596 (11)	0.80500 (11)	0.0178 (3)
H11	0.5651	0.3237	0.7378	0.021*
C12	0.58004 (16)	0.20868 (11)	0.85038 (11)	0.0171 (3)
C13	0.54399 (16)	0.18076 (11)	0.94917 (11)	0.0183 (3)
H13	0.5776	0.1138	0.9788	0.022*
C14	0.45772 (16)	0.25303 (11)	1.00359 (11)	0.0174 (3)
H14	0.4325	0.2359	1.0719	0.021*

Atomic displacement parameters (\AA^2)

	U^{11}	U^{22}	U^{33}	U^{12}	U^{13}	U^{23}
O1	0.0306 (6)	0.0227 (6)	0.0229 (6)	0.0100 (5)	0.0082 (4)	0.0054 (4)
O2	0.0261 (5)	0.0173 (5)	0.0178 (5)	0.0037 (4)	0.0071 (4)	-0.0005 (4)
O3	0.0320 (6)	0.0225 (6)	0.0170 (5)	0.0004 (4)	0.0110 (4)	0.0007 (4)
O4	0.0369 (6)	0.0231 (6)	0.0256 (6)	0.0117 (5)	0.0013 (5)	-0.0019 (5)
O5	0.0252 (5)	0.0326 (6)	0.0179 (5)	0.0014 (5)	0.0076 (4)	-0.0044 (4)
N1	0.0152 (5)	0.0186 (6)	0.0209 (6)	-0.0013 (5)	0.0057 (5)	-0.0055 (5)
N2	0.0182 (6)	0.0190 (6)	0.0156 (6)	0.0005 (5)	0.0063 (5)	-0.0032 (5)
N3	0.0165 (6)	0.0231 (7)	0.0180 (6)	0.0001 (5)	0.0003 (5)	-0.0053 (5)
C1	0.0145 (6)	0.0173 (7)	0.0190 (7)	-0.0017 (5)	0.0007 (5)	0.0010 (6)
C2	0.0169 (6)	0.0207 (7)	0.0142 (7)	-0.0020 (5)	0.0033 (5)	-0.0001 (6)
C3	0.0147 (6)	0.0165 (7)	0.0194 (7)	-0.0025 (5)	0.0027 (5)	-0.0030 (6)
C4	0.0184 (6)	0.0165 (7)	0.0196 (7)	-0.0003 (5)	0.0021 (5)	0.0027 (6)
C5	0.0179 (7)	0.0221 (8)	0.0139 (7)	-0.0035 (6)	0.0039 (5)	0.0001 (5)

supplementary materials

C6	0.0135 (6)	0.0182 (7)	0.0182 (7)	-0.0028 (5)	0.0036 (5)	-0.0020 (6)
C7	0.0141 (6)	0.0212 (7)	0.0183 (7)	-0.0040 (5)	0.0029 (5)	-0.0021 (6)
C8	0.0158 (6)	0.0166 (7)	0.0167 (7)	-0.0056 (5)	0.0039 (5)	-0.0020 (6)
C9	0.0135 (6)	0.0163 (7)	0.0169 (7)	-0.0038 (5)	0.0016 (5)	-0.0029 (5)
C10	0.0178 (6)	0.0162 (7)	0.0172 (7)	-0.0016 (5)	0.0023 (5)	0.0004 (5)
C11	0.0171 (6)	0.0220 (7)	0.0147 (7)	-0.0029 (6)	0.0033 (5)	-0.0014 (6)
C12	0.0134 (6)	0.0193 (7)	0.0185 (7)	-0.0008 (5)	0.0023 (5)	-0.0062 (6)
C13	0.0180 (7)	0.0180 (7)	0.0183 (7)	-0.0006 (6)	-0.0005 (5)	-0.0009 (6)
C14	0.0171 (7)	0.0208 (7)	0.0146 (7)	-0.0030 (5)	0.0025 (5)	-0.0008 (6)

Geometric parameters (Å, °)

O1—C1	1.3537 (17)	C4—C5	1.3729 (19)
O1—H1	0.860 (15)	C4—H4	0.9500
O2—C3	1.3576 (16)	C5—C6	1.409 (2)
O2—H2A	0.839 (14)	C5—H5	0.9500
O3—C8	1.2336 (16)	C6—C7	1.4454 (19)
O4—N3	1.2196 (16)	C7—H7	0.9500
O5—N3	1.2337 (15)	C8—C9	1.4999 (18)
N1—C7	1.2868 (19)	C9—C14	1.3927 (19)
N1—N2	1.3884 (16)	C9—C10	1.3939 (19)
N2—C8	1.3441 (19)	C10—C11	1.3847 (19)
N2—H2B	0.865 (13)	C10—H10	0.9500
N3—C12	1.4705 (17)	C11—C12	1.384 (2)
C1—C2	1.3880 (19)	C11—H11	0.9500
C1—C6	1.4121 (19)	C12—C13	1.3825 (19)
C2—C3	1.389 (2)	C13—C14	1.3817 (19)
C2—H2	0.9500	C13—H13	0.9500
C3—C4	1.3996 (19)	C14—H14	0.9500
C1—O1—H1	108.8 (13)	C1—C6—C7	123.09 (13)
C3—O2—H2A	108.2 (12)	N1—C7—C6	121.51 (13)
C7—N1—N2	116.19 (12)	N1—C7—H7	119.2
C8—N2—N1	118.83 (12)	C6—C7—H7	119.2
C8—N2—H2B	126.4 (11)	O3—C8—N2	122.48 (12)
N1—N2—H2B	114.7 (11)	O3—C8—C9	119.76 (13)
O4—N3—O5	123.19 (12)	N2—C8—C9	117.77 (12)
O4—N3—C12	118.79 (12)	C14—C9—C10	119.77 (12)
O5—N3—C12	118.00 (12)	C14—C9—C8	115.90 (12)
O1—C1—C2	116.57 (12)	C10—C9—C8	124.31 (13)
O1—C1—C6	122.64 (12)	C11—C10—C9	120.27 (13)
C2—C1—C6	120.78 (13)	C11—C10—H10	119.9
C1—C2—C3	119.94 (13)	C9—C10—H10	119.9
C1—C2—H2	120.0	C12—C11—C10	118.33 (13)
C3—C2—H2	120.0	C12—C11—H11	120.8
O2—C3—C2	122.02 (12)	C10—C11—H11	120.8
O2—C3—C4	117.61 (12)	C13—C12—C11	122.80 (13)
C2—C3—C4	120.36 (12)	C13—C12—N3	118.07 (13)
C5—C4—C3	119.47 (13)	C11—C12—N3	119.12 (12)
C5—C4—H4	120.3	C14—C13—C12	118.11 (13)

C3—C4—H4	120.3	C14—C13—H13	120.9
C4—C5—C6	121.79 (13)	C12—C13—H13	120.9
C4—C5—H5	119.1	C13—C14—C9	120.69 (13)
C6—C5—H5	119.1	C13—C14—H14	119.7
C5—C6—C1	117.65 (12)	C9—C14—H14	119.7
C5—C6—C7	119.27 (13)		

Hydrogen-bond geometry (\AA , $^\circ$)

<i>D</i> —H \cdots <i>A</i>	<i>D</i> —H	H \cdots <i>A</i>	<i>D</i> \cdots <i>A</i>	<i>D</i> —H \cdots <i>A</i>
O1—H1 \cdots N1	0.86 (2)	1.93 (2)	2.6818 (15)	146.(2)
O2—H2A \cdots O3 ⁱ	0.84 (1)	1.84 (2)	2.6759 (14)	173.(2)
N2—H2B \cdots O5 ⁱⁱ	0.87 (1)	2.28 (1)	3.0606 (16)	150.(1)
C2—H2 \cdots O3 ⁱ	0.95	2.47	3.1730 (17)	131
C4—H4 \cdots O4 ⁱⁱⁱ	0.95	2.54	3.3428 (17)	143
C7—H7 \cdots O5 ⁱⁱ	0.95	2.40	3.2082 (17)	143
C10—H10 \cdots O4 ⁱⁱ	0.95	2.52	3.3627 (18)	147

Symmetry codes: (i) $-x, y+1/2, -z+5/2$; (ii) $-x+1, y+1/2, -z+3/2$; (iii) $x-1, y+1, z$.

Fig. 1

