

Pyridinium tetrakis(1,1,1-trifluoropentane-2,4-dionate) dysprosite

Yan Wang,^{a,b*} Yuekui Wang,^b Jie Jia,^b Xiaoli Gao^b and Xiaoling Su^b

^aDepartment of Chemistry & Chemical Engineering, Luliang University, Luliang 033000, People's Republic of China, and ^bInstitute of Molecular Science, Key Laboratory of Chemical Biology and Molecular Engineering of the Education Ministry, Shanxi University, Taiyuan, Shanxi 030006, People's Republic of China
Correspondence e-mail: llswy74@yahoo.com.cn


Received 21 January 2011; accepted 10 February 2011

Key indicators: single-crystal X-ray study; $T = 298\text{ K}$; mean $\sigma(\text{C}-\text{C}) = 0.013\text{ \AA}$; disorder in main residue; R factor = 0.050; wR factor = 0.121; data-to-parameter ratio = 11.7.

In the anion of the title compound, $(\text{C}_5\text{H}_6\text{N})[\text{Dy}(\text{C}_5\text{H}_4\text{F}_3\text{O}_2)_4]$, the central metal ion, Dy^{3+} , is coordinated by four bidentate 1,1,1-trifluoropentane-2,4-dionate (TAA) ligands, forming an approximate square-antiprismatic configuration. The pyridinium cation is connected to the complex ion by an $\text{N}-\text{H}\cdots\text{O}$ hydrogen bond and electrostatic interactions in the crystal. There are two kinds of disorder in the structure, one involving rotational disorder of a CF_3 group [occupancy ratio 0.560 (15):0.440 (15)] and the other involving an exchange between a CF_3 group and CH_3 group within a given bidentate ligand (occupancy ratio 0.64:0.36).

Related literature

For applications of rare earth- β -diketone complexes, see: Chu & Elgavish (1995); Tsukube & Shinoda (2002); Iwamuro *et al.* (1997). For related structures, see: Ma *et al.* (2000); Tian *et al.* (2009).



Experimental

Crystal data

$(\text{C}_5\text{H}_6\text{N})[\text{Dy}(\text{C}_5\text{H}_4\text{F}_3\text{O}_2)_4]$
 $M_r = 854.94$
Monoclinic, $P2_1/c$
 $a = 10.619 (4)\text{ \AA}$
 $b = 19.799 (7)\text{ \AA}$

$c = 15.715 (6)\text{ \AA}$
 $\beta = 103.116 (6)^\circ$
 $V = 3217.8 (19)\text{ \AA}^3$
 $Z = 4$
Mo $K\alpha$ radiation

$\mu = 2.44\text{ mm}^{-1}$
 $T = 298\text{ K}$

$0.22 \times 0.22 \times 0.06\text{ mm}$

Data collection

Bruker SMART 1K CCD area detector diffractometer
Absorption correction: multi-scan (*SADABS*; Sheldrick, 2000)
 $T_{\min} = 0.616$, $T_{\max} = 0.868$

14449 measured reflections
5670 independent reflections
4126 reflections with $I > 2\sigma(I)$
 $R_{\text{int}} = 0.039$

Refinement

$R[F^2 > 2\sigma(F^2)] = 0.050$
 $wR(F^2) = 0.121$
 $S = 1.02$
5670 reflections
483 parameters

78 restraints
H-atom parameters constrained
 $\Delta\rho_{\max} = 0.95\text{ e \AA}^{-3}$
 $\Delta\rho_{\min} = -0.45\text{ e \AA}^{-3}$

Table 1
Selected bond lengths (\AA).

Dy1—O2	2.305 (5)	Dy1—O8	2.349 (4)
Dy1—O6	2.315 (5)	Dy1—O7	2.356 (5)
Dy1—O3	2.326 (5)	Dy1—O5	2.390 (5)
Dy1—O4	2.342 (5)	Dy1—O1	2.423 (5)

Table 2
Hydrogen-bond geometry (\AA , $^\circ$).

$D-\text{H}\cdots A$	$D-\text{H}$	$\text{H}\cdots A$	$D\cdots A$	$D-\text{H}\cdots A$
N1—H1 \cdots O5	0.86	2.10	2.947 (8)	167

Data collection: *SMART* (Bruker, 2000); cell refinement: *SAINT* (Bruker, 2000); data reduction: *SAINT*; program(s) used to solve structure: *SHELXS97* (Sheldrick, 2008); program(s) used to refine structure: *SHELXL97* (Sheldrick, 2008); molecular graphics: *ORTEP-3* (Farrugia, 1997); software used to prepare material for publication: *SHELXTL/PC* (Sheldrick, 2008).

This research was supported by the National Natural Science Foundation of China (grant No. 20673069) and the Natural Science Foundation of Shanxi province (grant No. 2007011021).

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: OM2404).

References

- Bruker (2000). *SMART* and *SAINT*. Bruker AXS Inc., Madison, Wisconsin, USA.
- Chu, W. J. & Elgavish, G. A. (1995). *NMR Biomed.* **8**, 159–163.
- Farrugia, L. J. (1997). *J. Appl. Cryst.* **30**, 565.
- Iwamuro, M., Hasegawa, Y., Wada, Y. & Murakoshi, K. (1997). *Chem. Lett.* **10**, 1067–1068.
- Ma, B. Q., Zhang, D. S., Gao, S., Jin, T. Z. & Yan, C. H. (2000). *Angew. Chem. Int. Ed.* **39**, 3644–3646.
- Sheldrick, G. M. (2000). *SADABS*. University of Göttingen, Germany.
- Sheldrick, G. M. (2008). *Acta Cryst. A* **64**, 112–122.
- Tian, L., Ren, N., Zhang, J.-J., Sun, S.-J., Ye, H.-M., Bai, J.-H. & Wang, R.-F. (2009). *J. Chem. Eng. Data* **54**, 69–74.
- Tsukube, H. & Shinoda, S. (2002). *Chem. Rev.* **102**, 2389–2404.

supplementary materials

Acta Cryst. (2011). E67, m341 [doi:10.1107/S1600536811005034]

Pyridinium tetrakis(1,1,1-trifluoropentane-2,4-dionato)dysprosate

Y. Wang, Y. Wang, J. Jia, X. Gao and X. Su

Comment

Rare earth- β -diketone complexes have attracted considerable attention in the past decades owing to their important applications as laser (Iwamuro *et al.*, 1997), fluorescent probe (Tsukube & Shinoda, 2002) and NMR reagents (Chu & Elgavish, 1995). As part of our interest in this field, we have been engaged in a major effort directed toward the development of syntheses of new lanthanide- β -diketon complexes.

The structure of the title Dy³⁺ complex is shown in Fig. 1. It contains an eight-coordinate dysprosium ion bonded to four TAA anions with bidentate chelation, forming the [Dy(TAA)₄]⁻ anions. These are connected to pyridinium cations by a N—H···O hydrogen bond. The co-ordination polyhedron may be described as an approximate square antiprism and the two sets of four O atoms (O2, O4, O5, O7) and (O1, O3, O6, O8) form the twisted upper and lower sides respectively. The Dy—O bond lengths are in the range of 2.305 (5)–2.423 (5) Å, average 2.351 (2) Å, which is consistent with other work in literature (Ma *et al.*, 2000; Tian *et al.*, 2009). The average angle of O—Dy—O is 100.25° and the average dihydral angle (C—O—Dy—O) is -10.49°.

Experimental

A mixture of Dy₂O₃ (0.186 g) and concentrated hydrochloric (5 mL) was heated and distilled to slight dryness, yielding a crystalline precipitate (DyCl₃). Then, the DyCl₃ solid was redissolved in 5 mL absolute ethanol, and heated with 10 mL of absolute ethanol solution containing HTAA (0.50 mL) and pyridine (0.32 mL) at about 363 K. The reaction mixture was maintained at ambient temperature for one month until yellow crystals formed.

Refinement

All F atoms were found to be disordered. There is disorder of the two different types: 1) disorder due to rotational disorder of the CF₃ group bonded to a single carbon. F4, F5, and F6 atoms was split into to two sets of positions using restraints on their anisotropic displacement parameters. The major and minor disorder components had refined occupancies of 0.56 (2) and 0.44 (2), respectively; 2) disorder due to exchange of CH₃ and CF₃ groups on the same ligand. Namely, F10, F11, and F12 as well as related H atoms were modelled over two sets of positions using restraint on their anisotropic displacement parameters. The major and minor disorder components had refined occupancies of 0.64 (1) and 0.36 (1), respectively. In the final refinement, the occupancies of these disordered atoms were fixed to aid convergence. Atoms F11B, F4, F5, F6, F4B, F5B, and F6B were refined anisotropically using 42 restraints (ISOR) and the geometrical parameters of CF₃ group were refined using 36 restraints (DFIX and DANG) because of the unacceptable parameters of their ellipsoids and distances between atoms. H atoms attached to C and N were placed in geometrically idealized positions with Csp²—H = 0.93, Csp³—H = 0.96, Nsp²—H = 0.86 Å, and constrained to ride on their carrier atoms, with Uiso~(H) = 1.2Ueq~(C & N) and Uiso~(H) = 1.5Ueq~(C).

supplementary materials

Figures




Fig. 1. A view of the structure of the complex with displacement ellipsoids drawn at the 30% probability level. H atoms without H-bond (dotting line) and minor disorder components were omitted for clarity.

Pyridinium tetrakis(1,1,1-trifluoropentane-2,4-dionato)dysprosate

Crystal data

(C ₅ H ₆ N)[Dy(C ₅ H ₄ F ₃ O ₂) ₄]	$F(000) = 1668$
$M_r = 854.94$	$D_x = 1.765 \text{ Mg m}^{-3}$
Monoclinic, $P2_1/c$	Mo $K\alpha$ radiation, $\lambda = 0.71073 \text{ \AA}$
Hall symbol: -P 2ybc	Cell parameters from 3207 reflections
$a = 10.619 (4) \text{ \AA}$	$\theta = 2.2\text{--}21.0^\circ$
$b = 19.799 (7) \text{ \AA}$	$\mu = 2.44 \text{ mm}^{-1}$
$c = 15.715 (6) \text{ \AA}$	$T = 298 \text{ K}$
$\beta = 103.116 (6)^\circ$	Plate, colorless
$V = 3217.8 (19) \text{ \AA}^3$	$0.22 \times 0.22 \times 0.06 \text{ mm}$
$Z = 4$	

Data collection

Bruker SMART 1K CCD area detector diffractometer	5670 independent reflections
Radiation source: fine-focus sealed tube graphite	4126 reflections with $I > 2\sigma(I)$
ω scans	$R_{\text{int}} = 0.039$
Absorption correction: multi-scan (<i>SADABS</i> ; Sheldrick, 2000)	$\theta_{\text{max}} = 25.0^\circ, \theta_{\text{min}} = 1.7^\circ$
$T_{\text{min}} = 0.616, T_{\text{max}} = 0.868$	$h = -12 \rightarrow 12$
14449 measured reflections	$k = -19 \rightarrow 23$
	$l = -18 \rightarrow 15$

Refinement

Refinement on F^2	Primary atom site location: structure-invariant direct methods
Least-squares matrix: full	Secondary atom site location: difference Fourier map
$R[F^2 > 2\sigma(F^2)] = 0.050$	Hydrogen site location: inferred from neighbouring sites
$wR(F^2) = 0.121$	H-atom parameters constrained
$S = 1.02$	$w = 1/[\sigma^2(F_o^2) + (0.0621P)^2]$ where $P = (F_o^2 + 2F_c^2)/3$

5670 reflections	$(\Delta/\sigma)_{\max} = 0.001$
483 parameters	$\Delta\rho_{\max} = 0.95 \text{ e \AA}^{-3}$
78 restraints	$\Delta\rho_{\min} = -0.45 \text{ e \AA}^{-3}$

Special details

Geometry. All e.s.d.'s (except the e.s.d. in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell e.s.d.'s are taken into account individually in the estimation of e.s.d.'s in distances, angles and torsion angles; correlations between e.s.d.'s in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell e.s.d.'s is used for estimating e.s.d.'s involving l.s. planes.

Refinement. Refinement of F^2 against ALL reflections. The weighted R -factor wR and goodness of fit S are based on F^2 , conventional R -factors R are based on F , with F set to zero for negative F^2 . The threshold expression of $F^2 > \sigma(F^2)$ is used only for calculating R -factors(gt) etc. and is not relevant to the choice of reflections for refinement. R -factors based on F^2 are statistically about twice as large as those based on F , and R -factors based on ALL data will be even larger.

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (\AA^2)

	<i>x</i>	<i>y</i>	<i>z</i>	$U_{\text{iso}}^*/U_{\text{eq}}$	Occ. (<1)
Dy1	0.54349 (3)	0.155070 (15)	0.274306 (19)	0.05328 (14)	
O1	0.4836 (5)	0.2430 (2)	0.3649 (3)	0.0698 (13)	
O2	0.3848 (5)	0.2201 (2)	0.1874 (3)	0.0664 (13)	
C1	0.3730 (10)	0.3186 (5)	0.4363 (6)	0.105 (3)	
H1A	0.4141	0.2953	0.4889	0.157*	
H1B	0.2835	0.3259	0.4360	0.157*	
H1C	0.4147	0.3614	0.4339	0.157*	
C2	0.3834 (8)	0.2772 (4)	0.3590 (5)	0.0656 (19)	
C3	0.2847 (8)	0.2808 (4)	0.2828 (5)	0.069 (2)	
H3	0.2086	0.3029	0.2860	0.083*	
C4	0.2930 (7)	0.2543 (4)	0.2051 (5)	0.0633 (19)	
C5	0.1851 (9)	0.2668 (5)	0.1261 (6)	0.088 (3)	
F1	0.2181 (6)	0.3041 (4)	0.0687 (4)	0.170 (3)	
F2	0.1395 (7)	0.2116 (4)	0.0866 (5)	0.165 (3)	
F3	0.0817 (7)	0.2960 (4)	0.1425 (4)	0.170 (3)	
O3	0.3793 (5)	0.1120 (2)	0.3346 (3)	0.0691 (13)	
O4	0.4291 (6)	0.0745 (3)	0.1777 (3)	0.0813 (15)	
C6	0.1664 (9)	0.1026 (6)	0.3557 (6)	0.116 (3)	
H6A	0.1652	0.0631	0.3908	0.173*	
H6B	0.0825	0.1092	0.3180	0.173*	
H6C	0.1886	0.1413	0.3930	0.173*	
C7	0.2647 (9)	0.0938 (4)	0.3014 (6)	0.076 (2)	
C8	0.2294 (10)	0.0660 (5)	0.2176 (7)	0.097 (3)	
H8	0.1438	0.0530	0.1968	0.116*	
C9	0.3145 (12)	0.0569 (4)	0.1642 (5)	0.090 (3)	
C10	0.2692 (12)	0.0137 (6)	0.0779 (8)	0.129 (4)	
F4	0.2652 (17)	0.0495 (7)	0.0147 (8)	0.139 (5)	0.560 (15)
F5	0.3418 (14)	-0.0434 (6)	0.0845 (8)	0.138 (5)	0.560 (15)
F6	0.1486 (13)	-0.0078 (8)	0.0698 (9)	0.152 (6)	0.560 (15)

supplementary materials

F4B	0.237 (2)	-0.0437 (7)	0.0915 (10)	0.136 (6)	0.440 (15)
F5B	0.158 (2)	0.0463 (12)	0.0364 (15)	0.201 (10)	0.440 (15)
F6B	0.3444 (16)	0.0219 (8)	0.0223 (10)	0.123 (6)	0.440 (15)
O5	0.6258 (5)	0.1812 (2)	0.1486 (3)	0.0685 (13)	
O6	0.6880 (5)	0.0700 (2)	0.2623 (3)	0.0687 (13)	
C11	0.7528 (9)	0.1973 (5)	0.0427 (5)	0.103 (3)	
H11A	0.6789	0.2041	-0.0045	0.155*	
H11B	0.8178	0.1727	0.0219	0.155*	
H11C	0.7869	0.2402	0.0652	0.155*	
C12	0.7130 (8)	0.1571 (4)	0.1151 (5)	0.0664 (19)	
C13	0.7781 (8)	0.0965 (4)	0.1434 (5)	0.075 (2)	
H13	0.8392	0.0817	0.1135	0.089*	
C14	0.7591 (7)	0.0577 (3)	0.2112 (5)	0.0641 (19)	
C15	0.8293 (10)	-0.0080 (5)	0.2295 (7)	0.093 (3)	
F7	0.9130 (8)	-0.0209 (3)	0.1817 (6)	0.170 (3)	
F8	0.7516 (7)	-0.0582 (3)	0.2151 (6)	0.185 (4)	
F9	0.8939 (9)	-0.0126 (4)	0.3073 (5)	0.194 (4)	
O7	0.7171 (5)	0.2319 (2)	0.3091 (3)	0.0692 (13)	
O8	0.6620 (5)	0.1262 (2)	0.4147 (3)	0.0624 (12)	
C16	0.9245 (11)	0.2782 (6)	0.3435 (7)	0.104 (3)	
H16A	0.9699	0.2688	0.2962	0.104*	0.64
H16B	1.0012	0.2854	0.3963	0.104*	0.64
H16C	0.8937	0.3261	0.3317	0.104*	0.64
F10B	1.0364 (15)	0.2718 (10)	0.3858 (12)	0.157 (8)	0.36
F11B	0.924 (2)	0.2781 (10)	0.2596 (11)	0.160 (7)	0.36
F12B	0.879 (2)	0.3390 (8)	0.3519 (18)	0.193 (12)	0.36
C17	0.8278 (8)	0.2252 (4)	0.3569 (5)	0.068 (2)	
C18	0.8665 (8)	0.1759 (4)	0.4206 (5)	0.070 (2)	
H18	0.9536	0.1735	0.4483	0.084*	
C19	0.7838 (7)	0.1309 (3)	0.4447 (4)	0.0589 (18)	
C20	0.8358 (10)	0.0834 (5)	0.5191 (6)	0.099 (3)	
H20A	0.7625	0.0425	0.5206	0.099*	0.36
H20B	0.8509	0.0975	0.5761	0.099*	0.36
H20C	0.9073	0.0496	0.5146	0.099*	0.36
F10	0.9621 (9)	0.0751 (5)	0.5361 (6)	0.141 (3)	0.64
F11	0.7907 (13)	0.0241 (5)	0.5025 (9)	0.243 (9)	0.64
F12	0.8159 (13)	0.1056 (7)	0.5915 (6)	0.194 (6)	0.64
N1	0.5608 (8)	0.3261 (3)	0.1375 (6)	0.096 (2)	
H1	0.5924	0.2859	0.1424	0.115*	
C21	0.5610 (9)	0.3615 (5)	0.2096 (6)	0.089 (3)	
H21	0.5963	0.3435	0.2645	0.106*	
C22	0.5089 (10)	0.4241 (4)	0.2015 (6)	0.088 (3)	
H22	0.5072	0.4495	0.2510	0.105*	
C23	0.4599 (9)	0.4493 (4)	0.1219 (6)	0.089 (3)	
H23	0.4224	0.4920	0.1162	0.106*	
C24	0.4647 (9)	0.4130 (4)	0.0494 (6)	0.088 (3)	
H24	0.4341	0.4316	-0.0057	0.106*	
C25	0.5124 (10)	0.3516 (4)	0.0571 (7)	0.089 (3)	
H25	0.5129	0.3259	0.0076	0.107*	

Atomic displacement parameters (\AA^2)

	U^{11}	U^{22}	U^{33}	U^{12}	U^{13}	U^{23}
Dy1	0.0585 (2)	0.0521 (2)	0.0466 (2)	0.00072 (16)	0.00630 (14)	0.00471 (15)
O1	0.067 (3)	0.074 (3)	0.062 (3)	0.006 (3)	0.002 (2)	-0.007 (2)
O2	0.062 (3)	0.078 (3)	0.054 (3)	0.011 (3)	0.004 (2)	0.009 (2)
C1	0.123 (9)	0.100 (6)	0.092 (7)	0.029 (6)	0.027 (6)	-0.018 (5)
C2	0.068 (5)	0.054 (4)	0.075 (5)	0.003 (4)	0.017 (4)	0.007 (4)
C3	0.068 (5)	0.063 (5)	0.076 (5)	0.015 (4)	0.013 (4)	0.009 (4)
C4	0.058 (5)	0.062 (4)	0.063 (5)	0.004 (4)	-0.001 (4)	0.020 (4)
C5	0.072 (6)	0.108 (7)	0.077 (6)	0.010 (6)	0.000 (5)	0.014 (6)
F1	0.125 (5)	0.248 (8)	0.120 (5)	-0.011 (5)	-0.008 (4)	0.111 (6)
F2	0.137 (6)	0.161 (6)	0.148 (6)	-0.001 (5)	-0.067 (4)	-0.004 (5)
F3	0.106 (5)	0.275 (9)	0.111 (5)	0.088 (6)	-0.010 (4)	0.003 (5)
O3	0.071 (4)	0.071 (3)	0.063 (3)	-0.009 (3)	0.009 (3)	0.010 (3)
O4	0.084 (4)	0.085 (4)	0.071 (3)	-0.020 (3)	0.008 (3)	-0.026 (3)
C6	0.078 (7)	0.159 (10)	0.122 (8)	-0.001 (7)	0.049 (6)	0.015 (7)
C7	0.075 (6)	0.065 (5)	0.083 (6)	-0.013 (4)	0.007 (5)	0.014 (4)
C8	0.081 (7)	0.105 (7)	0.096 (7)	-0.021 (5)	0.002 (6)	0.004 (6)
C9	0.123 (9)	0.069 (5)	0.063 (5)	-0.020 (6)	-0.008 (5)	-0.006 (4)
C10	0.130 (11)	0.126 (11)	0.122 (10)	-0.037 (9)	0.008 (8)	0.026 (8)
F4	0.147 (7)	0.143 (7)	0.122 (6)	-0.008 (5)	0.021 (5)	-0.001 (4)
F5	0.145 (7)	0.131 (7)	0.130 (6)	0.002 (4)	0.015 (4)	-0.025 (4)
F6	0.148 (7)	0.151 (7)	0.151 (7)	-0.022 (5)	0.020 (5)	-0.006 (5)
F4B	0.143 (8)	0.128 (7)	0.137 (7)	-0.008 (5)	0.030 (5)	-0.011 (5)
F5B	0.201 (11)	0.203 (11)	0.194 (11)	-0.002 (5)	0.033 (5)	-0.008 (5)
F6B	0.127 (7)	0.125 (7)	0.115 (7)	-0.006 (5)	0.025 (5)	-0.012 (5)
O5	0.083 (4)	0.067 (3)	0.058 (3)	0.006 (3)	0.021 (3)	0.011 (2)
O6	0.085 (4)	0.061 (3)	0.064 (3)	0.015 (3)	0.024 (3)	0.005 (2)
C11	0.118 (8)	0.129 (8)	0.078 (6)	0.001 (7)	0.053 (6)	0.021 (6)
C12	0.064 (5)	0.077 (5)	0.057 (4)	-0.002 (4)	0.013 (4)	-0.010 (4)
C13	0.078 (6)	0.081 (6)	0.071 (5)	0.001 (5)	0.030 (4)	-0.014 (4)
C14	0.065 (5)	0.051 (4)	0.070 (5)	-0.005 (4)	0.002 (4)	-0.018 (4)
C15	0.099 (8)	0.077 (6)	0.108 (8)	0.014 (6)	0.034 (6)	0.001 (6)
F7	0.184 (7)	0.125 (5)	0.231 (8)	0.067 (5)	0.112 (7)	0.020 (5)
F8	0.154 (7)	0.062 (3)	0.348 (12)	-0.006 (4)	0.074 (7)	-0.012 (5)
F9	0.267 (10)	0.153 (6)	0.129 (6)	0.118 (7)	-0.024 (6)	0.014 (5)
O7	0.061 (3)	0.072 (3)	0.071 (3)	-0.005 (3)	0.006 (3)	0.007 (3)
O8	0.063 (3)	0.067 (3)	0.053 (3)	0.004 (3)	0.005 (2)	0.012 (2)
C16	0.091 (8)	0.123 (9)	0.095 (7)	-0.029 (7)	0.015 (6)	0.000 (6)
F10B	0.084 (13)	0.21 (2)	0.160 (16)	-0.055 (13)	-0.010 (11)	0.055 (14)
F11B	0.158 (8)	0.168 (8)	0.157 (8)	-0.016 (5)	0.043 (5)	0.003 (5)
F12B	0.15 (2)	0.110 (14)	0.35 (4)	-0.061 (13)	0.14 (2)	-0.024 (16)
C17	0.067 (5)	0.074 (5)	0.064 (5)	-0.008 (4)	0.019 (4)	-0.016 (4)
C18	0.056 (5)	0.076 (5)	0.069 (5)	0.008 (4)	-0.002 (4)	-0.005 (4)
C19	0.060 (5)	0.058 (4)	0.050 (4)	0.017 (4)	-0.004 (3)	-0.006 (3)
C20	0.086 (7)	0.105 (8)	0.085 (7)	0.005 (6)	-0.025 (5)	0.028 (6)

supplementary materials

F10	0.107 (8)	0.146 (8)	0.145 (8)	0.029 (6)	-0.020 (6)	0.041 (6)
F11	0.243 (14)	0.124 (8)	0.253 (15)	-0.077 (9)	-0.170 (12)	0.118 (9)
F12	0.239 (14)	0.273 (15)	0.079 (6)	0.122 (12)	0.052 (8)	0.063 (8)
N1	0.114 (7)	0.059 (4)	0.134 (7)	0.011 (4)	0.069 (6)	0.016 (5)
C21	0.098 (7)	0.094 (7)	0.073 (6)	-0.007 (5)	0.017 (5)	0.020 (5)
C22	0.126 (8)	0.065 (5)	0.074 (6)	-0.003 (5)	0.024 (5)	-0.011 (5)
C23	0.098 (7)	0.064 (5)	0.105 (7)	0.017 (5)	0.024 (6)	-0.002 (5)
C24	0.109 (7)	0.080 (6)	0.072 (5)	-0.008 (5)	0.010 (5)	0.017 (5)
C25	0.116 (8)	0.067 (6)	0.100 (7)	-0.023 (5)	0.058 (6)	-0.016 (5)

Geometric parameters (\AA , $^\circ$)

Dy1—O2	2.305 (5)	C13—H13	0.9300
Dy1—O6	2.315 (5)	C14—C15	1.494 (11)
Dy1—O3	2.326 (5)	C15—F9	1.263 (10)
Dy1—O4	2.342 (5)	C15—F8	1.278 (10)
Dy1—O8	2.349 (4)	C15—F7	1.313 (9)
Dy1—O7	2.356 (5)	O7—C17	1.250 (9)
Dy1—O5	2.390 (5)	O8—C19	1.275 (8)
Dy1—O1	2.423 (5)	C16—F10B	1.230 (14)
O1—C2	1.247 (8)	C16—F12B	1.314 (14)
O2—C4	1.269 (8)	C16—F11B	1.317 (14)
C1—C2	1.492 (10)	C16—C17	1.517 (12)
C1—H1A	0.9600	C16—H16A	0.9912
C1—H1B	0.9600	C16—H16B	1.0321
C1—H1C	0.9600	C16—H16C	1.0056
C2—C3	1.403 (10)	F10B—H16A	1.4259
C3—C4	1.350 (10)	F10B—H16B	0.5180
C3—H3	0.9300	F11B—H16A	0.6894
C4—C5	1.507 (11)	F12B—H16C	0.4623
C5—F1	1.275 (9)	C17—C18	1.392 (10)
C5—F2	1.295 (9)	C18—C19	1.363 (10)
C5—F3	1.317 (9)	C18—H18	0.9300
O3—C7	1.264 (8)	C19—C20	1.505 (10)
O4—C9	1.238 (11)	C20—F11	1.272 (11)
C6—C7	1.501 (11)	C20—F12	1.281 (11)
C6—H6A	0.9600	C20—F10	1.317 (10)
C6—H6B	0.9600	C20—H20A	1.1271
C6—H6C	0.9600	C20—H20B	0.9180
C7—C8	1.397 (12)	C20—H20C	1.0263
C8—C9	1.378 (13)	F10—H20C	0.7857
C8—H8	0.9300	F11—H20A	0.5838
C9—C10	1.582 (14)	F11—H20C	1.3102
C10—F4	1.213 (12)	F12—H20B	0.5114
C10—F4B	1.220 (14)	N1—C21	1.333 (11)
C10—F6B	1.321 (14)	N1—C25	1.350 (12)
C10—F6	1.328 (13)	N1—H1	0.8600
C10—F5	1.359 (13)	C21—C22	1.350 (11)
C10—F5B	1.372 (16)	C21—H21	0.9300

O5—C12	1.258 (8)	C22—C23	1.338 (11)
O6—C14	1.245 (8)	C22—H22	0.9300
C11—C12	1.525 (10)	C23—C24	1.358 (11)
C11—H11A	0.9600	C23—H23	0.9300
C11—H11B	0.9600	C24—C25	1.312 (11)
C11—H11C	0.9600	C24—H24	0.9300
C12—C13	1.405 (10)	C25—H25	0.9300
C13—C14	1.365 (10)		
O2—Dy1—O6	140.08 (17)	F6—C10—C9	110.8 (11)
O2—Dy1—O3	86.05 (17)	F5—C10—C9	109.1 (10)
O6—Dy1—O3	109.78 (17)	F5B—C10—C9	102.3 (13)
O2—Dy1—O4	78.12 (19)	C12—O5—Dy1	134.9 (5)
O6—Dy1—O4	73.0 (2)	C14—O6—Dy1	134.8 (5)
O3—Dy1—O4	71.84 (18)	C12—C11—H11A	109.5
O2—Dy1—O8	148.85 (17)	C12—C11—H11B	109.5
O6—Dy1—O8	71.07 (16)	H11A—C11—H11B	109.5
O3—Dy1—O8	79.62 (17)	C12—C11—H11C	109.5
O4—Dy1—O8	122.14 (17)	H11A—C11—H11C	109.5
O2—Dy1—O7	101.87 (18)	H11B—C11—H11C	109.5
O6—Dy1—O7	89.42 (18)	O5—C12—C13	123.2 (7)
O3—Dy1—O7	139.54 (17)	O5—C12—C11	118.0 (7)
O4—Dy1—O7	148.56 (18)	C13—C12—C11	118.7 (7)
O8—Dy1—O7	73.40 (16)	C14—C13—C12	125.3 (7)
O2—Dy1—O5	75.24 (17)	C14—C13—H13	117.4
O6—Dy1—O5	73.54 (16)	C12—C13—H13	117.4
O3—Dy1—O5	149.75 (18)	O6—C14—C13	127.8 (7)
O4—Dy1—O5	81.05 (18)	O6—C14—C15	113.3 (7)
O8—Dy1—O5	127.59 (17)	C13—C14—C15	119.0 (8)
O7—Dy1—O5	68.84 (17)	F9—C15—F8	107.6 (10)
O2—Dy1—O1	72.12 (16)	F9—C15—F7	104.5 (9)
O6—Dy1—O1	146.89 (17)	F8—C15—F7	103.9 (8)
O3—Dy1—O1	72.74 (17)	F9—C15—C14	112.7 (8)
O4—Dy1—O1	134.69 (19)	F8—C15—C14	111.7 (8)
O8—Dy1—O1	77.24 (16)	F7—C15—C14	115.7 (8)
O7—Dy1—O1	72.29 (17)	C17—O7—Dy1	130.6 (5)
O5—Dy1—O1	121.60 (16)	C19—O8—Dy1	127.7 (4)
C2—O1—Dy1	132.8 (5)	F10B—C16—F12B	112.0 (17)
C4—O2—Dy1	131.5 (4)	F10B—C16—F11B	108.9 (15)
C2—C1—H1A	109.5	F12B—C16—F11B	100.6 (16)
C2—C1—H1B	109.5	F10B—C16—C17	117.0 (12)
H1A—C1—H1B	109.5	F12B—C16—C17	110.0 (11)
C2—C1—H1C	109.5	F11B—C16—C17	106.8 (12)
H1A—C1—H1C	109.5	C17—C16—H16A	115.7
H1B—C1—H1C	109.5	C17—C16—H16B	114.7
O1—C2—C3	123.5 (7)	H16A—C16—H16B	101.5
O1—C2—C1	117.3 (7)	C17—C16—H16C	118.4
C3—C2—C1	119.2 (8)	H16A—C16—H16C	103.4
C4—C3—C2	124.2 (7)	H16B—C16—H16C	100.6
C4—C3—H3	117.9	H16A—F10B—H16B	96.0

supplementary materials

C2—C3—H3	117.9	C16—F11B—H16A	47.4
O2—C4—C3	128.1 (7)	O7—C17—C18	126.1 (7)
O2—C4—C5	112.7 (7)	O7—C17—C16	114.7 (8)
C3—C4—C5	119.2 (7)	C18—C17—C16	119.1 (8)
F1—C5—F2	106.4 (9)	C19—C18—C17	123.6 (7)
F1—C5—F3	105.0 (8)	C19—C18—H18	118.2
F2—C5—F3	103.1 (9)	C17—C18—H18	118.2
F1—C5—C4	113.5 (8)	O8—C19—C18	127.5 (6)
F2—C5—C4	112.8 (7)	O8—C19—C20	113.8 (7)
F3—C5—C4	115.0 (8)	C18—C19—C20	118.6 (7)
C7—O3—Dy1	132.6 (5)	F11—C20—F12	111.6 (13)
C9—O4—Dy1	130.5 (6)	F11—C20—F10	104.2 (11)
C7—C6—H6A	109.5	F12—C20—F10	103.2 (9)
C7—C6—H6B	109.5	F11—C20—C19	111.3 (8)
H6A—C6—H6B	109.5	F12—C20—C19	111.8 (9)
C7—C6—H6C	109.5	F10—C20—C19	114.4 (9)
H6A—C6—H6C	109.5	C19—C20—H20A	109.4
H6B—C6—H6C	109.5	C19—C20—H20B	121.4
O3—C7—C8	121.9 (8)	H20A—C20—H20B	99.6
O3—C7—C6	117.5 (8)	C19—C20—H20C	120.1
C8—C7—C6	120.6 (9)	H20A—C20—H20C	93.3
C9—C8—C7	123.6 (9)	H20B—C20—H20C	107.3
C9—C8—H8	118.2	C20—F10—H20C	51.2
C7—C8—H8	118.2	C20—F11—H20A	62.4
O4—C9—C8	127.8 (8)	C20—F11—H20C	46.8
O4—C9—C10	113.3 (9)	H20A—F11—H20C	105.2
C8—C9—C10	118.7 (10)	C21—N1—C25	121.7 (7)
F4—C10—F4B	136.8 (15)	C21—N1—H1	119.2
F4—C10—F6B	45.6 (9)	C25—N1—H1	119.2
F4B—C10—F6B	117.9 (15)	N1—C21—C22	118.8 (8)
F4—C10—F6	104.8 (13)	N1—C21—H21	120.6
F4B—C10—F6	54.0 (10)	C22—C21—H21	120.6
F6B—C10—F6	133.7 (13)	C23—C22—C21	119.7 (8)
F4—C10—F5	117.5 (14)	C23—C22—H22	120.1
F4B—C10—F5	52.5 (10)	C21—C22—H22	120.1
F6B—C10—F5	74.6 (12)	C22—C23—C24	120.3 (8)
F6—C10—F5	105.0 (12)	C22—C23—H23	119.8
F4—C10—F5B	57.9 (11)	C24—C23—H23	119.8
F4B—C10—F5B	106.0 (15)	C25—C24—C23	120.0 (9)
F6B—C10—F5B	102.1 (14)	C25—C24—H24	120.0
F6—C10—F5B	53.2 (11)	C23—C24—H24	120.0
F5—C10—F5B	147.1 (15)	C24—C25—N1	119.4 (8)
F4—C10—C9	109.5 (10)	C24—C25—H25	120.3
F4B—C10—C9	113.3 (11)	N1—C25—H25	120.3
F6B—C10—C9	112.8 (11)		
O2—Dy1—O1—C2	24.6 (6)	O7—Dy1—O6—C14	-67.3 (7)
O6—Dy1—O1—C2	-166.8 (6)	O5—Dy1—O6—C14	0.8 (6)
O3—Dy1—O1—C2	-66.9 (6)	O1—Dy1—O6—C14	-122.4 (6)
O4—Dy1—O1—C2	-26.9 (7)	Dy1—O5—C12—C13	9.0 (11)

O8—Dy1—O1—C2	-149.8 (7)	Dy1—O5—C12—C11	-168.6 (5)
O7—Dy1—O1—C2	133.8 (7)	O5—C12—C13—C14	-2.2 (12)
O5—Dy1—O1—C2	83.7 (7)	C11—C12—C13—C14	175.4 (8)
O6—Dy1—O2—C4	161.1 (6)	Dy1—O6—C14—C13	3.7 (12)
O3—Dy1—O2—C4	44.6 (6)	Dy1—O6—C14—C15	-176.3 (5)
O4—Dy1—O2—C4	116.9 (6)	C12—C13—C14—O6	-4.2 (13)
O8—Dy1—O2—C4	-17.8 (8)	C12—C13—C14—C15	175.7 (8)
O7—Dy1—O2—C4	-95.3 (6)	O6—C14—C15—F9	-54.8 (11)
O5—Dy1—O2—C4	-159.4 (6)	C13—C14—C15—F9	125.2 (9)
O1—Dy1—O2—C4	-28.5 (6)	O6—C14—C15—F8	66.5 (10)
Dy1—O1—C2—C1	169.3 (5)	C13—C14—C15—F8	-113.5 (9)
C1—C2—C3—C4	169.1 (8)	O6—C14—C15—F7	-175.0 (8)
Dy1—O2—C4—C5	-158.1 (5)	C13—C14—C15—F7	5.0 (12)
C2—C3—C4—C5	-174.7 (7)	O2—Dy1—O7—C17	177.5 (6)
C3—C4—C5—F1	112.4 (10)	O6—Dy1—O7—C17	-41.1 (6)
O2—C4—C5—F2	55.4 (10)	O3—Dy1—O7—C17	79.6 (7)
C3—C4—C5—F2	-126.4 (9)	O4—Dy1—O7—C17	-95.9 (7)
O2—C4—C5—F3	173.3 (8)	O8—Dy1—O7—C17	29.3 (6)
O2—Dy1—O3—C7	42.0 (7)	O5—Dy1—O7—C17	-113.6 (6)
O6—Dy1—O3—C7	-100.4 (7)	O1—Dy1—O7—C17	110.9 (6)
O4—Dy1—O3—C7	-36.8 (7)	O2—Dy1—O8—C19	-119.0 (6)
O8—Dy1—O3—C7	-165.8 (7)	O6—Dy1—O8—C19	61.8 (5)
O7—Dy1—O3—C7	145.7 (6)	O3—Dy1—O8—C19	177.0 (6)
O5—Dy1—O3—C7	-9.3 (9)	O4—Dy1—O8—C19	116.3 (5)
O1—Dy1—O3—C7	114.5 (7)	O7—Dy1—O8—C19	-33.5 (5)
O2—Dy1—O4—C9	-60.3 (7)	O5—Dy1—O8—C19	11.7 (6)
O6—Dy1—O4—C9	147.6 (7)	O1—Dy1—O8—C19	-108.5 (6)
O3—Dy1—O4—C9	29.4 (7)	Dy1—O7—C17—C18	-18.4 (11)
O8—Dy1—O4—C9	93.9 (7)	Dy1—O7—C17—C16	163.5 (6)
O7—Dy1—O4—C9	-153.7 (6)	F10B—C16—C17—O7	-176.4 (14)
O5—Dy1—O4—C9	-137.0 (7)	F12B—C16—C17—O7	54.3 (17)
O1—Dy1—O4—C9	-10.8 (8)	F11B—C16—C17—O7	-54.0 (14)
Dy1—O3—C7—C6	-148.8 (6)	F10B—C16—C17—C18	5.4 (18)
C6—C7—C8—C9	179.6 (9)	F12B—C16—C17—C18	-123.9 (16)
Dy1—O4—C9—C10	166.1 (6)	F11B—C16—C17—C18	127.7 (12)
C7—C8—C9—C10	169.6 (8)	O7—C17—C18—C19	-5.8 (12)
O4—C9—C10—F6B	-20.3 (15)	C16—C17—C18—C19	172.2 (8)
C8—C9—C10—F6B	164.5 (13)	Dy1—O8—C19—C18	28.9 (10)
O4—C9—C10—F6	175.6 (11)	Dy1—O8—C19—C20	-155.5 (6)
C8—C9—C10—F6	0.5 (15)	C17—C18—C19—O8	-0.1 (12)
O4—C9—C10—F5	60.6 (12)	C17—C18—C19—C20	-175.5 (7)
C8—C9—C10—F5	-114.6 (12)	O8—C19—C20—F11	46.0 (14)
O4—C9—C10—F5B	-129.3 (15)	C18—C19—C20—F11	-138.1 (12)
O2—Dy1—O5—C12	-162.0 (7)	O8—C19—C20—F12	-79.5 (12)
O6—Dy1—O5—C12	-7.2 (7)	C18—C19—C20—F12	96.5 (12)
O3—Dy1—O5—C12	-108.4 (7)	O8—C19—C20—F10	163.7 (8)
O4—Dy1—O5—C12	-82.0 (7)	C18—C19—C20—F10	-20.3 (13)
O8—Dy1—O5—C12	41.9 (7)	C25—N1—C21—C22	1.3 (13)
O7—Dy1—O5—C12	88.7 (7)	N1—C21—C22—C23	-0.7 (14)

supplementary materials

O1—Dy1—O5—C12	140.3 (6)	C21—C22—C23—C24	-1.4 (15)
O2—Dy1—O6—C14	40.7 (8)	C22—C23—C24—C25	3.0 (14)
O3—Dy1—O6—C14	149.1 (6)	C23—C24—C25—N1	-2.4 (14)
O4—Dy1—O6—C14	86.2 (7)	C21—N1—C25—C24	0.3 (14)
O8—Dy1—O6—C14	-139.9 (7)		

Hydrogen-bond geometry (\AA , °)

$D\text{—H}\cdots A$	$D\text{—H}$	$\text{H}\cdots A$	$D\cdots A$	$D\text{—H}\cdots A$
N1—H1…O5	0.86	2.10	2.947 (8)	167

Fig. 1

