

Triclinic, $P\bar{1}$	$V = 327.54 (5) \text{ \AA}^3$
$a = 6.3676 (6) \text{ \AA}$	$Z = 2$
$b = 7.3635 (7) \text{ \AA}$	Mo $K\alpha$ radiation
$c = 7.6798 (7) \text{ \AA}$	$\mu = 6.20 \text{ mm}^{-1}$
$\alpha = 65.554 (9)^\circ$	$T = 100 \text{ K}$
$\beta = 87.705 (8)^\circ$	$0.20 \times 0.10 \times 0.05 \text{ mm}$
$\gamma = 88.884 (8)^\circ$	

4-Bromo-1-nitrobenzene

Qamar Ali,^a M. Raza Shah^a and Seik Weng Ng^{b*}

^aH.E.J. Research Institute of Chemistry, International Center for Chemical and Biological Sciences, University of Karachi, Karachi 7527, Pakistan, and ^bDepartment of Chemistry, University of Malaya, 50603 Kuala Lumpur, Malaysia
Correspondence e-mail: seikweng@um.edu.my

Received 22 January 2011; accepted 25 January 2011

Key indicators: single-crystal X-ray study; $T = 100 \text{ K}$; mean $\sigma(\text{C}-\text{C}) = 0.006 \text{ \AA}$; R factor = 0.045; wR factor = 0.119; data-to-parameter ratio = 15.7.

The non-H atoms of the title molecule, $C_6H_4BrNO_2$, are essentially coplanar with an r.m.s. deviation of 0.040 \AA . In the crystal, $\pi-\pi$ stacking occurs between parallel benzene rings of adjacent molecules with centroid–centroid distances of 3.643 (3) and 3.741 (3) \AA . Weak intermolecular C–H \cdots O hydrogen bonding and short Br \cdots O contacts [3.227 (4) 3.401 (4) \AA] are also observed in the crystal structure. The crystal studied was a non-morohedral twin with a 26.1 (6)% minor component.

Related literature

For the structure of 2-bromonitrobenzene, see: Fronczek (2006). For the structure of 3-bromonitrobenzene, see: Charlton & Trotter (1963).



Experimental

Crystal data

$C_6H_4BrNO_2$

$M_r = 202.01$

Data collection

Agilent SuperNova Dual diffractometer with an Atlas detector
Absorption correction: multi-scan (*CrysAlis PRO*; Agilent, 2010)
 $T_{\min} = 0.414$, $T_{\max} = 1.000$

2142 measured reflections
1443 independent reflections
1365 reflections with $I > 2\sigma(I)$
 $R_{\text{int}} = 0.052$

Refinement

$R[F^2 > 2\sigma(F^2)] = 0.045$
 $wR(F^2) = 0.119$
 $S = 1.07$
1443 reflections

92 parameters
H-atom parameters constrained
 $\Delta\rho_{\max} = 0.91 \text{ e \AA}^{-3}$
 $\Delta\rho_{\min} = -1.58 \text{ e \AA}^{-3}$

Table 1
Hydrogen-bond geometry (\AA , $^\circ$).

$D-\text{H}\cdots A$	$D-\text{H}$	$\text{H}\cdots A$	$D\cdots A$	$D-\text{H}\cdots A$
C3—H3 \cdots O1 ⁱ	0.95	2.52	3.359 (6)	147
C5—H5 \cdots O2 ⁱⁱ	0.95	2.54	3.276 (6)	135

Symmetry codes: (i) $-x, -y + 1, -z + 1$; (ii) $-x + 1, -y, -z + 2$.

Data collection: *CrysAlis PRO* (Agilent, 2010); cell refinement: *CrysAlis PRO*; data reduction: *CrysAlis PRO*; program(s) used to solve structure: *SHELXS97* (Sheldrick, 2008); program(s) used to refine structure: *SHELXL97* (Sheldrick, 2008); molecular graphics: *X-SEED* (Barbour, 2001); software used to prepare material for publication: *publCIF* (Westrip, 2010).

We thank the Higher Education Commission of Pakistan and the University of Malaya for supporting this study.

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: XU5150).

References

- Agilent (2010). *CrysAlis PRO*. Agilent Technologies, Yarnton, England.
Barbour, L. J. (2001). *J. Supramol. Chem.* **1**, 189–191.
Charlton, T. L. & Trotter, J. (1963). *Acta Cryst.* **16**, 313.
Fronczek, F. R. (2006). Private communication (refcode 264855). CCDC, Cambridge, England.
Sheldrick, G. M. (2008). *Acta Cryst. A* **64**, 112–122.
Westrip, S. P. (2010). *J. Appl. Cryst.* **43**, 920–925.

supplementary materials

Acta Cryst. (2011). E67, o548 [doi:10.1107/S1600536811003394]

4-Bromo-1-nitrobenzene

Q. Ali, M. Raza Shah and S. W. Ng

Comment

4-Bromo-1-nitrobenzene (Scheme I) was synthesized as a precursor that will be used in the synthesis of 4,4'-bis(aminophenoxy)biphenyl (the compound is also commercially available: <http://www.chemindustry.com/chemicals/815494.html>). The molecule is flat (Fig. 1) as the nitro substituent is co-planar with the aromatic ring. π - π stacking occurs between parallel benzene rings of adjacent molecules, centroids distance between C1-ring and C1ⁱ-ring (symmetry code: (i) 1-x, -5, 1-z) is 3.643 (3) Å and that between C1-ring and C1ⁱⁱ-ring (symmetry code: (ii) 1-x, 1-y, 1-z) is 3.741 (3) Å. Intermolecular weak C—H \cdots O hydrogen bonding (Table 1) and the short Br \cdots O contacts [3.227 (4), 3.401 (4) Å] are observed in the crystal structure.

Experimental

The nitrating mixture consisted of 5 ml conc. HNO₃ and 5 ml conc. H₂SO₄ kept at 273 K. Bromobenzene (2.6 ml) was added. The temperature was then raised to about 333 K for 3 h. The mixture was added to water (200 ml); the organic compound was extracted by using dichloromethane. The solvent was dried and then allowed to evaporate to yield the product in 70% yield.

Refinement

Carbon-bound H-atoms were placed in calculated positions [C—H 0.95 Å, $U_{\text{iso}}(\text{H})$ 1.2 $U_{\text{eq}}(\text{C})$] and were included in the refinement in the riding model approximation.

The crystal is a non-merohedral twin; the separation of the two domains was effected by *CrysAlis PRO* (Agilent, 2010).

Figures




Fig. 1. Thermal ellipsoid plot (Barbour, 2001) of C₆H₄BrNO₂ at the 70% probability level; hydrogen atoms are drawn as spheres of arbitrary radius.

4-Bromo-1-nitrobenzene

Crystal data

C ₆ H ₄ BrNO ₂	Z = 2
M_r = 202.01	$F(000)$ = 196
Triclinic, $P\bar{1}$	D_x = 2.048 Mg m ⁻³

supplementary materials

Hall symbol: -P 1

$a = 6.3676 (6)$ Å

$b = 7.3635 (7)$ Å

$c = 7.6798 (7)$ Å

$\alpha = 65.554 (9)^\circ$

$\beta = 87.705 (8)^\circ$

$\gamma = 88.884 (8)^\circ$

$V = 327.54 (5)$ Å³

Mo $K\alpha$ radiation, $\lambda = 0.71073$ Å

Cell parameters from 1590 reflections

$\theta = 2.9\text{--}28.3^\circ$

$\mu = 6.20$ mm⁻¹

$T = 100$ K

Block, colorless

$0.20 \times 0.10 \times 0.05$ mm

Data collection

Agilent SuperNova Dual
diffractometer with an Atlas detector

1443 independent reflections

Radiation source: SuperNova (Mo) X-ray Source

1365 reflections with $I > 2\sigma(I)$

Mirror

$R_{\text{int}} = 0.052$

Detector resolution: 10.4041 pixels mm⁻¹

$\theta_{\text{max}} = 27.5^\circ, \theta_{\text{min}} = 2.9^\circ$

ω scans

$h = -8 \rightarrow 8$

Absorption correction: multi-scan
(*CrysAlis PRO*; Agilent, 2010)

$k = -9 \rightarrow 9$

$T_{\text{min}} = 0.414, T_{\text{max}} = 1.000$

$l = -9 \rightarrow 9$

2142 measured reflections

Refinement

Refinement on F^2

Primary atom site location: structure-invariant direct
methods

Least-squares matrix: full

Secondary atom site location: difference Fourier map

$R[F^2 > 2\sigma(F^2)] = 0.045$

Hydrogen site location: inferred from neighbouring
sites

$wR(F^2) = 0.119$

H-atom parameters constrained

$S = 1.07$

$w = 1/[\sigma^2(F_o^2) + (0.0671P)^2 + 0.4717P]$

where $P = (F_o^2 + 2F_c^2)/3$

1443 reflections

$(\Delta/\sigma)_{\text{max}} = 0.001$

92 parameters

$\Delta\rho_{\text{max}} = 0.91$ e Å⁻³

0 restraints

$\Delta\rho_{\text{min}} = -1.58$ e Å⁻³

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (Å²)

	x	y	z	$U_{\text{iso}}^*/U_{\text{eq}}$
Br1	0.83556 (7)	0.24763 (7)	0.14671 (6)	0.01978 (19)
O1	0.0573 (5)	0.3167 (6)	0.7390 (5)	0.0238 (8)
O2	0.3071 (6)	0.2117 (6)	0.9389 (5)	0.0245 (8)
N1	0.2383 (6)	0.2623 (6)	0.7787 (5)	0.0151 (7)
C6	0.7151 (7)	0.1751 (7)	0.5288 (7)	0.0173 (9)
H6	0.8526	0.1212	0.5575	0.021*
C5	0.5810 (7)	0.1811 (7)	0.6722 (6)	0.0146 (9)
H5	0.6251	0.1326	0.8004	0.018*
C2	0.4465 (7)	0.3256 (7)	0.2969 (7)	0.0173 (9)

H2	0.4024	0.3750	0.1687	0.021*
C3	0.3121 (7)	0.3291 (7)	0.4408 (6)	0.0159 (9)
H3	0.1731	0.3792	0.4130	0.019*
C1	0.6478 (7)	0.2483 (6)	0.3429 (6)	0.0157 (9)
C4	0.3809 (7)	0.2594 (6)	0.6254 (6)	0.0125 (8)

Atomic displacement parameters (\AA^2)

	U^{11}	U^{22}	U^{33}	U^{12}	U^{13}	U^{23}
Br1	0.0204 (3)	0.0213 (3)	0.0206 (3)	-0.00244 (19)	0.00662 (18)	-0.0122 (2)
O1	0.0149 (16)	0.038 (2)	0.0232 (18)	0.0024 (15)	0.0001 (13)	-0.0174 (16)
O2	0.0299 (19)	0.032 (2)	0.0131 (16)	0.0028 (16)	-0.0003 (14)	-0.0107 (15)
N1	0.0173 (18)	0.0152 (18)	0.0156 (18)	-0.0013 (14)	0.0010 (14)	-0.0093 (15)
C6	0.015 (2)	0.017 (2)	0.021 (2)	-0.0008 (17)	0.0000 (17)	-0.0089 (19)
C5	0.016 (2)	0.015 (2)	0.014 (2)	0.0014 (16)	-0.0023 (16)	-0.0076 (17)
C2	0.020 (2)	0.018 (2)	0.016 (2)	-0.0004 (18)	0.0001 (17)	-0.0094 (18)
C3	0.018 (2)	0.016 (2)	0.015 (2)	0.0020 (17)	-0.0041 (17)	-0.0072 (17)
C1	0.019 (2)	0.014 (2)	0.019 (2)	-0.0005 (18)	0.0015 (18)	-0.0113 (19)
C4	0.0133 (19)	0.015 (2)	0.012 (2)	-0.0005 (16)	0.0005 (15)	-0.0090 (17)

Geometric parameters (\AA , °)

Br1—C1	1.887 (4)	C5—C4	1.387 (6)
O1—N1	1.220 (5)	C5—H5	0.9500
O2—N1	1.226 (5)	C2—C3	1.379 (6)
N1—C4	1.464 (5)	C2—C1	1.390 (6)
C6—C1	1.384 (6)	C2—H2	0.9500
C6—C5	1.381 (6)	C3—C4	1.380 (6)
C6—H6	0.9500	C3—H3	0.9500
O1—N1—O2	123.6 (4)	C1—C2—H2	120.6
O1—N1—C4	117.9 (4)	C4—C3—C2	119.6 (4)
O2—N1—C4	118.4 (4)	C4—C3—H3	120.2
C1—C6—C5	119.5 (4)	C2—C3—H3	120.2
C1—C6—H6	120.2	C6—C1—C2	121.5 (4)
C5—C6—H6	120.2	C6—C1—Br1	119.2 (3)
C4—C5—C6	118.7 (4)	C2—C1—Br1	119.3 (3)
C4—C5—H5	120.6	C3—C4—C5	121.8 (4)
C6—C5—H5	120.6	C3—C4—N1	119.7 (4)
C3—C2—C1	118.8 (4)	C5—C4—N1	118.4 (4)
C3—C2—H2	120.6		
C1—C6—C5—C4	0.5 (7)	C2—C3—C4—N1	-179.8 (4)
C1—C2—C3—C4	1.1 (7)	C6—C5—C4—C3	0.7 (7)
C5—C6—C1—C2	-0.9 (7)	C6—C5—C4—N1	179.0 (4)
C5—C6—C1—Br1	177.9 (3)	O1—N1—C4—C3	4.1 (6)
C3—C2—C1—C6	0.1 (7)	O2—N1—C4—C3	-175.3 (4)
C3—C2—C1—Br1	-178.7 (3)	O1—N1—C4—C5	-174.3 (4)
C2—C3—C4—C5	-1.5 (7)	O2—N1—C4—C5	6.3 (6)

supplementary materials

Hydrogen-bond geometry (Å, °)

<i>D—H···A</i>	<i>D—H</i>	<i>H···A</i>	<i>D···A</i>	<i>D—H···A</i>
C3—H3···O1 ⁱ	0.95	2.52	3.359 (6)	147
C5—H5···O2 ⁱⁱ	0.95	2.54	3.276 (6)	135

Symmetry codes: (i) $-x, -y+1, -z+1$; (ii) $-x+1, -y, -z+2$.

Fig. 1

