

Dimethyl 5,6,7-trimethoxy-2-methyl-1,2-dihydroquinoline-2,4-dicarboxylate

Zeynep Gültekin,^a Wolfgang Frey^b and Tuncer Hökelek^{c*}

^aDepartment of Chemistry, Çankırı Karatekin University, TR-18100 Çankırı, Turkey,

^bUniversität Stuttgart, Pfaffenwaldring 55, D-70569 Stuttgart, Germany, and

^cDepartment of Physics, Hacettepe University, 06800 Beytepe Ankara, Turkey

Correspondence e-mail: merzifon@hacettepe.edu.tr


Received 27 January 2011; accepted 1 February 2011

Key indicators: single-crystal X-ray study; $T = 294$ K; mean $\sigma(\text{C}-\text{C}) = 0.005$ Å; R factor = 0.070; wR factor = 0.156; data-to-parameter ratio = 14.5.

In the title compound, $\text{C}_{17}\text{H}_{21}\text{NO}_7$, the dihydropyridine ring assumes a screw-boat conformation. In the crystal, intermolecular $\text{C}-\text{H}\cdots\text{O}$ hydrogen bonds link the molecules, forming supramolecular chains running along the b axis.

Related literature

For the preparation of 1,2-dihydroquinoline, see: Edwards *et al.* (1998); Yan *et al.* (2004); Petasis & Butkevich (2009); Johnson *et al.* (1989); Gültekin *et al.* (2010); Waldmann *et al.* (2008). For the biological activity of dihydroquinolines, see: Elmore *et al.* (2001); Dillard *et al.* (1973); Muren & Weissmann (1971). For ring puckering parameters, see: Cremer & Pople (1975).



Experimental

Crystal data

$\text{C}_{17}\text{H}_{21}\text{NO}_7$

$M_r = 351.35$

Orthorhombic, $Pbca$

$a = 10.476$ (2) Å

$b = 16.552$ (4) Å

$c = 20.238$ (4) Å

$V = 3509.2$ (13) Å³

$Z = 8$

Mo $K\alpha$ radiation

$\mu = 0.10$ mm⁻¹

$T = 294$ K

$0.6 \times 0.4 \times 0.15$ mm

Data collection

Nicolet P3 diffractometer

3447 measured reflections

3447 independent reflections

1839 reflections with $I > 2\sigma(I)$

3 standard reflections every 50

reflections

intensity decay: 1%

Refinement

$R[F^2 > 2\sigma(F^2)] = 0.070$

$wR(F^2) = 0.156$

$S = 1.07$

3447 reflections

237 parameters

H atoms treated by a mixture of

independent and constrained

refinement

$\Delta\rho_{\text{max}} = 0.18$ e Å⁻³

$\Delta\rho_{\text{min}} = -0.17$ e Å⁻³

Table 1

Hydrogen-bond geometry (Å, °).

$D-H\cdots A$	$D-H$	$H\cdots A$	$D\cdots A$	$D-H\cdots A$
$\text{C14}-\text{H14A}\cdots\text{O1}^i$	0.96	2.51	3.251 (6)	134

Symmetry code: (i) $-x + \frac{1}{2}, y - \frac{1}{2}, z$.

Data collection: XSCANS (Siemens, 1996); cell refinement: XSCANS; data reduction: SHELXTL (Sheldrick, 2008); program(s) used to solve structure: SHELXS97 (Sheldrick, 2008); program(s) used to refine structure: SHELXL97 (Sheldrick, 2008); molecular graphics: ORTEP-3 for Windows (Farrugia, 1997); software used to prepare material for publication: WinGX (Farrugia, 1999).

This research was carried out at RWTH Aachen University. The authors thank Professor Magnus Rueping of RWTH Aachen University, Germany, for helpful discussions.

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: XU5153).

References

- Cremer, D. & Pople, J. A. (1975). *J. Am. Chem. Soc.* **97**, 1354–1358.
- Dillard, R. D., Pavey, D. E. & Benslay, D. N. (1973). *J. Med. Chem.* **16**, 251–253.
- Edwards, J. P., Ringenberg, J. D. & Jones, T. K. (1998). *Tetrahedron Lett.* **39**, 5139–5142.
- Elmore, S. W., Coghlan, M. J., Anderson, D. D., Pratt, J. K., Green, B. E., Wang, A. X., Stashko, M. A., Lin, C. W., Tyree, C. M., Miner, J. N., Jacobson, P. B., Wilcox, D. M. & Lane, B. C. (2001). *J. Med. Chem.* **44**, 4481–4491.
- Farrugia, L. J. (1997). *J. Appl. Cryst.* **30**, 565.
- Farrugia, L. J. (1999). *J. Appl. Cryst.* **32**, 837–838.
- Gültekin, Z., Frey, W., Tercan, B. & Hökelek, T. (2010). *Acta Cryst.* **E66**, o2891–o2892.
- Johnson, J. V., Rauckman, B. S., Baccanari, D. P. & Roth, B. (1989). *J. Med. Chem.* **32**, 1942–1949.
- Muren, J. F. & Weissmann, A. (1971). *J. Med. Chem.* **14**, 49–53.
- Petasis, N. A. & Butkevich, A. N. (2009). *J. Organomet. Chem.* **694**, 1747–1753.
- Sheldrick, G. M. (2008). *Acta Cryst.* **A64**, 112–122.
- Siemens (1996). XSCANS. Siemens Analytical X-ray Instruments Inc., Madison, Wisconsin, USA.
- Waldmann, H., Karunakar, G. V. & Kumar, K. (2008). *Org. Lett.* **10**, 2159–2162.
- Yan, M.-C., Tu, Z.-J., Lin, C.-C., Ko, S.-K., Hsu, J.-M. & Yao, C.-F. (2004). *J. Org. Chem.* **69**, 1565–1570.

supplementary materials

Acta Cryst. (2011). E67, o576 [doi:10.1107/S1600536811004028]

Dimethyl 5,6,7-trimethoxy-2-methyl-1,2-dihydroquinoline-2,4-dicarboxylate

Z. Gültekin, W. Frey and T. Hökelek

Comment

Dihydroquinolines have been widely studied and found an important structural unit in synthetic organic and medicinal chemistry (Elmore *et al.*, 2001; Dillard *et al.*, 1973; Muren & Weissmann, 1971). Many dihydroquinoline derivatives have been reported in the literature (Edwards *et al.*, 1998; Yan *et al.*, 2004; Petasis & Butkevich, 2009; Gültekin *et al.*, 2010) and some of them have biological effects. For example, 2,2,4-substituted 1,2-dihydroquinolines have been shown antibacterial activities (Johnson *et al.*, 1989).

In the title compound, (I), (Fig. 1), the ring A (C1-C4/C9/N1) is not planar; the puckering parameters (Cremer & Pople, 1975) $Q_T = 0.379$ (3) Å, $\varphi = 21.5$ (6)° and $\theta = 66.4$ (5)° suggesting a screw-boat conformation. In the crystal structure, intermolecular C-H...O hydrogen bonds (Table 1) link the molecules to form infinite chains along the b-axis (Fig. 2).

Experimental

The title compound was synthesized by the literature method (Waldmann *et al.*, 2008). 3,4,5-dimethoxyaniline (100 mg, 1 eq) was dissolved in chloroform (1.5 ml) in a screw-capped test tube and Bi(OTf)₃ (5 mol%, 0.05 eq) was added to the mixture. The mixture was stirred at room temperature for 4 h until the starting material was completely consumed as monitored by TLC. The resultant residue was directly purified by flash chromatography on silica (EtOAc:Cyclohexane 2:98) gave in 83% yield as a yellow solid. Recrystallized over pentane and ethyl acetate (70:30) gave yellow crystalline solid R_f 0.16 (2:1 Cyclohexane/EtOAc) mp 394-395 K.

Figures




Fig. 1. The molecular structure of the title compound with the atom-numbering scheme. Displacement ellipsoids are drawn at the 50% probability level.




Fig. 2. A partial packing diagram viewed down the c-axis. Hydrogen bonds are shown as dashed lines.

Dimethyl 5,6,7-trimethoxy-2-methyl-1,2-dihydroquinoline-2,4-dicarboxylate

Crystal data

$C_{17}H_{21}NO_7$	$F(000) = 1488$
$M_r = 351.35$	$D_x = 1.330 \text{ Mg m}^{-3}$
Orthorhombic, <i>Pbca</i>	Mo $K\alpha$ radiation, $\lambda = 0.71073 \text{ \AA}$
Hall symbol: -P 2ac 2ab	Cell parameters from 40 reflections
$a = 10.476 (2) \text{ \AA}$	$\theta = 10\text{--}12^\circ$
$b = 16.552 (4) \text{ \AA}$	$\mu = 0.10 \text{ mm}^{-1}$
$c = 20.238 (4) \text{ \AA}$	$T = 294 \text{ K}$
$V = 3509.2 (13) \text{ \AA}^3$	Plates, colourless
$Z = 8$	$0.6 \times 0.4 \times 0.15 \text{ mm}$

Data collection

Nicolet P3 diffractometer	$R_{\text{int}} = 0.000$
Radiation source: fine-focus sealed tube graphite	$\theta_{\text{max}} = 26.0^\circ$, $\theta_{\text{min}} = 2.0^\circ$
Wyckoff scan	$h = 0 \rightarrow 12$
3447 measured reflections	$k = 0 \rightarrow 20$
3447 independent reflections	$l = 0 \rightarrow 24$
1839 reflections with $I > 2\sigma(I)$	3 standard reflections every 50 reflections
	intensity decay: 1%

Refinement

Refinement on F^2	Secondary atom site location: difference Fourier map
Least-squares matrix: full	Hydrogen site location: inferred from neighbouring sites
$R[F^2 > 2\sigma(F^2)] = 0.070$	H atoms treated by a mixture of independent and constrained refinement
$wR(F^2) = 0.156$	$w = 1/[\sigma^2(F_o^2) + (0.0359P)^2 + 3.112P]$
$S = 1.07$	where $P = (F_o^2 + 2F_c^2)/3$
3447 reflections	$(\Delta/\sigma)_{\text{max}} < 0.001$
237 parameters	$\Delta\rho_{\text{max}} = 0.18 \text{ e \AA}^{-3}$
0 restraints	$\Delta\rho_{\text{min}} = -0.17 \text{ e \AA}^{-3}$
Primary atom site location: structure-invariant direct methods	Extinction correction: <i>SHELXL97</i> (Sheldrick, 2008), $F_c^* = kFc[1 + 0.001xFc^2\lambda^3/\sin(2\theta)]^{-1/4}$
	Extinction coefficient: 0.0031 (3)

Special details

Geometry. All esds (except the esd in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell esds are taken into account individually in the estimation of esds in distances, angles and torsion angles; correlations between esds

in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell esds is used for estimating esds involving l.s. planes.

Refinement. Refinement of F^2 against ALL reflections. The weighted R-factor wR and goodness of fit S are based on F^2 , conventional R-factors R are based on F, with F set to zero for negative F^2 . The threshold expression of $F^2 > 2\sigma(F^2)$ is used only for calculating R-factors(gt) etc. and is not relevant to the choice of reflections for refinement. R-factors based on F^2 are statistically about twice as large as those based on F, and R- factors based on ALL data will be even larger.

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (\AA^2)

	x	y	z	$U_{\text{iso}}^*/U_{\text{eq}}$
O1	0.5908 (3)	1.03546 (16)	0.62717 (15)	0.0881 (10)
O2	0.6233 (3)	0.93649 (16)	0.55547 (13)	0.0713 (8)
O3	0.2547 (3)	0.7747 (2)	0.51994 (16)	0.1002 (11)
O4	0.1503 (3)	0.75844 (14)	0.61564 (14)	0.0745 (8)
O5	0.0584 (2)	0.91803 (14)	0.58046 (11)	0.0540 (6)
O6	-0.0366 (2)	1.05816 (15)	0.63970 (12)	0.0637 (7)
O7	0.1046 (3)	1.13852 (16)	0.72647 (13)	0.0736 (8)
N1	0.4420 (3)	0.9400 (2)	0.70665 (15)	0.0590 (9)
H21	0.485 (4)	0.973 (2)	0.733 (2)	0.091 (16)*
C1	0.5191 (3)	0.9018 (2)	0.65584 (19)	0.0525 (9)
C2	0.4333 (3)	0.8467 (2)	0.61660 (18)	0.0536 (9)
H2	0.4696	0.8031	0.5946	0.064*
C3	0.3086 (3)	0.85768 (19)	0.61218 (16)	0.0479 (8)
C4	0.2492 (3)	0.92839 (19)	0.64312 (15)	0.0445 (8)
C5	0.1281 (3)	0.9578 (2)	0.62794 (16)	0.0457 (8)
C6	0.0824 (3)	1.0285 (2)	0.65599 (17)	0.0489 (9)
C7	0.1567 (3)	1.0694 (2)	0.70186 (17)	0.0541 (9)
C8	0.2776 (3)	1.0412 (2)	0.71861 (17)	0.0566 (10)
H8	0.3272	1.0688	0.7493	0.068*
C9	0.3233 (3)	0.9712 (2)	0.68871 (16)	0.0489 (9)
C10	0.6265 (4)	0.8538 (2)	0.6884 (2)	0.0730 (12)
H10A	0.6801	0.8899	0.7132	0.109*
H10B	0.6763	0.8274	0.6550	0.109*
H10C	0.5907	0.8141	0.7176	0.109*
C11	0.5798 (3)	0.9663 (2)	0.61161 (19)	0.0528 (9)
C12	0.6849 (5)	0.9931 (3)	0.5108 (2)	0.1013 (16)
H12A	0.6974	0.9678	0.4686	0.152*
H12B	0.7660	1.0088	0.5287	0.152*
H12C	0.6319	1.0400	0.5055	0.152*
C13	0.2345 (4)	0.7940 (2)	0.5758 (2)	0.0613 (10)
C14	0.0782 (5)	0.6925 (3)	0.5864 (3)	0.118 (2)
H14A	0.0260	0.6676	0.6197	0.177*
H14B	0.1362	0.6532	0.5686	0.177*
H14C	0.0249	0.7130	0.5517	0.177*
C15	-0.0622 (3)	0.8868 (2)	0.6031 (2)	0.0755 (12)
H15A	-0.1022	0.8569	0.5682	0.113*
H15B	-0.1164	0.9308	0.6160	0.113*
H15C	-0.0483	0.8519	0.6403	0.113*

supplementary materials

C16	-0.0349 (5)	1.1165 (3)	0.5882 (2)	0.0906 (15)
H16A	-0.1179	1.1406	0.5842	0.136*
H16B	-0.0124	1.0907	0.5474	0.136*
H16C	0.0268	1.1576	0.5984	0.136*
C17	0.1819 (4)	1.1883 (2)	0.7671 (2)	0.0896 (15)
H17A	0.1360	1.2367	0.7781	0.134*
H17B	0.2587	1.2022	0.7438	0.134*
H17C	0.2033	1.1597	0.8069	0.134*

Atomic displacement parameters (\AA^2)

	U^{11}	U^{22}	U^{33}	U^{12}	U^{13}	U^{23}
O1	0.111 (3)	0.0508 (17)	0.103 (2)	-0.0080 (17)	0.014 (2)	0.0001 (16)
O2	0.0779 (19)	0.0685 (17)	0.0676 (18)	-0.0126 (15)	0.0092 (15)	-0.0002 (15)
O3	0.101 (2)	0.115 (3)	0.085 (2)	-0.028 (2)	0.012 (2)	-0.047 (2)
O4	0.0813 (18)	0.0507 (15)	0.092 (2)	-0.0248 (15)	-0.0035 (17)	-0.0025 (15)
O5	0.0507 (14)	0.0598 (15)	0.0516 (14)	-0.0123 (12)	-0.0065 (12)	-0.0034 (12)
O6	0.0493 (15)	0.0704 (17)	0.0712 (18)	0.0042 (13)	0.0024 (13)	-0.0002 (14)
O7	0.0750 (19)	0.0670 (17)	0.0789 (19)	0.0009 (15)	0.0051 (15)	-0.0253 (15)
N1	0.0520 (19)	0.073 (2)	0.0524 (19)	-0.0034 (18)	-0.0093 (16)	-0.0077 (17)
C1	0.050 (2)	0.048 (2)	0.060 (2)	0.0010 (18)	-0.0078 (18)	0.0047 (18)
C2	0.056 (2)	0.0443 (19)	0.060 (2)	0.0016 (18)	-0.0018 (19)	0.0009 (17)
C3	0.055 (2)	0.0426 (19)	0.046 (2)	-0.0088 (17)	-0.0019 (17)	0.0038 (16)
C4	0.0478 (18)	0.0465 (18)	0.0391 (17)	-0.0091 (17)	0.0019 (16)	-0.0005 (16)
C5	0.0453 (19)	0.052 (2)	0.0395 (18)	-0.0142 (17)	-0.0014 (16)	-0.0009 (16)
C6	0.0425 (19)	0.053 (2)	0.051 (2)	-0.0064 (17)	0.0045 (17)	-0.0020 (17)
C7	0.055 (2)	0.052 (2)	0.055 (2)	-0.0057 (19)	0.0102 (19)	-0.0077 (18)
C8	0.058 (2)	0.063 (2)	0.049 (2)	-0.013 (2)	0.0029 (18)	-0.0144 (19)
C9	0.047 (2)	0.056 (2)	0.043 (2)	-0.0094 (18)	-0.0022 (17)	0.0025 (17)
C10	0.064 (2)	0.069 (3)	0.086 (3)	0.003 (2)	-0.017 (2)	0.020 (2)
C11	0.045 (2)	0.050 (2)	0.063 (2)	0.0026 (18)	-0.0061 (19)	0.0045 (19)
C12	0.104 (4)	0.115 (4)	0.085 (3)	-0.029 (3)	0.017 (3)	0.027 (3)
C13	0.060 (2)	0.054 (2)	0.070 (3)	-0.008 (2)	-0.001 (2)	-0.009 (2)
C14	0.106 (4)	0.072 (3)	0.176 (6)	-0.041 (3)	-0.005 (4)	-0.029 (3)
C15	0.050 (2)	0.075 (3)	0.102 (3)	-0.016 (2)	-0.006 (2)	-0.009 (2)
C16	0.094 (3)	0.072 (3)	0.106 (4)	0.006 (3)	-0.017 (3)	0.021 (3)
C17	0.104 (4)	0.073 (3)	0.092 (4)	-0.006 (3)	0.000 (3)	-0.038 (3)

Geometric parameters (\AA , $^\circ$)

O1—C11	1.193 (4)	C5—C6	1.386 (5)
O2—C11	1.320 (4)	C6—C7	1.387 (5)
O2—C12	1.453 (5)	C7—C8	1.391 (5)
O3—C13	1.194 (4)	C8—C9	1.393 (5)
O4—C13	1.332 (4)	C8—H8	0.9300
O4—C14	1.453 (5)	C10—H10A	0.9600
O5—C5	1.374 (4)	C10—H10B	0.9600
O5—C15	1.440 (4)	C10—H10C	0.9600
O6—C6	1.380 (4)	C12—H12A	0.9600

O6—C16	1.421 (5)	C12—H12B	0.9600
O7—C7	1.362 (4)	C12—H12C	0.9600
O7—C17	1.418 (4)	C14—H14A	0.9600
N1—C1	1.453 (4)	C14—H14B	0.9600
N1—C9	1.394 (4)	C14—H14C	0.9600
N1—H21	0.89 (4)	C15—H15A	0.9600
C1—C2	1.507 (5)	C15—H15B	0.9600
C1—C10	1.527 (5)	C15—H15C	0.9600
C1—C11	1.532 (5)	C16—H16A	0.9600
C2—C3	1.322 (4)	C16—H16B	0.9600
C2—H2	0.9300	C16—H16C	0.9600
C3—C4	1.466 (5)	C17—H17A	0.9600
C3—C13	1.502 (5)	C17—H17B	0.9600
C4—C5	1.393 (5)	C17—H17C	0.9600
C4—C9	1.399 (4)		
C11—O2—C12	116.6 (3)	C1—C10—H10C	109.5
C13—O4—C14	115.5 (4)	H10A—C10—H10B	109.5
C5—O5—C15	114.5 (3)	H10A—C10—H10C	109.5
C6—O6—C16	113.9 (3)	H10B—C10—H10C	109.5
C7—O7—C17	118.1 (3)	O1—C11—O2	123.6 (4)
C1—N1—H21	114 (3)	O1—C11—C1	123.7 (4)
C9—N1—C1	118.2 (3)	O2—C11—C1	112.7 (3)
C9—N1—H21	112 (3)	O2—C12—H12A	109.5
N1—C1—C2	107.7 (3)	O2—C12—H12B	109.5
N1—C1—C10	109.3 (3)	O2—C12—H12C	109.5
N1—C1—C11	109.9 (3)	H12A—C12—H12B	109.5
C2—C1—C10	110.7 (3)	H12A—C12—H12C	109.5
C2—C1—C11	111.2 (3)	H12B—C12—H12C	109.5
C10—C1—C11	108.0 (3)	O3—C13—O4	124.9 (4)
C1—C2—H2	118.6	O3—C13—C3	124.1 (4)
C3—C2—C1	122.8 (3)	O4—C13—C3	110.9 (3)
C3—C2—H2	118.6	O4—C14—H14A	109.5
C2—C3—C4	120.1 (3)	O4—C14—H14B	109.5
C2—C3—C13	116.6 (3)	O4—C14—H14C	109.5
C4—C3—C13	123.4 (3)	H14A—C14—H14B	109.5
C5—C4—C3	124.9 (3)	H14A—C14—H14C	109.5
C5—C4—C9	118.3 (3)	H14B—C14—H14C	109.5
C9—C4—C3	116.7 (3)	O5—C15—H15A	109.5
O5—C5—C4	118.1 (3)	O5—C15—H15B	109.5
O5—C5—C6	120.5 (3)	O5—C15—H15C	109.5
C6—C5—C4	121.3 (3)	H15A—C15—H15B	109.5
O6—C6—C5	120.9 (3)	H15A—C15—H15C	109.5
O6—C6—C7	119.6 (3)	H15B—C15—H15C	109.5
C5—C6—C7	119.5 (3)	O6—C16—H16A	109.5
O7—C7—C6	115.4 (3)	O6—C16—H16B	109.5
O7—C7—C8	123.9 (3)	O6—C16—H16C	109.5
C6—C7—C8	120.7 (3)	H16A—C16—H16B	109.5
C7—C8—C9	119.1 (3)	H16A—C16—H16C	109.5
C7—C8—H8	120.5	H16B—C16—H16C	109.5

supplementary materials

C9—C8—H8	120.5	O7—C17—H17A	109.5
N1—C9—C4	118.7 (3)	O7—C17—H17B	109.5
C8—C9—N1	120.1 (3)	O7—C17—H17C	109.5
C8—C9—C4	121.1 (3)	H17A—C17—H17B	109.5
C1—C10—H10A	109.5	H17A—C17—H17C	109.5
C1—C10—H10B	109.5	H17B—C17—H17C	109.5
C12—O2—C11—O1	-1.3 (6)	C2—C3—C4—C9	-14.3 (5)
C12—O2—C11—C1	-178.8 (3)	C13—C3—C4—C5	-19.0 (5)
C14—O4—C13—O3	-1.0 (6)	C13—C3—C4—C9	164.4 (3)
C14—O4—C13—C3	-176.6 (3)	C2—C3—C13—O3	-57.5 (6)
C15—O5—C5—C4	120.6 (3)	C2—C3—C13—O4	118.1 (4)
C15—O5—C5—C6	-64.2 (4)	C4—C3—C13—O3	123.8 (5)
C16—O6—C6—C5	-94.5 (4)	C4—C3—C13—O4	-60.6 (4)
C16—O6—C6—C7	86.5 (4)	C3—C4—C5—O5	-0.1 (5)
C17—O7—C7—C6	-172.3 (3)	C3—C4—C5—C6	-175.2 (3)
C17—O7—C7—C8	5.6 (5)	C9—C4—C5—O5	176.5 (3)
C9—N1—C1—C2	-45.4 (4)	C9—C4—C5—C6	1.3 (5)
C9—N1—C1—C10	-165.7 (3)	C3—C4—C9—N1	-6.3 (4)
C9—N1—C1—C11	76.0 (4)	C3—C4—C9—C8	177.2 (3)
C1—N1—C9—C8	-145.0 (3)	C5—C4—C9—N1	176.8 (3)
C1—N1—C9—C4	38.5 (5)	C5—C4—C9—C8	0.3 (5)
N1—C1—C2—C3	24.7 (5)	O5—C5—C6—O6	3.6 (5)
C10—C1—C2—C3	144.2 (4)	O5—C5—C6—C7	-177.4 (3)
C11—C1—C2—C3	-95.8 (4)	C4—C5—C6—O6	178.7 (3)
N1—C1—C11—O1	18.8 (5)	C4—C5—C6—C7	-2.3 (5)
N1—C1—C11—O2	-163.7 (3)	C5—C6—C7—O7	179.6 (3)
C2—C1—C11—O1	138.0 (4)	C5—C6—C7—C8	1.7 (5)
C2—C1—C11—O2	-44.5 (4)	O6—C6—C7—O7	-1.4 (5)
C10—C1—C11—O1	-100.4 (4)	O6—C6—C7—C8	-179.3 (3)
C10—C1—C11—O2	77.1 (4)	O7—C7—C8—C9	-177.8 (3)
C1—C2—C3—C4	3.5 (5)	C6—C7—C8—C9	0.0 (5)
C1—C2—C3—C13	-175.2 (3)	C7—C8—C9—N1	-177.4 (3)
C2—C3—C4—C5	162.3 (3)	C7—C8—C9—C4	-1.0 (5)

Hydrogen-bond geometry (\AA , $^\circ$)

$D-H\cdots A$	$D-H$	$H\cdots A$	$D\cdots A$	$D-H\cdots A$
C14—H14A \cdots O1 ⁱ	0.96	2.51	3.251 (6)	134

Symmetry codes: (i) $-x+1/2, y-1/2, z$.

Fig. 1




Fig. 2

