

4-Bromo-2-[(*E*)-(4-fluorophenyl)imino-methyl]phenol

Amir Adabi Ardakani,^a Reza Kia,^{b*} Hadi Kargar^c and Muhammad Nawaz Tahir^{d*}

^aIslamic Azad University, Ardakan Branch, Iran, ^bX-ray Crystallography Laboratory, Plasma Physics Research Center, Science and Research Branch, Islamic Azad University, Tehran, Iran, ^cChemistry Department, Payame Noor University, Tehran 19395-4697, I. R. of Iran, and ^dDepartment of Physics, University of Sargodha, Punjab, Pakistan

Correspondence e-mail: zsrkk@yahoo.com, rkia@srbiau.ac.ir, dmntahir_uos@yahoo.com


Received 2 February 2011; accepted 6 February 2011

Key indicators: single-crystal X-ray study; $T = 296\text{ K}$; mean $\sigma(\text{C}-\text{C}) = 0.003\text{ \AA}$; R factor = 0.032; wR factor = 0.083; data-to-parameter ratio = 18.1.

In the title compound, $\text{C}_{13}\text{H}_9\text{BrFNO}$, the dihedral angle between the substituted benzene rings is $9.00(11)^\circ$. Strong intramolecular $\text{O}-\text{H} \cdots \text{N}$ hydrogen bonds generate $S(6)$ ring motifs.

Related literature

For standard bond lengths, see: Allen *et al.* (1987). For hydrogen-bond motifs, see: Bernstein *et al.* (1995).



Experimental

Crystal data

$\text{C}_{13}\text{H}_9\text{BrFNO}$

$M_r = 294.12$

Monoclinic, $P2_1/n$
 $a = 4.4820(2)\text{ \AA}$
 $b = 20.8088(9)\text{ \AA}$
 $c = 12.2189(5)\text{ \AA}$
 $\beta = 94.570(2)^\circ$
 $V = 1135.97(8)\text{ \AA}^3$

$Z = 4$
Mo $K\alpha$ radiation
 $\mu = 3.61\text{ mm}^{-1}$
 $T = 296\text{ K}$
 $0.35 \times 0.17 \times 0.11\text{ mm}$

Data collection

Bruker SMART APEXII CCD area-detector diffractometer
Absorption correction: multi-scan (*SADABS*; Bruker, 2005)
 $S = 0.365$, $T_{\max} = 0.692$

10561 measured reflections
2792 independent reflections
1958 reflections with $I > 2\sigma(I)$
 $R_{\text{int}} = 0.034$

Refinement

$R[F^2 > 2\sigma(F^2)] = 0.032$
 $wR(F^2) = 0.083$
 $S = 1.02$
2792 reflections

154 parameters
H-atom parameters constrained
 $\Delta\rho_{\text{max}} = 0.44\text{ e \AA}^{-3}$
 $\Delta\rho_{\text{min}} = -0.26\text{ e \AA}^{-3}$

Table 1
Hydrogen-bond geometry (\AA , $^\circ$).

$D-\text{H} \cdots A$	$D-\text{H}$	$\text{H} \cdots A$	$D \cdots A$	$D-\text{H} \cdots A$
O1—H1 \cdots N1	0.82	1.89	2.612 (2)	146

Data collection: *APEX2* (Bruker, 2005); cell refinement: *SAINT* (Bruker, 2005); data reduction: *SAINT*; program(s) used to solve structure: *SHELXTL* (Sheldrick, 2008); program(s) used to refine structure: *SHELXTL*; molecular graphics: *SHELXTL*; software used to prepare material for publication: *SHELXTL* and *PLATON* (Spek, 2009).

AAA thanks the Islamic Azad University, Ardakan Branch for the research facilities (this paper was extracted from a research project supported by IAU). RK thanks the Science and Research Branch, Islamic Azad University, Tehran. HK thanks the PNU for financial support. MNT thanks Sargodha University for the research facilities.

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: JH2266).

References

- Allen, F. H., Kennard, O., Watson, D. G., Brammer, L., Orpen, A. G. & Taylor, R. (1987). *J. Chem. Soc. Perkin Trans. 2*, pp. S1–19.
Bernstein, J., Davis, R. E., Shimoni, L. & Chang, N.-L. (1995). *Angew. Chem. Int. Ed. Engl.* **34**, 1555–1573.
Bruker (2005). *APEX2*, *SAINT* and *SADABS*. Bruker AXS Inc., Madison, Wisconsin, USA.
Sheldrick, G. M. (2008). *Acta Cryst. A* **64**, 112–122.
Spek, A. L. (2009). *Acta Cryst. D* **65**, 148–155.

supplementary materials

Acta Cryst. (2011). E67, o598 [doi:10.1107/S1600536811004429]

4-Bromo-2-[(E)-(4-fluorophenyl)iminomethyl]phenol

A. A. Ardakani, R. Kia, H. Kargar and M. N. Tahir

Comment

Schiff base ligands are one of the most prevalent systems in coordination chemistry. As part of a general study of Schiff bases, we have determined the crystal structure of the title compound.

The asymmetric unit of the title compound, Fig. 1, comprises a potentially bidentate Schiff base ligand. The bond lengths (Allen *et al.*, 1987) and angles are within the normal ranges. The dihedral angle between the substituted benzene rings is 9.00 (11) Å. Strong intramolecular O—H···N hydrogen bonds generate *S*(6) ring motifs (Bernstein *et al.*, 1995).

Experimental

The title compound was synthesized by adding 5-bromo-salicylaldehyde (2 mmol) to a solution of *p*-fluoroaniline (2 mmol) in ethanol (20 ml). The mixture was refluxed with stirring for half an hour. The resulting light-yellow solution was filtered. Pale-yellow single crystals suitable for *X*-ray diffraction were recrystallized from ethanol by slow evaporation of the solvents at room temperature over several days.

Refinement

H atoms of the hydroxy groups were located by a rotating model and constrained to refine with the parent atoms with $U_{\text{iso}}(\text{H}) = 1.5 U_{\text{eq}}(\text{O})$, see Table 1. The remaining H atoms were positioned geometrically with C—H = 0.93 Å and included in a riding model approximation with $U_{\text{iso}}(\text{H}) = 1.2 U_{\text{eq}}(\text{C})$.

Figures




Fig. 1. The asymmetric unit of the title compound, showing 40% probability displacement ellipsoids and the atomic numbering. Intramolecular hydrogen bond is drawn as dashed lines.




Fig. 2. The packing diagram of the title compound, viewed down the *c*-axis.

supplementary materials

4-Bromo-2-[(*E*)-(4-fluorophenyl)iminomethyl]phenol

Crystal data

C ₁₃ H ₉ BrFNO	<i>F</i> (000) = 584
<i>M_r</i> = 294.12	<i>D_x</i> = 1.720 Mg m ⁻³
Monoclinic, <i>P2₁/n</i>	Mo <i>Kα</i> radiation, λ = 0.71073 Å
Hall symbol: -P 2yn	Cell parameters from 2520 reflections
<i>a</i> = 4.4820 (2) Å	θ = 2.5–28.5°
<i>b</i> = 20.8088 (9) Å	μ = 3.61 mm ⁻¹
<i>c</i> = 12.2189 (5) Å	<i>T</i> = 296 K
β = 94.570 (2)°	Prism, pale-yellow
<i>V</i> = 1135.97 (8) Å ³	0.35 × 0.17 × 0.11 mm
<i>Z</i> = 4	

Data collection

Bruker SMART APEXII CCD area-detector diffractometer	2792 independent reflections
Radiation source: fine-focus sealed tube	1958 reflections with $I > 2\sigma(I)$
graphite	$R_{\text{int}} = 0.034$
φ and ω scans	$\theta_{\text{max}} = 28.3^\circ$, $\theta_{\text{min}} = 1.9^\circ$
Absorption correction: multi-scan (<i>SADABS</i> ; Bruker, 2005)	$h = -5 \rightarrow 5$
$T_{\text{min}} = 0.365$, $T_{\text{max}} = 0.692$	$k = -27 \rightarrow 27$
10561 measured reflections	$l = -16 \rightarrow 16$

Refinement

Refinement on F^2	Primary atom site location: structure-invariant direct methods
Least-squares matrix: full	Secondary atom site location: difference Fourier map
$R[F^2 > 2\sigma(F^2)] = 0.032$	Hydrogen site location: inferred from neighbouring sites
$wR(F^2) = 0.083$	H-atom parameters constrained
$S = 1.02$	$w = 1/[\sigma^2(F_o^2) + (0.0413P)^2 + 0.1115P]$ where $P = (F_o^2 + 2F_c^2)/3$
2792 reflections	$(\Delta/\sigma)_{\text{max}} = 0.001$
154 parameters	$\Delta\rho_{\text{max}} = 0.44 \text{ e \AA}^{-3}$
0 restraints	$\Delta\rho_{\text{min}} = -0.26 \text{ e \AA}^{-3}$

Special details

Geometry. All esds (except the esd in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell esds are taken into account individually in the estimation of esds in distances, angles and torsion angles; correlations between esds

in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell esds is used for estimating esds involving l.s. planes.

Refinement. Refinement of F^2 against ALL reflections. The weighted R-factor wR and goodness of fit S are based on F^2 , conventional R-factors R are based on F, with F set to zero for negative F^2 . The threshold expression of $F^2 > 2\text{sigma}(F^2)$ is used only for calculating R-factors(gt) etc. and is not relevant to the choice of reflections for refinement. R-factors based on F^2 are statistically about twice as large as those based on F, and R-factors based on ALL data will be even larger.

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (\AA^2)

	x	y	z	$U_{\text{iso}}^*/U_{\text{eq}}$
Br1	1.14491 (6)	0.420695 (13)	0.55664 (2)	0.06007 (13)
F1	-0.3896 (4)	-0.01392 (7)	0.62967 (14)	0.0714 (5)
O1	0.7822 (4)	0.23124 (8)	0.89264 (12)	0.0502 (4)
H1	0.6582	0.2057	0.8661	0.075*
N1	0.4202 (4)	0.17993 (8)	0.73875 (14)	0.0357 (4)
C1	0.7310 (5)	0.26970 (10)	0.70681 (16)	0.0322 (5)
C2	0.8605 (5)	0.27266 (11)	0.81485 (16)	0.0356 (5)
C3	1.0742 (5)	0.31885 (11)	0.84342 (17)	0.0415 (5)
H3	1.1617	0.3204	0.9150	0.050*
C4	1.1590 (5)	0.36250 (11)	0.76728 (18)	0.0418 (5)
H4	1.3019	0.3936	0.7873	0.050*
C5	1.0300 (5)	0.35970 (10)	0.66087 (18)	0.0377 (5)
C6	0.8198 (5)	0.31402 (10)	0.63019 (17)	0.0369 (5)
H6	0.7360	0.3126	0.5581	0.044*
C7	0.5052 (4)	0.22206 (10)	0.67230 (17)	0.0343 (5)
H7	0.4212	0.2224	0.6002	0.041*
C8	0.2060 (4)	0.13210 (10)	0.70442 (17)	0.0332 (5)
C9	0.1021 (5)	0.12019 (11)	0.59655 (18)	0.0418 (5)
H9	0.1690	0.1454	0.5406	0.050*
C10	-0.0999 (5)	0.07125 (11)	0.57144 (19)	0.0460 (6)
H10	-0.1702	0.0633	0.4990	0.055*
C11	-0.1946 (5)	0.03478 (11)	0.6549 (2)	0.0445 (6)
C12	-0.0996 (5)	0.04499 (12)	0.7619 (2)	0.0483 (6)
H12	-0.1684	0.0195	0.8171	0.058*
C13	0.1021 (5)	0.09426 (12)	0.78670 (19)	0.0443 (6)
H13	0.1687	0.1021	0.8595	0.053*

Atomic displacement parameters (\AA^2)

	U^{11}	U^{22}	U^{33}	U^{12}	U^{13}	U^{23}
Br1	0.0733 (2)	0.05159 (18)	0.05448 (18)	-0.01654 (13)	-0.00037 (13)	0.01097 (12)
F1	0.0782 (11)	0.0552 (10)	0.0788 (11)	-0.0331 (9)	-0.0069 (9)	0.0023 (8)
O1	0.0592 (10)	0.0587 (11)	0.0313 (8)	-0.0167 (8)	-0.0051 (7)	0.0033 (8)
N1	0.0310 (10)	0.0401 (11)	0.0354 (9)	0.0009 (8)	-0.0014 (7)	-0.0038 (8)
C1	0.0303 (11)	0.0348 (11)	0.0312 (10)	0.0019 (9)	0.0005 (8)	-0.0036 (9)
C2	0.0348 (12)	0.0405 (12)	0.0315 (11)	0.0018 (9)	0.0023 (9)	-0.0039 (9)
C3	0.0422 (13)	0.0505 (14)	0.0305 (11)	-0.0031 (11)	-0.0043 (9)	-0.0084 (10)

supplementary materials

C4	0.0393 (13)	0.0388 (13)	0.0467 (13)	-0.0048 (10)	-0.0005 (10)	-0.0125 (10)
C5	0.0418 (13)	0.0314 (11)	0.0402 (12)	0.0017 (10)	0.0045 (10)	0.0001 (9)
C6	0.0382 (12)	0.0387 (12)	0.0327 (11)	0.0021 (10)	-0.0036 (9)	-0.0026 (9)
C7	0.0326 (11)	0.0372 (12)	0.0322 (10)	0.0025 (9)	-0.0028 (9)	-0.0055 (9)
C8	0.0275 (11)	0.0361 (12)	0.0357 (11)	0.0019 (9)	-0.0004 (9)	-0.0001 (9)
C9	0.0495 (14)	0.0409 (13)	0.0344 (11)	-0.0059 (11)	-0.0005 (10)	0.0009 (10)
C10	0.0517 (15)	0.0451 (14)	0.0397 (12)	-0.0089 (11)	-0.0066 (11)	-0.0024 (10)
C11	0.0407 (13)	0.0364 (13)	0.0554 (14)	-0.0050 (10)	-0.0027 (11)	0.0008 (11)
C12	0.0501 (15)	0.0462 (15)	0.0492 (14)	-0.0058 (12)	0.0068 (11)	0.0128 (11)
C13	0.0425 (13)	0.0523 (15)	0.0370 (12)	-0.0029 (11)	-0.0035 (10)	0.0038 (10)

Geometric parameters (\AA , $^\circ$)

Br1—C5	1.898 (2)	C5—C6	1.370 (3)
F1—C11	1.358 (3)	C6—H6	0.9300
O1—C2	1.350 (2)	C7—H7	0.9300
O1—H1	0.8176	C8—C9	1.385 (3)
N1—C7	1.274 (3)	C8—C13	1.387 (3)
N1—C8	1.423 (3)	C9—C10	1.381 (3)
C1—C6	1.395 (3)	C9—H9	0.9300
C1—C2	1.400 (3)	C10—C11	1.366 (3)
C1—C7	1.455 (3)	C10—H10	0.9300
C2—C3	1.382 (3)	C11—C12	1.359 (3)
C3—C4	1.375 (3)	C12—C13	1.384 (3)
C3—H3	0.9300	C12—H12	0.9300
C4—C5	1.381 (3)	C13—H13	0.9300
C4—H4	0.9300		
C2—O1—H1	110.1	N1—C7—H7	119.3
C7—N1—C8	121.51 (18)	C1—C7—H7	119.3
C6—C1—C2	118.99 (19)	C9—C8—C13	118.8 (2)
C6—C1—C7	119.02 (19)	C9—C8—N1	125.01 (19)
C2—C1—C7	121.99 (19)	C13—C8—N1	116.18 (18)
O1—C2—C3	118.68 (18)	C10—C9—C8	120.5 (2)
O1—C2—C1	121.60 (19)	C10—C9—H9	119.7
C3—C2—C1	119.7 (2)	C8—C9—H9	119.7
C4—C3—C2	120.78 (19)	C11—C10—C9	118.8 (2)
C4—C3—H3	119.6	C11—C10—H10	120.6
C2—C3—H3	119.6	C9—C10—H10	120.6
C3—C4—C5	119.4 (2)	F1—C11—C12	118.8 (2)
C3—C4—H4	120.3	F1—C11—C10	118.5 (2)
C5—C4—H4	120.3	C12—C11—C10	122.7 (2)
C6—C5—C4	120.9 (2)	C11—C12—C13	118.4 (2)
C6—C5—Br1	119.87 (16)	C11—C12—H12	120.8
C4—C5—Br1	119.20 (17)	C13—C12—H12	120.8
C5—C6—C1	120.14 (19)	C12—C13—C8	120.9 (2)
C5—C6—H6	119.9	C12—C13—H13	119.6
C1—C6—H6	119.9	C8—C13—H13	119.6
N1—C7—C1	121.38 (19)		
C6—C1—C2—O1	179.65 (18)	C6—C1—C7—N1	178.87 (18)

C7—C1—C2—O1	0.1 (3)	C2—C1—C7—N1	-1.6 (3)
C6—C1—C2—C3	-0.4 (3)	C7—N1—C8—C9	9.7 (3)
C7—C1—C2—C3	180.0 (2)	C7—N1—C8—C13	-171.9 (2)
O1—C2—C3—C4	-179.3 (2)	C13—C8—C9—C10	-0.4 (3)
C1—C2—C3—C4	0.8 (3)	N1—C8—C9—C10	178.1 (2)
C2—C3—C4—C5	-0.5 (3)	C8—C9—C10—C11	-0.2 (4)
C3—C4—C5—C6	-0.2 (3)	C9—C10—C11—F1	-179.0 (2)
C3—C4—C5—Br1	179.31 (17)	C9—C10—C11—C12	0.5 (4)
C4—C5—C6—C1	0.5 (3)	F1—C11—C12—C13	179.2 (2)
Br1—C5—C6—C1	-179.00 (15)	C10—C11—C12—C13	-0.2 (4)
C2—C1—C6—C5	-0.2 (3)	C11—C12—C13—C8	-0.3 (4)
C7—C1—C6—C5	179.4 (2)	C9—C8—C13—C12	0.6 (4)
C8—N1—C7—C1	-178.11 (18)	N1—C8—C13—C12	-177.9 (2)

Hydrogen-bond geometry (Å, °)

D—H···A	D—H	H···A	D···A	D—H···A
O1—H1···N1	0.82	1.89	2.612 (2)	146

supplementary materials

Fig. 1




Fig. 2

