

4,4'-Dimethoxy-2,2'-[2,2-dimethyl-propane-1,3-diylbis(nitrilomethanylidyne)]diphenol

Hadi Kargar,^a Reza Kia,^{b*} Elham Pahlavani^a and Muhammad Nawaz Tahir^{c*}

^aChemistry Department, Payame Noor University, Tehran 19395-4697, I. R. of Iran,

^bX-ray Crystallography Laboratory, Plasma Physics Research Center, Science and Research Branch, Islamic Azad University, Tehran, Iran, and ^cDepartment of Physics, University of Sargodha, Punjab, Pakistan

Correspondence e-mail: rkia@srbiau.ac.ir, zsrkk@yahoo.com, dmntahir_uos@yahoo.com


Received 30 January 2011; accepted 8 February 2011

Key indicators: single-crystal X-ray study; $T = 296\text{ K}$; mean $\sigma(\text{C}-\text{C}) = 0.004\text{ \AA}$; R factor = 0.031; wR factor = 0.058; data-to-parameter ratio = 13.7.

In the title compound, $C_{21}H_{26}N_2O_4$, the dihedral angle between the substituted benzene rings is $30.47(15)^\circ$. Two strong intramolecular O—H···N hydrogen bonds generate two $S(6)$ ring motifs.

Related literature

For hydrogen-bond motifs, see: Bernstein *et al.* (1995). For related structures, see: Kargar *et al.* (2009, 2010).



Experimental

Crystal data



$M_r = 370.44$

Monoclinic, $P2_1/c$

$a = 10.660(2)\text{ \AA}$

$b = 21.742(4)\text{ \AA}$

$c = 9.2767(19)\text{ \AA}$

$\beta = 108.03(3)^\circ$

$V = 2044.5(7)\text{ \AA}^3$

$Z = 4$
Mo $K\alpha$ radiation
 $\mu = 0.08\text{ mm}^{-1}$

$T = 296\text{ K}$
 $0.23 \times 0.15 \times 0.08\text{ mm}$

Data collection

Stoe IPDS 2T Image Plate diffractometer
Absorption correction: multi-scan (*MULABS* in *PLATON*; Blessing, 1995)
 $T_{\min} = 0.965$, $T_{\max} = 1.000$

7094 measured reflections
3375 independent reflections
967 reflections with $I > 2\sigma(I)$
 $R_{\text{int}} = 0.054$

Refinement

$R[F^2 > 2\sigma(F^2)] = 0.031$
 $wR(F^2) = 0.058$
 $S = 0.57$
3375 reflections

246 parameters
H-atom parameters constrained
 $\Delta\rho_{\max} = 0.08\text{ e \AA}^{-3}$
 $\Delta\rho_{\min} = -0.11\text{ e \AA}^{-3}$

Table 1
Hydrogen-bond geometry (\AA , $^\circ$).

$D-\text{H}\cdots A$	$D-\text{H}$	$\text{H}\cdots A$	$D\cdots A$	$D-\text{H}\cdots A$
O1—H1···N1	0.81	1.88	2.593 (3)	147
O2—H2···N2	0.83	1.90	2.604 (3)	143

Data collection: *X-AREA* (Stoe & Cie, 2009); cell refinement: *X-AREA*; data reduction: *X-AREA*; program(s) used to solve structure: *SHELXTL* (Sheldrick, 2008); program(s) used to refine structure: *SHELXTL*; molecular graphics: *SHELXTL*; software used to prepare material for publication: *SHELXTL* and *PLATON* (Spek, 2009).

HK and EP thank PNU for financial support. RK thanks the Science and Research Branch, Islamic Azad University, Tehran. MNT thanks Sargodha University for the research facilities. RK thanks the Chemistry Department, University of Isfahan, for the use of Stoe IPDS 2T diffractometer facility

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: TK2716).

References

- Bernstein, J., Davis, R. E., Shimoni, L. & Chang, N.-L. (1995). *Angew. Chem. Int. Ed. Engl.* **34**, 1555–1573.
Blessing, R. H. (1995). *Acta Cryst. A* **51**, 33–38.
Kargar, H., Kia, R., Jamshidvand, A. & Fun, H.-K. (2009). *Acta Cryst. E* **65**, o776–o777.
Kargar, H., Kia, R., Ullah Khan, I. & Sahraei, A. (2010). *Acta Cryst. E* **66**, o539.
Sheldrick, G. M. (2008). *Acta Cryst. A* **64**, 112–122.
Spek, A. L. (2009). *Acta Cryst. D* **65**, 148–155.
Stoe & Cie (2009). *X-AREA* Stoe & Cie, Darmstadt, Germany.

supplementary materials

Acta Cryst. (2011). E67, o614 [doi:10.1107/S1600536811004776]

4,4'-Dimethoxy-2,2'-[2,2-dimethylpropane-1,3-diylbis(nitrilomethanylylidene)]diphenol

H. Kargar, R. Kia, E. Pahlavani and M. N. Tahir

Comment

Schiff base ligands are one of the most prevalent systems in coordination chemistry. As part of a general study of potentially tetradenate Schiff bases (Kargar *et al.*, 2009; Kargar *et al.*, 2010), we have determined the crystal structure of the title compound.

The asymmetric unit of the title compound, Fig. 1, comprises a potentially tetradenate Schiff base ligand. The bond lengths are comparable to previously reported structures (Kargar *et al.*, 2009; Kargar *et al.*, 2010). The dihedral angle between the two benzene rings is 30.47 (15) °. Strong intramolecular O—H···N hydrogen bonds (Table 1) generate two S(6) ring motifs (Bernstein *et al.*, 1995).

Experimental

The title compound was synthesized by adding 5-methoxy-salicylaldehyde (4 mmol) to a solution of 2,2-dimethyl-1,3-propanediamine (2 mmol) in ethanol (20 ml). The mixture was refluxed with stirring for 30 min. The resultant yellow solution was filtered. Yellow crystals were obtained by slow evaporation of its ethanol solution at room temperature over several days.

Refinement

H atoms of the hydroxy groups were located in a difference Fourier map and constrained at those positions with $U_{\text{iso}}(\text{H}) = 1.5 U_{\text{eq}}(\text{O})$, see Table 1 for distances. The remaining H atoms were positioned geometrically with C—H = 0.93–0.97 Å and included in a riding model approximation with $U_{\text{iso}}(\text{H}) = 1.2$ or $1.5 U_{\text{eq}}(\text{C})$. A rotating group model was used only for the methyl groups of the methoxy substituents.

Figures




Fig. 1. The molecular structure of the title compound, showing 30% probability displacement ellipsoids and the atomic numbering. Intramolecular hydrogen bonds are drawn as dashed lines.

supplementary materials

4,4'-Dimethoxy-2,2'-[2,2-dimethylpropane-1,3- diylbis(nitrilomethanlylidene)]diphenol

Crystal data

C ₂₁ H ₂₆ N ₂ O ₄	$F(000) = 792$
$M_r = 370.44$	$D_x = 1.204 \text{ Mg m}^{-3}$
Monoclinic, $P2_1/c$	Mo $K\alpha$ radiation, $\lambda = 0.71073 \text{ \AA}$
Hall symbol: -P 2ybc	Cell parameters from 3220 reflections
$a = 10.660 (2) \text{ \AA}$	$\theta = 2.0\text{--}24.2^\circ$
$b = 21.742 (4) \text{ \AA}$	$\mu = 0.08 \text{ mm}^{-1}$
$c = 9.2767 (19) \text{ \AA}$	$T = 296 \text{ K}$
$\beta = 108.03 (3)^\circ$	Plate, yellow
$V = 2044.5 (7) \text{ \AA}^3$	$0.23 \times 0.15 \times 0.08 \text{ mm}$
$Z = 4$	

Data collection

Stoe IPDS 2T Image Plate diffractometer	3375 independent reflections
Radiation source: fine-focus sealed tube graphite	967 reflections with $I > 2\sigma(I)$
Detector resolution: 0.15 mm pixels mm ⁻¹	$R_{\text{int}} = 0.054$
ω scans	$\theta_{\text{max}} = 25.0^\circ, \theta_{\text{min}} = 2.0^\circ$
Absorption correction: multi-scan (<i>MULABS</i> in <i>PLATON</i> ; Blessing, 1995)	$h = -12\text{--}12$
$T_{\text{min}} = 0.965, T_{\text{max}} = 1.000$	$k = -22\text{--}25$
7094 measured reflections	$l = -11\text{--}10$

Refinement

Refinement on F^2	Primary atom site location: structure-invariant direct methods
Least-squares matrix: full	Secondary atom site location: difference Fourier map
$R[F^2 > 2\sigma(F^2)] = 0.031$	Hydrogen site location: inferred from neighbouring sites
$wR(F^2) = 0.058$	H-atom parameters constrained
$S = 0.57$	$w = 1/[\sigma^2(F_o^2) + (0.0172P)^2]$ where $P = (F_o^2 + 2F_c^2)/3$
3375 reflections	$(\Delta/\sigma)_{\text{max}} = 0.001$
246 parameters	$\Delta\rho_{\text{max}} = 0.08 \text{ e \AA}^{-3}$
0 restraints	$\Delta\rho_{\text{min}} = -0.11 \text{ e \AA}^{-3}$

Special details

Geometry. All e.s.d.'s (except the e.s.d. in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell e.s.d.'s are taken into account individually in the estimation of e.s.d.'s in distances, angles and torsion angles; correlations

between e.s.d.'s in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell e.s.d.'s is used for estimating e.s.d.'s involving l.s. planes.

Refinement. Refinement of F^2 against ALL reflections. The weighted R -factor wR and goodness of fit S are based on F^2 , conventional R -factors R are based on F , with F set to zero for negative F^2 . The threshold expression of $F^2 > \sigma(F^2)$ is used only for calculating R -factors(gt) etc. and is not relevant to the choice of reflections for refinement. R -factors based on F^2 are statistically about twice as large as those based on F , and R -factors based on ALL data will be even larger.

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (\AA^2)

	<i>x</i>	<i>y</i>	<i>z</i>	$U_{\text{iso}}^*/U_{\text{eq}}$
O1	-0.05846 (17)	0.01529 (7)	0.75824 (19)	0.0807 (6)
H1	-0.0997	0.0250	0.6727	0.121*
O2	-0.1706 (2)	0.31943 (8)	0.4769 (2)	0.0874 (6)
H2	-0.2189	0.2892	0.4703	0.131*
O3	0.4634 (2)	0.07957 (10)	0.8988 (2)	0.1064 (8)
O4	0.3081 (2)	0.30533 (10)	0.9255 (3)	0.0936 (7)
N1	-0.1073 (3)	0.07562 (9)	0.5057 (3)	0.0629 (7)
N2	-0.2263 (3)	0.21006 (10)	0.5539 (3)	0.0662 (7)
C1	0.0688 (3)	0.03241 (12)	0.7858 (4)	0.0601 (8)
C2	0.1605 (3)	0.01492 (12)	0.9224 (3)	0.0685 (8)
H2A	0.1336	-0.0083	0.9917	0.082*
C3	0.2893 (3)	0.03161 (12)	0.9549 (3)	0.0750 (9)
H3A	0.3502	0.0195	1.0460	0.090*
C4	0.3300 (3)	0.06678 (14)	0.8524 (4)	0.0720 (9)
C5	0.2413 (3)	0.08523 (12)	0.7179 (3)	0.0705 (9)
H5A	0.2691	0.1088	0.6498	0.085*
C6	0.1091 (3)	0.06829 (11)	0.6844 (3)	0.0557 (8)
C7	0.0141 (3)	0.08981 (11)	0.5447 (3)	0.0621 (8)
H7A	0.0435	0.1151	0.4811	0.075*
C8	-0.1960 (3)	0.10405 (11)	0.3705 (3)	0.0689 (8)
H8A	-0.2106	0.0760	0.2856	0.083*
H8B	-0.1554	0.1410	0.3468	0.083*
C9	-0.3286 (3)	0.12057 (12)	0.3915 (3)	0.0647 (8)
C10	-0.3096 (2)	0.15556 (12)	0.5410 (3)	0.0686 (8)
H10A	-0.3951	0.1680	0.5473	0.082*
H10B	-0.2698	0.1282	0.6255	0.082*
C11	-0.1179 (3)	0.21062 (13)	0.6614 (3)	0.0647 (9)
H11A	-0.0969	0.1777	0.7287	0.078*
C12	-0.0261 (3)	0.26170 (13)	0.6810 (3)	0.0550 (8)
C13	-0.0544 (3)	0.31384 (15)	0.5877 (3)	0.0673 (9)
C14	0.0387 (4)	0.36057 (13)	0.6113 (4)	0.0794 (11)
H14A	0.0205	0.3954	0.5500	0.095*
C15	0.1556 (4)	0.35591 (14)	0.7224 (4)	0.0804 (10)
H15A	0.2167	0.3876	0.7359	0.096*
C16	0.1861 (4)	0.30500 (14)	0.8166 (4)	0.0677 (9)
C17	0.0946 (3)	0.25820 (12)	0.7948 (3)	0.0628 (8)
H17A	0.1140	0.2238	0.8573	0.075*
C18	-0.4084 (2)	0.06177 (12)	0.3967 (3)	0.0982 (10)

supplementary materials

H18A	-0.3598	0.0366	0.4803	0.147*
H18B	-0.4234	0.0393	0.3038	0.147*
H18C	-0.4916	0.0729	0.4092	0.147*
C19	-0.4063 (3)	0.16042 (12)	0.2556 (3)	0.0975 (10)
H19A	-0.3572	0.1971	0.2522	0.146*
H19B	-0.4901	0.1713	0.2666	0.146*
H19C	-0.4199	0.1376	0.1634	0.146*
C20	0.5161 (3)	0.11165 (14)	0.7998 (4)	0.1184 (13)
H20A	0.6098	0.1155	0.8448	0.178*
H20B	0.4771	0.1518	0.7810	0.178*
H20C	0.4974	0.0896	0.7059	0.178*
C21	0.3454 (3)	0.25225 (13)	1.0160 (3)	0.1204 (13)
H21A	0.4318	0.2581	1.0868	0.181*
H21B	0.3463	0.2175	0.9525	0.181*
H21C	0.2834	0.2450	1.0703	0.181*

Atomic displacement parameters (\AA^2)

	U^{11}	U^{22}	U^{33}	U^{12}	U^{13}	U^{23}
O1	0.0710 (15)	0.0916 (13)	0.0825 (14)	-0.0025 (12)	0.0279 (14)	0.0249 (11)
O2	0.1028 (17)	0.0767 (14)	0.0822 (16)	0.0121 (13)	0.0279 (14)	0.0188 (12)
O3	0.0738 (17)	0.154 (2)	0.0852 (18)	-0.0283 (16)	0.0154 (16)	0.0102 (14)
O4	0.0945 (18)	0.0784 (16)	0.0994 (19)	-0.0159 (14)	0.0174 (16)	0.0104 (14)
N1	0.0633 (17)	0.0646 (15)	0.0623 (18)	0.0079 (15)	0.0216 (16)	0.0016 (13)
N2	0.078 (2)	0.0597 (16)	0.0666 (19)	0.0014 (16)	0.0305 (16)	-0.0021 (14)
C1	0.060 (2)	0.0550 (19)	0.070 (2)	0.0004 (17)	0.027 (2)	-0.0008 (17)
C2	0.081 (2)	0.070 (2)	0.062 (2)	0.006 (2)	0.034 (2)	0.0144 (17)
C3	0.076 (3)	0.085 (2)	0.063 (2)	0.003 (2)	0.021 (2)	0.0031 (18)
C4	0.061 (2)	0.084 (2)	0.073 (3)	-0.015 (2)	0.024 (2)	-0.007 (2)
C5	0.069 (2)	0.082 (2)	0.064 (2)	-0.0074 (19)	0.025 (2)	0.0053 (18)
C6	0.062 (2)	0.0521 (18)	0.059 (2)	0.0031 (17)	0.027 (2)	0.0024 (16)
C7	0.079 (2)	0.0551 (19)	0.063 (2)	-0.0008 (19)	0.036 (2)	0.0015 (16)
C8	0.072 (2)	0.077 (2)	0.060 (2)	0.0013 (18)	0.025 (2)	-0.0030 (17)
C9	0.058 (2)	0.0713 (19)	0.061 (2)	-0.0044 (18)	0.0138 (19)	-0.0057 (17)
C10	0.058 (2)	0.082 (2)	0.071 (2)	0.0058 (18)	0.0274 (18)	-0.0003 (17)
C11	0.086 (3)	0.058 (2)	0.062 (2)	0.003 (2)	0.039 (2)	-0.0006 (17)
C12	0.071 (2)	0.0477 (18)	0.053 (2)	0.0033 (18)	0.0293 (19)	-0.0010 (17)
C13	0.080 (3)	0.067 (2)	0.061 (2)	0.012 (2)	0.031 (2)	0.001 (2)
C14	0.112 (3)	0.051 (2)	0.089 (3)	0.004 (2)	0.050 (3)	0.017 (2)
C15	0.105 (3)	0.057 (2)	0.092 (3)	-0.007 (2)	0.048 (3)	0.002 (2)
C16	0.086 (3)	0.057 (2)	0.066 (2)	0.007 (2)	0.032 (2)	0.0060 (19)
C17	0.082 (2)	0.0454 (19)	0.065 (2)	-0.0025 (19)	0.029 (2)	0.0042 (16)
C18	0.085 (2)	0.099 (2)	0.102 (3)	-0.023 (2)	0.017 (2)	-0.015 (2)
C19	0.093 (2)	0.108 (2)	0.078 (2)	0.017 (2)	0.006 (2)	0.012 (2)
C20	0.082 (3)	0.161 (3)	0.113 (3)	-0.043 (2)	0.032 (2)	0.014 (2)
C21	0.114 (3)	0.098 (3)	0.119 (3)	-0.014 (2)	-0.008 (2)	0.028 (2)

Geometric parameters (Å, °)

O1—C1	1.353 (3)	C9—C19	1.543 (3)
O1—H1	0.8074	C9—C18	1.544 (3)
O2—C13	1.348 (3)	C10—H10A	0.9700
O2—H2	0.8251	C10—H10B	0.9700
O3—C4	1.381 (3)	C11—C12	1.454 (3)
O3—C20	1.402 (3)	C11—H11A	0.9300
O4—C16	1.377 (3)	C12—C17	1.391 (3)
O4—C21	1.410 (3)	C12—C13	1.401 (3)
N1—C7	1.269 (3)	C13—C14	1.390 (3)
N1—C8	1.455 (3)	C14—C15	1.353 (4)
N2—C11	1.272 (3)	C14—H14A	0.9300
N2—C10	1.463 (3)	C15—C16	1.385 (3)
C1—C6	1.389 (3)	C15—H15A	0.9300
C1—C2	1.392 (3)	C16—C17	1.380 (3)
C2—C3	1.360 (3)	C17—H17A	0.9300
C2—H2A	0.9300	C18—H18A	0.9600
C3—C4	1.390 (3)	C18—H18B	0.9600
C3—H3A	0.9300	C18—H18C	0.9600
C4—C5	1.371 (3)	C19—H19A	0.9600
C5—C6	1.395 (3)	C19—H19B	0.9600
C5—H5A	0.9300	C19—H19C	0.9600
C6—C7	1.453 (3)	C20—H20A	0.9600
C7—H7A	0.9300	C20—H20B	0.9600
C8—C9	1.528 (3)	C20—H20C	0.9600
C8—H8A	0.9700	C21—H21A	0.9600
C8—H8B	0.9700	C21—H21B	0.9600
C9—C10	1.539 (3)	C21—H21C	0.9600
C1—O1—H1	108.8	N2—C11—C12	121.2 (3)
C13—O2—H2	112.9	N2—C11—H11A	119.4
C4—O3—C20	118.6 (3)	C12—C11—H11A	119.4
C16—O4—C21	117.4 (2)	C17—C12—C13	119.0 (3)
C7—N1—C8	118.4 (2)	C17—C12—C11	118.8 (3)
C11—N2—C10	116.9 (3)	C13—C12—C11	122.2 (3)
O1—C1—C6	121.9 (3)	O2—C13—C14	119.8 (3)
O1—C1—C2	118.5 (3)	O2—C13—C12	121.0 (3)
C6—C1—C2	119.5 (3)	C14—C13—C12	119.2 (3)
C3—C2—C1	120.3 (3)	C15—C14—C13	120.6 (3)
C3—C2—H2A	119.8	C15—C14—H14A	119.7
C1—C2—H2A	119.8	C13—C14—H14A	119.7
C2—C3—C4	120.1 (3)	C14—C15—C16	121.5 (3)
C2—C3—H3A	119.9	C14—C15—H15A	119.3
C4—C3—H3A	119.9	C16—C15—H15A	119.3
C5—C4—O3	125.3 (3)	O4—C16—C17	125.1 (3)
C5—C4—C3	120.7 (3)	O4—C16—C15	116.3 (3)
O3—C4—C3	114.0 (3)	C17—C16—C15	118.6 (3)
C4—C5—C6	119.3 (3)	C16—C17—C12	121.1 (3)

supplementary materials

C4—C5—H5A	120.3	C16—C17—H17A	119.4
C6—C5—H5A	120.3	C12—C17—H17A	119.4
C1—C6—C5	120.0 (3)	C9—C18—H18A	109.5
C1—C6—C7	120.5 (3)	C9—C18—H18B	109.5
C5—C6—C7	119.4 (3)	H18A—C18—H18B	109.5
N1—C7—C6	123.0 (3)	C9—C18—H18C	109.5
N1—C7—H7A	118.5	H18A—C18—H18C	109.5
C6—C7—H7A	118.5	H18B—C18—H18C	109.5
N1—C8—C9	111.7 (2)	C9—C19—H19A	109.5
N1—C8—H8A	109.3	C9—C19—H19B	109.5
C9—C8—H8A	109.3	H19A—C19—H19B	109.5
N1—C8—H8B	109.3	C9—C19—H19C	109.5
C9—C8—H8B	109.3	H19A—C19—H19C	109.5
H8A—C8—H8B	107.9	H19B—C19—H19C	109.5
C8—C9—C10	111.2 (2)	O3—C20—H20A	109.5
C8—C9—C19	108.1 (2)	O3—C20—H20B	109.5
C10—C9—C19	110.3 (2)	H20A—C20—H20B	109.5
C8—C9—C18	110.4 (2)	O3—C20—H20C	109.5
C10—C9—C18	107.6 (2)	H20A—C20—H20C	109.5
C19—C9—C18	109.2 (2)	H20B—C20—H20C	109.5
N2—C10—C9	112.4 (2)	O4—C21—H21A	109.5
N2—C10—H10A	109.1	O4—C21—H21B	109.5
C9—C10—H10A	109.1	H21A—C21—H21B	109.5
N2—C10—H10B	109.1	O4—C21—H21C	109.5
C9—C10—H10B	109.1	H21A—C21—H21C	109.5
H10A—C10—H10B	107.9	H21B—C21—H21C	109.5
O1—C1—C2—C3	−179.5 (2)	C11—N2—C10—C9	−116.9 (3)
C6—C1—C2—C3	−1.5 (4)	C8—C9—C10—N2	54.6 (3)
C1—C2—C3—C4	0.6 (4)	C19—C9—C10—N2	−65.3 (3)
C20—O3—C4—C5	−2.9 (4)	C18—C9—C10—N2	175.7 (2)
C20—O3—C4—C3	176.3 (3)	C10—N2—C11—C12	177.96 (19)
C2—C3—C4—C5	0.2 (4)	N2—C11—C12—C17	−176.1 (3)
C2—C3—C4—O3	−179.0 (3)	N2—C11—C12—C13	3.0 (4)
O3—C4—C5—C6	179.1 (3)	C17—C12—C13—O2	−179.4 (2)
C3—C4—C5—C6	−0.1 (4)	C11—C12—C13—O2	1.5 (4)
O1—C1—C6—C5	179.6 (2)	C17—C12—C13—C14	0.1 (3)
C2—C1—C6—C5	1.7 (4)	C11—C12—C13—C14	−179.0 (2)
O1—C1—C6—C7	1.1 (4)	O2—C13—C14—C15	179.7 (3)
C2—C1—C6—C7	−176.8 (2)	C12—C13—C14—C15	0.2 (4)
C4—C5—C6—C1	−0.9 (4)	C13—C14—C15—C16	−0.4 (5)
C4—C5—C6—C7	177.6 (3)	C21—O4—C16—C17	3.8 (4)
C8—N1—C7—C6	174.6 (2)	C21—O4—C16—C15	−175.7 (2)
C1—C6—C7—N1	−2.5 (4)	C14—C15—C16—O4	179.9 (3)
C5—C6—C7—N1	179.0 (3)	C14—C15—C16—C17	0.3 (4)
C7—N1—C8—C9	−140.7 (2)	O4—C16—C17—C12	−179.6 (2)
N1—C8—C9—C10	48.8 (3)	C15—C16—C17—C12	−0.1 (4)
N1—C8—C9—C19	170.1 (2)	C13—C12—C17—C16	−0.1 (4)
N1—C8—C9—C18	−70.5 (3)	C11—C12—C17—C16	179.0 (2)

Hydrogen-bond geometry (Å, °)

$D-H\cdots A$	$D-H$	$H\cdots A$	$D\cdots A$	$D-H\cdots A$
O1—H1···N1	0.81	1.88	2.593 (3)	147
O2—H2···N2	0.83	1.90	2.604 (3)	143

supplementary materials

Fig. 1

