

Methidathion: *S*-(5-methoxy-2-oxo-2,3-dihydro-1,3,4-thiadiazol-3-yl)methyl *O,O*-dimethyl phosphorodithioate

Hyunjee Kim,^a Yong Woon Shin,^b Ki-Min Park,^{a*} Jineun Kim^a and Tae Ho Kim^{a*}

^aDepartment of Chemistry and Research Institute of Natural Sciences, Gyeongsang National University, Jinju 660-701, Republic of Korea, and ^bTest & Analytical Laboratory, Korea Food & Drug Administration, 123-7 Yongdang-dong, Busan 608-829, Republic of Korea

Correspondence e-mail: kmpark@gnu.ac.kr, thkim@gnu.ac.kr


Received 10 February 2011; accepted 12 February 2011

Key indicators: single-crystal X-ray study; $T = 173\text{ K}$; mean $\sigma(\text{O}-\text{C}) = 0.003\text{ \AA}$; R factor = 0.033; wR factor = 0.089; data-to-parameter ratio = 19.1.

The title compound, $\text{C}_6\text{H}_{11}\text{N}_2\text{O}_4\text{PS}_3$, crystallizes with two independent molecules in the asymmetric unit. The dihedral angles between the thiadiazole ring planes and the PS_2 planes of the phosphorodithioate group are $86.51(5)$ and $56.33(5)^\circ$ in the two molecules. In the crystal, weak intermolecular $\text{S}\cdots\text{S}$ [$3.570(8)\text{ \AA}$] interactions and $\text{C}-\text{H}\cdots\text{O}$ and $\text{C}-\text{H}\cdots\text{N}$ hydrogen bonds contribute to the stabilization of the packing.

Related literature

For the toxicity and insecticidal activity of the title compound, see: Altuntas *et al.* (2002). For related structures, see: Rohrbaugh *et al.* (1976).



Experimental

Crystal data

$\text{C}_6\text{H}_{11}\text{N}_2\text{O}_4\text{PS}_3$	$V = 2528.68(6)\text{ \AA}^3$
$M_r = 302.32$	$Z = 8$
Monoclinic, $P2_1/c$	Mo $K\alpha$ radiation
$a = 12.3944(2)\text{ \AA}$	$\mu = 0.71\text{ mm}^{-1}$
$b = 10.8056(1)\text{ \AA}$	$T = 173\text{ K}$
$c = 19.3631(3)\text{ \AA}$	$0.30 \times 0.27 \times 0.19\text{ mm}$
$\beta = 102.815(1)^\circ$	

Data collection

Bruker APEXII CCD diffractometer	20918 measured reflections
Absorption correction: multi-scan (<i>SADABS</i> ; Sheldrick, 1996)	5513 independent reflections
$T_{\min} = 0.815$, $T_{\max} = 0.877$	4682 reflections with $I > 2\sigma(I)$
	$R_{\text{int}} = 0.024$

Refinement

$R[F^2 > 2\sigma(F^2)] = 0.033$	289 parameters
$wR(F^2) = 0.089$	H-atom parameters constrained
$S = 1.07$	$\Delta\rho_{\max} = 0.98\text{ e \AA}^{-3}$
5513 reflections	$\Delta\rho_{\min} = -0.55\text{ e \AA}^{-3}$

Table 1
Hydrogen-bond geometry (\AA , $^\circ$).

$D-\text{H}\cdots A$	$D-\text{H}$	$\text{H}\cdots A$	$D\cdots A$	$D-\text{H}\cdots A$
C1—H1A \cdots O7 ⁱ	0.98	2.53	3.464 (3)	159
C1—H1B \cdots O5 ⁱⁱ	0.98	2.53	3.484 (3)	164
C2—H2B \cdots O6 ⁱⁱⁱ	0.98	2.59	3.373 (3)	138
C3—H3A \cdots N2 ⁱ	0.99	2.56	3.506 (3)	161
C9—H9B \cdots O3	0.99	2.42	3.246 (2)	140

Symmetry codes: (i) $-x, -y, -z$; (ii) $x - 1, y, z$; (iii) $-x + 1, y + \frac{1}{2}, -z + \frac{1}{2}$.

Data collection: *APEX2* (Bruker, 2006); cell refinement: *SAINT* (Bruker, 2006); data reduction: *SAINT*; program(s) used to solve structure: *SHELXTL* (Sheldrick, 2008); program(s) used to refine structure: *SHELXTL*; molecular graphics: *SHELXTL* and *DIAMOND* (Brandenburg, 1998); software used to prepare material for publication: *SHELXTL*.

This research was supported by the Basic Science Research Program through the National Research Foundation of Korea (NRF) funded by the Ministry of Education, Science and Technology (No. 2010–0009089).

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: JH2268).

References

- Altuntas, I., Delibas, N., Demirci, M., Kilinc, I. & Tamer, N. (2002). *Arch. Toxicol.* **76**, 470–473.
- Brandenburg, K. (1998). *DIAMOND*. Crystal Impact GbR, Bonn, Germany.
- Bruker (2006). *APEX2* and *SAINT*. Bruker AXS Inc., Madison, Wisconsin, USA.
- Rohrbaugh, W. J., Meyers, E. K. & Jacobson, R. A. (1976). *J. Agric. Food Chem.* **24**, 713–717.
- Sheldrick, G. M. (1996). *SADABS*. University of Göttingen, Germany.
- Sheldrick, G. M. (2008). *Acta Cryst. A* **64**, 112–122.

supplementary materials

Acta Cryst. (2011). E67, o653 [doi:10.1107/S1600536811005241]

Methidathion: *S*-(5-methoxy-2-oxo-2,3-dihydro-1,3,4-thiadiazol-3-yl)methyl *O,O*-dimethyl phosphorodithioate

H. Kim, Y. W. Shin, K.-M. Park, J. Kim and T. H. Kim

Comment

Methidathion (systematic name: *S*-2,3-dihydro-5-methoxy-2-oxo-1,3,4-thiadiazol-3-ylmethyl *O,O*-dimethyl phosphorodithioate), is one of the most widely used organophosphate insecticides in agriculture and public health programmes (Altuntas *et al.* 2002). However its crystal structure has not been reported yet.

In the title compound (Scheme 1, Fig. 1), crystallizes with two independent molecules in the asymmetric unit. The dihedral angles between the thiadiazol ring planes and the PS_2 planes of the phosphorodithioate group are $86.51(5)^\circ$ and $56.33(5)^\circ$ in the two molecules. All bond lengths and bond angles are normal and comparable to those observed in similar structures (Rohrbaugh *et al.* 1976).

In the crystal structure, as shown in Fig. 2, weak intermolecular C—H···O, C—H···N hydrogen bonds (Table 1) and S···S interactions with 3.3372 (8) Å are observed (Table 1). These intermolecular interactions may be contribute to the stabilization of the packing.

Experimental

The title compound was purchased from the Dr. Ehrenstorfer GmbH Company. Slow evaporation of a solution in CH_2Cl_2 gave single crystals suitable for X-ray analysis.

Refinement

All H-atoms were positioned geometrically and refined using a riding model with $d(C—H) = 0.99 \text{ \AA}$, $U_{\text{iso}} = 1.2U_{\text{eq}}(C)$ for CH_2 and $d(C—H) = 0.98 \text{ \AA}$, $U_{\text{iso}} = 1.5U_{\text{eq}}(C)$ for CH_3 groups.

Figures




Fig. 1. The molecular structure of the title compound with the atom numbering scheme. Displacement ellipsoids are drawn at the 50% probability level. H atoms are presented as a small spheres of arbitrary radius.

supplementary materials




Fig. 2. Crystal packing of the title compound with intermolecular C—H···O, C—H···N hydrogen bonds and S···S interactions shown as dashed lines. H atoms not involved in intermolecular interactions have been omitted for clarity. [Symmetry codes: (i) $-x, -y, -z$; (ii) $x + 1, y, z$; (iii) $-x + 1, -y, -z$; (iv) $-x + 1, -y + 1, -z$; (v) $-x, -y + 1, -z$.]

S-(5-methoxy-2-oxo-2,3-dihydro-1,3,4-thiadiazol-3-yl)methyl O,O-dimethyl phosphorodithioate

Crystal data

$C_6H_{11}N_2O_4PS_3$	$F(000) = 1248$
$M_r = 302.32$	$D_x = 1.588 \text{ Mg m}^{-3}$
Monoclinic, $P2_1/c$	Mo $K\alpha$ radiation, $\lambda = 0.71073 \text{ \AA}$
Hall symbol: -P 2ybc	Cell parameters from 9930 reflections
$a = 12.3944 (2) \text{ \AA}$	$\theta = 2.5\text{--}28.2^\circ$
$b = 10.8056 (1) \text{ \AA}$	$\mu = 0.71 \text{ mm}^{-1}$
$c = 19.3631 (3) \text{ \AA}$	$T = 173 \text{ K}$
$\beta = 102.815 (1)^\circ$	Block, colourless
$V = 2528.68 (6) \text{ \AA}^3$	$0.30 \times 0.27 \times 0.19 \text{ mm}$
$Z = 8$	

Data collection

Bruker APEXII CCD diffractometer	5513 independent reflections
Radiation source: fine-focus sealed tube graphite	4682 reflections with $I > 2\sigma(I)$
φ and ω scans	$R_{\text{int}} = 0.024$
Absorption correction: multi-scan (<i>SADABS</i> ; Sheldrick, 1996)	$\theta_{\text{max}} = 27.0^\circ, \theta_{\text{min}} = 1.7^\circ$
$T_{\text{min}} = 0.815, T_{\text{max}} = 0.877$	$h = -15 \rightarrow 11$
20918 measured reflections	$k = -11 \rightarrow 13$
	$l = -24 \rightarrow 24$

Refinement

Refinement on F^2	Primary atom site location: structure-invariant direct methods
Least-squares matrix: full	Secondary atom site location: difference Fourier map
$R[F^2 > 2\sigma(F^2)] = 0.033$	Hydrogen site location: inferred from neighbouring sites
$wR(F^2) = 0.089$	H-atom parameters constrained
$S = 1.07$	$w = 1/[\sigma^2(F_o^2) + (0.0424P)^2 + 1.2139P]$
5513 reflections	where $P = (F_o^2 + 2F_c^2)/3$
289 parameters	$(\Delta/\sigma)_{\text{max}} = 0.001$
	$\Delta\rho_{\text{max}} = 0.98 \text{ e \AA}^{-3}$

0 restraints

 $\Delta\rho_{\min} = -0.55 \text{ e } \text{\AA}^{-3}$ *Special details*

Geometry. All e.s.d.'s (except the e.s.d. in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell e.s.d.'s are taken into account individually in the estimation of e.s.d.'s in distances, angles and torsion angles; correlations between e.s.d.'s in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell e.s.d.'s is used for estimating e.s.d.'s involving l.s. planes.

Refinement. Refinement of F^2 against ALL reflections. The weighted R -factor wR and goodness of fit S are based on F^2 , conventional R -factors R are based on F , with F set to zero for negative F^2 . The threshold expression of $F^2 > \sigma(F^2)$ is used only for calculating R -factors(gt) etc. and is not relevant to the choice of reflections for refinement. R -factors based on F^2 are statistically about twice as large as those based on F , and R -factors based on ALL data will be even larger.

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (\AA^2)

	x	y	z	$U_{\text{iso}}^*/U_{\text{eq}}$
P1	0.02509 (4)	0.14885 (4)	0.16800 (3)	0.02754 (12)
P2	0.60901 (4)	-0.02370 (5)	0.25072 (3)	0.02917 (13)
S1	0.01168 (5)	0.17732 (6)	0.26343 (3)	0.04165 (15)
S2	0.15513 (4)	0.02823 (5)	0.16946 (3)	0.03550 (14)
S3	0.16030 (4)	-0.35504 (4)	0.03186 (3)	0.03270 (13)
S4	0.57301 (6)	-0.06480 (7)	0.33906 (3)	0.05122 (17)
S5	0.46376 (4)	0.01769 (5)	0.17901 (3)	0.03259 (13)
S6	0.48755 (4)	0.34626 (5)	-0.00393 (3)	0.03301 (13)
O1	-0.07804 (11)	0.09769 (14)	0.11286 (7)	0.0355 (3)
O2	0.04167 (15)	0.26370 (13)	0.12160 (8)	0.0459 (4)
O3	0.31042 (11)	-0.17257 (13)	0.05006 (8)	0.0363 (3)
O4	-0.05059 (11)	-0.37521 (12)	0.03211 (8)	0.0338 (3)
O5	0.69519 (11)	0.08288 (13)	0.25149 (7)	0.0326 (3)
O6	0.67189 (12)	-0.12180 (13)	0.21387 (8)	0.0347 (3)
O7	0.35232 (12)	0.15068 (14)	-0.01300 (8)	0.0412 (4)
O8	0.67746 (11)	0.41371 (13)	0.07718 (8)	0.0351 (3)
N1	0.13193 (12)	-0.12835 (14)	0.05709 (8)	0.0243 (3)
N2	0.02836 (13)	-0.18037 (14)	0.05340 (9)	0.0263 (3)
N3	0.51734 (13)	0.14990 (15)	0.06949 (9)	0.0283 (4)
N4	0.60999 (13)	0.22068 (15)	0.09789 (8)	0.0269 (3)
C1	-0.14361 (19)	-0.0019 (2)	0.13290 (14)	0.0459 (6)
H1A	-0.2042	-0.0227	0.0927	0.069*
H1B	-0.1742	0.0246	0.1730	0.069*
H1C	-0.0967	-0.0748	0.1465	0.069*
C2	0.1050 (3)	0.3680 (3)	0.14990 (15)	0.0700 (9)
H2A	0.1047	0.4285	0.1122	0.105*
H2B	0.1813	0.3423	0.1703	0.105*
H2C	0.0731	0.4055	0.1869	0.105*
C3	0.14752 (16)	0.00052 (17)	0.07483 (10)	0.0273 (4)
H3A	0.0853	0.0487	0.0464	0.033*
H3B	0.2167	0.0298	0.0626	0.033*

supplementary materials

C4	0.21602 (15)	-0.20269 (17)	0.04761 (10)	0.0259 (4)
C5	0.03364 (15)	-0.29610 (17)	0.04052 (10)	0.0263 (4)
C6	-0.15623 (17)	-0.31980 (19)	0.03375 (14)	0.0383 (5)
H6A	-0.2133	-0.3842	0.0272	0.057*
H6B	-0.1755	-0.2586	-0.0043	0.057*
H6C	-0.1515	-0.2791	0.0795	0.057*
C7	0.6764 (2)	0.2031 (2)	0.27938 (12)	0.0426 (5)
H7A	0.7376	0.2585	0.2760	0.064*
H7B	0.6721	0.1949	0.3291	0.064*
H7C	0.6069	0.2375	0.2520	0.064*
C8	0.6314 (2)	-0.2470 (2)	0.20117 (15)	0.0536 (7)
H8A	0.6811	-0.2940	0.1780	0.080*
H8B	0.5570	-0.2456	0.1705	0.080*
H8C	0.6287	-0.2864	0.2464	0.080*
C9	0.50949 (17)	0.02671 (18)	0.09544 (10)	0.0291 (4)
H9A	0.5829	-0.0133	0.1020	0.035*
H9B	0.4572	-0.0210	0.0591	0.035*
C10	0.43898 (16)	0.19916 (19)	0.01633 (11)	0.0305 (4)
C11	0.60347 (16)	0.32287 (18)	0.06415 (10)	0.0275 (4)
C12	0.7657 (2)	0.3958 (2)	0.13932 (13)	0.0494 (6)
H12A	0.8163	0.4666	0.1448	0.074*
H12B	0.8063	0.3199	0.1339	0.074*
H12C	0.7342	0.3887	0.1813	0.074*

Atomic displacement parameters (\AA^2)

	U^{11}	U^{22}	U^{33}	U^{12}	U^{13}	U^{23}
P1	0.0317 (3)	0.0231 (2)	0.0290 (3)	0.00059 (19)	0.0094 (2)	-0.0009 (2)
P2	0.0291 (3)	0.0324 (3)	0.0282 (3)	-0.0032 (2)	0.0112 (2)	0.0027 (2)
S1	0.0511 (4)	0.0445 (3)	0.0318 (3)	-0.0008 (3)	0.0143 (3)	-0.0075 (2)
S2	0.0322 (3)	0.0394 (3)	0.0307 (3)	0.0102 (2)	-0.0021 (2)	-0.0081 (2)
S3	0.0272 (3)	0.0230 (2)	0.0480 (3)	0.00344 (19)	0.0086 (2)	-0.0059 (2)
S4	0.0610 (4)	0.0630 (4)	0.0361 (3)	-0.0113 (3)	0.0247 (3)	0.0075 (3)
S5	0.0229 (2)	0.0389 (3)	0.0378 (3)	-0.0022 (2)	0.0106 (2)	-0.0046 (2)
S6	0.0327 (3)	0.0342 (3)	0.0299 (3)	0.0100 (2)	0.0021 (2)	0.0020 (2)
O1	0.0268 (7)	0.0473 (9)	0.0317 (8)	0.0021 (6)	0.0051 (6)	0.0004 (7)
O2	0.0710 (11)	0.0257 (7)	0.0442 (9)	-0.0051 (7)	0.0200 (8)	0.0005 (7)
O3	0.0253 (7)	0.0347 (8)	0.0505 (9)	-0.0015 (6)	0.0119 (7)	-0.0092 (7)
O4	0.0269 (7)	0.0220 (6)	0.0529 (9)	-0.0008 (5)	0.0098 (7)	-0.0015 (6)
O5	0.0290 (7)	0.0361 (8)	0.0335 (8)	-0.0074 (6)	0.0089 (6)	-0.0008 (6)
O6	0.0360 (8)	0.0310 (7)	0.0394 (8)	0.0035 (6)	0.0131 (7)	0.0057 (6)
O7	0.0283 (8)	0.0419 (9)	0.0463 (9)	0.0062 (7)	-0.0068 (7)	-0.0131 (7)
O8	0.0297 (7)	0.0351 (8)	0.0396 (8)	-0.0035 (6)	0.0054 (6)	0.0071 (7)
N1	0.0219 (8)	0.0211 (7)	0.0304 (8)	-0.0001 (6)	0.0067 (7)	-0.0015 (6)
N2	0.0240 (8)	0.0231 (8)	0.0321 (9)	-0.0004 (6)	0.0071 (7)	-0.0003 (7)
N3	0.0264 (8)	0.0274 (8)	0.0283 (9)	0.0007 (6)	0.0002 (7)	-0.0022 (7)
N4	0.0238 (8)	0.0301 (8)	0.0262 (8)	0.0000 (6)	0.0041 (7)	-0.0002 (7)
C1	0.0336 (12)	0.0538 (14)	0.0530 (15)	-0.0132 (10)	0.0157 (11)	-0.0132 (12)

C2	0.098 (2)	0.0519 (16)	0.0517 (16)	-0.0435 (16)	-0.0026 (16)	0.0031 (13)
C3	0.0284 (10)	0.0230 (9)	0.0306 (10)	-0.0004 (8)	0.0069 (8)	-0.0014 (8)
C4	0.0275 (10)	0.0241 (9)	0.0261 (9)	0.0023 (7)	0.0057 (8)	-0.0020 (8)
C5	0.0256 (10)	0.0239 (9)	0.0292 (10)	0.0028 (7)	0.0056 (8)	0.0007 (8)
C6	0.0264 (11)	0.0295 (10)	0.0604 (15)	-0.0019 (8)	0.0127 (10)	-0.0041 (10)
C7	0.0519 (14)	0.0384 (12)	0.0360 (12)	-0.0105 (10)	0.0069 (11)	-0.0075 (10)
C8	0.0713 (18)	0.0287 (12)	0.0661 (17)	-0.0002 (11)	0.0269 (15)	0.0012 (11)
C9	0.0292 (10)	0.0279 (10)	0.0300 (10)	0.0007 (8)	0.0065 (8)	-0.0053 (8)
C10	0.0274 (10)	0.0327 (10)	0.0299 (10)	0.0088 (8)	0.0031 (9)	-0.0074 (8)
C11	0.0240 (9)	0.0324 (10)	0.0263 (10)	0.0050 (8)	0.0063 (8)	0.0003 (8)
C12	0.0428 (14)	0.0520 (14)	0.0460 (14)	-0.0172 (11)	-0.0056 (11)	0.0082 (12)

Geometric parameters (Å, °)

P1—O2	1.5720 (15)	N2—C5	1.280 (2)
P1—O1	1.5730 (15)	N3—C10	1.357 (3)
P1—S1	1.9166 (7)	N3—N4	1.388 (2)
P1—S2	2.0681 (7)	N3—C9	1.434 (2)
P2—O5	1.5685 (14)	N4—C11	1.277 (2)
P2—O6	1.5770 (15)	C1—H1A	0.9800
P2—S4	1.9135 (7)	C1—H1B	0.9800
P2—S5	2.0640 (8)	C1—H1C	0.9800
S2—C3	1.838 (2)	C2—H2A	0.9800
S3—C5	1.7363 (19)	C2—H2B	0.9800
S3—C4	1.7852 (19)	C2—H2C	0.9800
S5—C9	1.832 (2)	C3—H3A	0.9900
S6—C11	1.739 (2)	C3—H3B	0.9900
S6—C10	1.774 (2)	C6—H6A	0.9800
O1—C1	1.452 (3)	C6—H6B	0.9800
O2—C2	1.413 (3)	C6—H6C	0.9800
O3—C4	1.205 (2)	C7—H7A	0.9800
O4—C5	1.331 (2)	C7—H7B	0.9800
O4—C6	1.446 (2)	C7—H7C	0.9800
O5—C7	1.446 (3)	C8—H8A	0.9800
O6—C8	1.445 (3)	C8—H8B	0.9800
O7—C10	1.217 (2)	C8—H8C	0.9800
O8—C11	1.329 (2)	C9—H9A	0.9900
O8—C12	1.448 (3)	C9—H9B	0.9900
N1—C4	1.360 (2)	C12—H12A	0.9800
N1—N2	1.389 (2)	C12—H12B	0.9800
N1—C3	1.437 (2)	C12—H12C	0.9800
O2—P1—O1	94.54 (9)	N1—C3—H3B	109.3
O2—P1—S1	118.33 (6)	S2—C3—H3B	109.3
O1—P1—S1	118.66 (6)	H3A—C3—H3B	107.9
O2—P1—S2	107.77 (7)	O3—C4—N1	126.98 (18)
O1—P1—S2	107.54 (6)	O3—C4—S3	126.24 (15)
S1—P1—S2	108.78 (3)	N1—C4—S3	106.78 (13)
O5—P2—O6	95.40 (8)	N2—C5—O4	125.36 (17)
O5—P2—S4	117.08 (6)	N2—C5—S3	117.62 (15)

supplementary materials

O6—P2—S4	119.02 (6)	O4—C5—S3	117.00 (13)
O5—P2—S5	109.75 (6)	O4—C6—H6A	109.5
O6—P2—S5	106.63 (6)	O4—C6—H6B	109.5
S4—P2—S5	108.07 (3)	H6A—C6—H6B	109.5
C3—S2—P1	102.80 (7)	O4—C6—H6C	109.5
C5—S3—C4	88.23 (9)	H6A—C6—H6C	109.5
C9—S5—P2	102.19 (7)	H6B—C6—H6C	109.5
C11—S6—C10	88.05 (9)	O5—C7—H7A	109.5
C1—O1—P1	119.85 (14)	O5—C7—H7B	109.5
C2—O2—P1	122.67 (16)	H7A—C7—H7B	109.5
C5—O4—C6	114.88 (15)	O5—C7—H7C	109.5
C7—O5—P2	119.87 (13)	H7A—C7—H7C	109.5
C8—O6—P2	121.24 (14)	H7B—C7—H7C	109.5
C11—O8—C12	114.92 (16)	O6—C8—H8A	109.5
C4—N1—N2	118.77 (15)	O6—C8—H8B	109.5
C4—N1—C3	122.56 (16)	H8A—C8—H8B	109.5
N2—N1—C3	118.57 (14)	O6—C8—H8C	109.5
C5—N2—N1	108.60 (15)	H8A—C8—H8C	109.5
C10—N3—N4	118.32 (16)	H8B—C8—H8C	109.5
C10—N3—C9	122.48 (17)	N3—C9—S5	114.67 (13)
N4—N3—C9	119.16 (16)	N3—C9—H9A	108.6
C11—N4—N3	108.72 (16)	S5—C9—H9A	108.6
O1—C1—H1A	109.5	N3—C9—H9B	108.6
O1—C1—H1B	109.5	S5—C9—H9B	108.6
H1A—C1—H1B	109.5	H9A—C9—H9B	107.6
O1—C1—H1C	109.5	O7—C10—N3	126.5 (2)
H1A—C1—H1C	109.5	O7—C10—S6	126.18 (16)
H1B—C1—H1C	109.5	N3—C10—S6	107.35 (14)
O2—C2—H2A	109.5	N4—C11—O8	125.39 (18)
O2—C2—H2B	109.5	N4—C11—S6	117.48 (15)
H2A—C2—H2B	109.5	O8—C11—S6	117.12 (14)
O2—C2—H2C	109.5	O8—C12—H12A	109.5
H2A—C2—H2C	109.5	O8—C12—H12B	109.5
H2B—C2—H2C	109.5	H12A—C12—H12B	109.5
N1—C3—S2	111.76 (13)	O8—C12—H12C	109.5
N1—C3—H3A	109.3	H12A—C12—H12C	109.5
S2—C3—H3A	109.3	H12B—C12—H12C	109.5
O2—P1—S2—C3	-55.63 (9)	C3—N1—C4—O3	-2.7 (3)
O1—P1—S2—C3	45.23 (9)	N2—N1—C4—S3	0.3 (2)
S1—P1—S2—C3	174.93 (7)	C3—N1—C4—S3	176.58 (14)
O5—P2—S5—C9	61.04 (9)	C5—S3—C4—O3	179.15 (19)
O6—P2—S5—C9	-41.17 (9)	C5—S3—C4—N1	-0.10 (14)
S4—P2—S5—C9	-170.21 (7)	N1—N2—C5—O4	-178.69 (17)
O2—P1—O1—C1	-169.51 (15)	N1—N2—C5—S3	0.2 (2)
S1—P1—O1—C1	-43.64 (17)	C6—O4—C5—N2	4.0 (3)
S2—P1—O1—C1	80.24 (15)	C6—O4—C5—S3	-174.91 (15)
O1—P1—O2—C2	162.2 (2)	C4—S3—C5—N2	-0.09 (17)
S1—P1—O2—C2	36.1 (2)	C4—S3—C5—O4	178.94 (16)
S2—P1—O2—C2	-87.7 (2)	C10—N3—C9—S5	-101.82 (19)

O6—P2—O5—C7	178.26 (15)	N4—N3—C9—S5	80.34 (19)
S4—P2—O5—C7	−55.19 (16)	P2—S5—C9—N3	−105.14 (14)
S5—P2—O5—C7	68.42 (15)	N4—N3—C10—O7	−178.22 (18)
O5—P2—O6—C8	177.37 (18)	C9—N3—C10—O7	3.9 (3)
S4—P2—O6—C8	52.25 (19)	N4—N3—C10—S6	3.0 (2)
S5—P2—O6—C8	−70.14 (18)	C9—N3—C10—S6	−174.84 (14)
C4—N1—N2—C5	−0.3 (2)	C11—S6—C10—O7	179.08 (19)
C3—N1—N2—C5	−176.80 (17)	C11—S6—C10—N3	−2.15 (14)
C10—N3—N4—C11	−2.3 (2)	N3—N4—C11—O8	179.59 (17)
C9—N3—N4—C11	175.65 (16)	N3—N4—C11—S6	0.3 (2)
C4—N1—C3—S2	−101.78 (18)	C12—O8—C11—N4	−6.0 (3)
N2—N1—C3—S2	74.52 (18)	C12—O8—C11—S6	173.21 (16)
P1—S2—C3—N1	−120.69 (12)	C10—S6—C11—N4	1.10 (16)
N2—N1—C4—O3	−178.97 (19)	C10—S6—C11—O8	−178.22 (15)

Hydrogen-bond geometry (Å, °)

<i>D</i> —H··· <i>A</i>	<i>D</i> —H	H··· <i>A</i>	<i>D</i> ··· <i>A</i>	<i>D</i> —H··· <i>A</i>
C1—H1A···O7 ⁱ	0.98	2.53	3.464 (3)	159
C1—H1B···O5 ⁱⁱ	0.98	2.53	3.484 (3)	164
C2—H2B···O6 ⁱⁱⁱ	0.98	2.59	3.373 (3)	138
C3—H3A···N2 ⁱ	0.99	2.56	3.506 (3)	161
C6—H6B···O7 ⁱ	0.98	2.57	2.996 (2)	106
C9—H9B···O3	0.99	2.42	3.246 (2)	140

Symmetry codes: (i) $-x, -y, -z$; (ii) $x-1, y, z$; (iii) $-x+1, y+1/2, -z+1/2$.

supplementary materials

Fig. 1




Fig. 2

