

**catena-Poly[bis(propane-1,3-diaminium)
[[aqua(sulfato- κ O)bis(sulfato- κ^2 O,O')-
cerate(IV)]- μ -sulfato- κ^3 O,O':O']]
dihydrate]**

Ali Farooq Meer,^a Saeed Ahmad,^a Shahzad Sharif,^b
Islam Ullah Khan^b and Seik Weng Ng^{c*}

^aDepartment of Chemistry, University of Engineering and Technology, Lahore 54890, Pakistan, ^bMaterials Chemistry Laboratory, Department of Chemistry, Government College University, 54000 Lahore, Pakistan, and ^cDepartment of Chemistry, University of Malaya, 50603 Kuala Lumpur, Malaysia
Correspondence e-mail: seikweng@um.edu.my


Received 11 February 2011; accepted 4 March 2011

Key indicators: single-crystal X-ray study; $T = 295$ K; mean $\sigma(\text{C}-\text{C}) = 0.006$ Å;
R factor = 0.028; wR factor = 0.071; data-to-parameter ratio = 17.3.

The Ce^{IV} atom in the title salt, $\{(\text{H}_3\text{NCH}_2\text{CH}_2\text{CH}_2\text{NH}_3)_2[\text{Ce}(\text{SO}_4)_4(\text{H}_2\text{O})]\cdot 2\text{H}_2\text{O}\}_n$, exists in a monocapped square-anti-prismatic coordination geometry. The water-coordinated metal atom is bonded to four sulfate ions; one of them is monodentate and two function in a chelating mode. The fourth is also chelating but it uses one of the other two O atoms to bind to an adjacent metal atom, generating a polyanionic chain. The cations are linked to the polyanionic chain as well as to the uncoordinated water molecules, resulting in an O–H···O and N–H···O hydrogen-bonded three-dimensional network.

Related literature

For $(\text{C}_2\text{H}_{10}\text{N}_2)_5[\text{Ce}_2(\text{SO}_4)_9]\cdot 3\text{H}_2\text{O}$, see: Jabeen *et al.* (2010).



Experimental

Crystal data

$(\text{C}_3\text{H}_{12}\text{N}_2)_2[\text{Ce}(\text{SO}_4)_4(\text{H}_2\text{O})]\cdot 2\text{H}_2\text{O}$	$c = 12.8688 (2)$ Å
$M_r = 730.70$	$\beta = 99.535 (1)^\circ$
Monoclinic, $P2_1/c$	$V = 2321.71 (6)$ Å ³
$a = 8.9459 (1)$ Å	$Z = 4$
$b = 20.4497 (3)$ Å	Mo $K\alpha$ radiation

$\mu = 2.41$ mm^{−1}
 $T = 295$ K

$0.30 \times 0.15 \times 0.10$ mm

Data collection

Bruker Kappa APEXII
diffractometer
Absorption correction: multi-scan
(SADABS; Sheldrick, 1996)
 $T_{\min} = 0.531$, $T_{\max} = 0.794$

21238 measured reflections
5296 independent reflections
4663 reflections with $I > 2\sigma(I)$
 $R_{\text{int}} = 0.024$

Refinement

$R[F^2 > 2\sigma(F^2)] = 0.028$
 $wR(F^2) = 0.071$
 $S = 1.14$
5296 reflections

307 parameters
H-atom parameters constrained
 $\Delta\rho_{\max} = 1.24$ e Å^{−3}
 $\Delta\rho_{\min} = -1.41$ e Å^{−3}

Table 1
Hydrogen-bond geometry (Å, °).

D–H···A	D–H	H···A	D···A	D–H···A
O1w–H1w2···O12 ⁱ	0.84	2.05	2.854 (4)	162
O2w–H2w1···O11 ⁱ	0.84	2.24	2.935 (5)	140
O2w–H2w2···O15 ⁱⁱ	0.84	2.08	2.902 (5)	167
O3w–H3w1···O3 ⁱ	0.84	1.90	2.729 (5)	169
O3w–H3w2···O2w	0.85	1.98	2.802 (5)	163
N1–H11···O9 ⁱ	0.86	2.04	2.881 (4)	167
N1–H12···O3w	0.86	1.95	2.806 (5)	171
N1–H13···O13 ⁱⁱⁱ	0.86	2.19	3.036 (4)	166
N2–H21···O16 ^{iv}	0.86	2.09	2.904 (5)	157
N2–H22···O16 ⁱⁱ	0.86	2.25	2.978 (5)	142
N2–H23···O2	0.86	2.49	3.180 (6)	137
N2–H23···O5	0.86	2.44	2.989 (4)	122
N3–H31···O6 ^v	0.86	2.04	2.866 (4)	162
N3–H32···O4	0.86	2.37	2.882 (6)	119
N3–H32···O2w ^{vi}	0.86	2.37	3.085 (6)	141
N3–H33···O1	0.86	2.33	3.004 (5)	135
N3–H33···O10	0.86	2.43	3.191 (6)	147
N4–H41···O6 ^{vii}	0.86	2.34	2.986 (4)	132
N4–H41···O15 ^{viii}	0.86	2.25	2.951 (5)	138
N4–H43···O4 ^{viii}	0.86	2.00	2.851 (5)	173

Symmetry codes: (i) $x - 1, y, z$; (ii) $-x + 1, y - \frac{1}{2}, -z + \frac{1}{2}$; (iii) $x - 1, -y + \frac{1}{2}, z - \frac{1}{2}$; (iv) $x, -y + \frac{1}{2}, z - \frac{1}{2}$; (v) $x, -y + \frac{1}{2}, z + \frac{1}{2}$; (vi) $-x + 1, -y, -z + 1$; (vii) $x + 1, -y + \frac{1}{2}, z + \frac{1}{2}$; (viii) $-x + 2, -y, -z + 1$.

Data collection: APEX2 (Bruker, 2009); cell refinement: SAINT (Bruker, 2009); data reduction: SAINT; program(s) used to solve structure: SHELLS97 (Sheldrick, 2008); program(s) used to refine structure: SHELLXL97 (Sheldrick, 2008); molecular graphics: X-SEED (Barbour, 2001); software used to prepare material for publication: publCIF (Westrip, 2010).

We thank the Higher Education Commission of Pakistan, GC University and the University of Malaya for supporting this study.

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: BT5471).

References

- Barbour, L. J. (2001). *J. Supramol. Chem.* **1**, 189–191.
- Bruker (2009). APEX2 and SAINT. Bruker AXS Inc., Madison, Wisconsin, USA.
- Jabeen, N., Ahmad, S., Meer, A. F., Khan, I. U. & Ng, S. W. (2010). *Acta Cryst. E66*, m797–m798.
- Sheldrick, G. M. (1996). SADABS. University of Göttingen, Germany.
- Sheldrick, G. M. (2008). *Acta Cryst. A64*, 112–122.
- Westrip, S. P. (2010). *J. Appl. Cryst.* **43**, 920–925.

supplementary materials

Acta Cryst. (2011). E67, m413 [doi:10.1107/S1600536811008324]

catena-Poly[bis(propane-1,3-diaminium) [[aqua(sulfato- κO)bis(sulfato- $\kappa^2 O,O'$)cerate(IV)]- μ -sulfato- $\kappa^3 O,O':O''$] dihydrate]

A. F. Meer, S. Ahmad, S. Sharif, I. U. Khan and S. W. Ng

Comment

A previous study reports the isolation of pentakis(ethylenediammonium) μ -sulfato-bis[trisulfatocerate(IV) trihydrate, which was synthesized by the reaction of 1,2-diaminoethane with cerium(IV) sulfate (Jabeen *et al.*, 2010). The two independent Ce atoms exist in a nine-coordinate geometry this is best described as a tricapped trigonal prism. Shortening the cationic chain results in the formation of a polyanion. The compound obtained with 1,3-diaminopropane in place of 1,2-diaminoethane is $2(\text{H}_3\text{NCH}_2\text{CH}_2\text{CH}_2\text{NH}_3)[\text{Ce}(\text{H}_2\text{O})(\text{SO}_4)_4] \cdot 2\text{H}_2\text{O}$ (Scheme I, Fig. 1). One of the sulfate ions behaving in a bridging mode to link adjacent cerate ions into a chain. The cations are linked to the polyanionic chain as well as to the lattice water molecules to result in a hydrogen-bonded three-dimensional network. The metal atom shows monocapped square-antiprismatic geometry (Fig. 2).

Experimental

1,3-Diaminopropane (0.148 g, 2 mmol) was placed in a 1 N sulfuric acid solution (5 ml) of cerium(IV) sulfate tetrahydrate (0.202 g, 0.5 mmol). The yellow solution was filtered and then set aside for the growth of crystals.

Refinement

Carbon- and nitrogen-bound H-atoms were placed in calculated positions (C–H 0.93 to 0.97 Å; N–H 0.86 Å) and were included in the refinement in the riding model approximation, with $U(\text{H})$ set to 1.2–1.5 $U(\text{C},\text{N})$. The H-atoms of the water molecules were placed in chemically sensible positions on the basis of hydrogen bonding interactions (O–H 0.84 Å) and their temperature factors were similarly tied. The final difference Fourier map had a peak at 0.85 Å from Ce1 and a hole at 1.18 Å from O10.

The (1 0 0), (1 1 0) and (0 2 0) reflections were omitted as they were affected by the beam stop. The reflections (-1 3 10), (-1 7 8), (0 7 6), (0 1 6), (-1 8 12) and (-1 2 12) were omitted because of bad disagreement between the calculated and observed intensities.

Figures




Fig. 1. Thermal ellipsoid plot (Barbour, 2001) of a portion of polymeric $(\text{H}_3\text{NCH}_2\text{CH}_2\text{CH}_2\text{NH}_3)[\text{Ce}(\text{H}_2\text{O})(\text{SO}_4)_4] \cdot 2\text{H}_2\text{O}$ at the 50% probability level; hydrogen atoms are drawn as spheres of arbitrary radius.

supplementary materials




Fig. 2. Monocapped square-antiprismatic geometry of Ce(IV).

catena-Poly[bis(propane-1,3-diaminium) [[aqua(sulfato- κ O)bis(sulfato- κ^2 O,O')cerate(IV)]- μ -sulfato- κ^3 O,O':O''] dihydrate]

Crystal data

$(C_3H_{12}N_2)_2[Ce(SO_4)_4(H_2O)] \cdot 2H_2O$	$F(000) = 1472$
$M_r = 730.70$	$D_x = 2.090 \text{ Mg m}^{-3}$
Monoclinic, $P2_1/c$	Mo $K\alpha$ radiation, $\lambda = 0.71073 \text{ \AA}$
Hall symbol: -P 2ybc	Cell parameters from 9960 reflections
$a = 8.9459 (1) \text{ \AA}$	$\theta = 2.8\text{--}28.3^\circ$
$b = 20.4497 (3) \text{ \AA}$	$\mu = 2.41 \text{ mm}^{-1}$
$c = 12.8688 (2) \text{ \AA}$	$T = 295 \text{ K}$
$\beta = 99.535 (1)^\circ$	Prism, yellow
$V = 2321.71 (6) \text{ \AA}^3$	$0.30 \times 0.15 \times 0.10 \text{ mm}$
$Z = 4$	

Data collection

Bruker Kappa APEXII diffractometer	5296 independent reflections
Radiation source: fine-focus sealed tube graphite	4663 reflections with $I > 2\sigma(I)$
φ and ω scans	$R_{\text{int}} = 0.024$
Absorption correction: multi-scan (<i>SADABS</i> ; Sheldrick, 1996)	$\theta_{\text{max}} = 27.5^\circ, \theta_{\text{min}} = 1.9^\circ$
$T_{\text{min}} = 0.531, T_{\text{max}} = 0.794$	$h = -11 \rightarrow 11$
21238 measured reflections	$k = -26 \rightarrow 20$
	$l = -16 \rightarrow 16$

Refinement

Refinement on F^2	Primary atom site location: structure-invariant direct methods
Least-squares matrix: full	Secondary atom site location: difference Fourier map
$R[F^2 > 2\sigma(F^2)] = 0.028$	Hydrogen site location: inferred from neighbouring sites
$wR(F^2) = 0.071$	H-atom parameters constrained
$S = 1.14$	$w = 1/[\sigma^2(F_o^2) + (0.0167P)^2 + 7.615P]$
5296 reflections	where $P = (F_o^2 + 2F_c^2)/3$
	$(\Delta/\sigma)_{\text{max}} = 0.001$

307 parameters $\Delta\rho_{\max} = 1.24 \text{ e Å}^{-3}$
 0 restraints $\Delta\rho_{\min} = -1.41 \text{ e Å}^{-3}$

Special details

Geometry. All e.s.d.'s (except the e.s.d. in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell e.s.d.'s are taken into account individually in the estimation of e.s.d.'s in distances, angles and torsion angles; correlations between e.s.d.'s in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell e.s.d.'s is used for estimating e.s.d.'s involving l.s. planes.

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (\AA^2)

	<i>x</i>	<i>y</i>	<i>z</i>	$U_{\text{iso}}^*/U_{\text{eq}}$
Ce1	0.682986 (19)	0.252538 (8)	0.326048 (13)	0.01407 (6)
S1	0.58776 (10)	0.08286 (4)	0.35192 (8)	0.02668 (19)
S2	0.57080 (9)	0.24695 (4)	0.08631 (6)	0.01880 (16)
S3	1.02446 (9)	0.22812 (4)	0.34548 (7)	0.02170 (17)
S4	0.66130 (10)	0.39960 (4)	0.35617 (7)	0.02449 (18)
O1	0.6789 (3)	0.14558 (12)	0.3672 (2)	0.0248 (5)
O2	0.4287 (3)	0.10079 (15)	0.3213 (4)	0.0588 (11)
O3	0.6376 (4)	0.04469 (15)	0.2697 (3)	0.0429 (7)
O4	0.6144 (4)	0.04863 (16)	0.4518 (3)	0.0530 (9)
O5	0.6001 (3)	0.19145 (12)	0.16128 (19)	0.0266 (5)
O6	0.6232 (3)	0.30497 (11)	0.15445 (18)	0.0227 (5)
O7	0.4153 (3)	0.25303 (17)	0.0394 (2)	0.0439 (8)
O8	0.6718 (3)	0.24052 (12)	0.00700 (19)	0.0246 (5)
O9	0.9026 (3)	0.23795 (12)	0.24892 (19)	0.0236 (5)
O10	0.9303 (3)	0.23354 (13)	0.4321 (2)	0.0253 (5)
O11	1.0895 (3)	0.16402 (14)	0.3425 (2)	0.0384 (7)
O12	1.1351 (3)	0.27980 (14)	0.3520 (2)	0.0323 (6)
O13	0.8002 (3)	0.35764 (12)	0.3595 (2)	0.0236 (5)
O14	0.5368 (3)	0.35021 (12)	0.3413 (2)	0.0231 (5)
O15	0.6465 (3)	0.44422 (15)	0.2677 (3)	0.0486 (8)
O16	0.6665 (3)	0.43396 (15)	0.4552 (3)	0.0436 (8)
O1w	0.4185 (3)	0.23056 (12)	0.3117 (2)	0.0245 (5)
H1w1	0.4039	0.1903	0.3164	0.037*
H1w2	0.3478	0.2512	0.3315	0.037*
O2w	0.1571 (4)	0.02367 (19)	0.3392 (3)	0.0621 (10)
H2w1	0.1834	0.0622	0.3559	0.093*
H2w2	0.2243	0.0057	0.3107	0.093*
O3w	-0.1048 (4)	0.02558 (18)	0.1859 (3)	0.0572 (9)
H3w1	-0.1861	0.0260	0.2112	0.086*
H3w2	-0.0325	0.0329	0.2360	0.086*
N1	-0.0516 (4)	0.14453 (17)	0.0895 (3)	0.0332 (7)
H11	-0.0698	0.1765	0.1290	0.050*
H12	-0.0780	0.1084	0.1155	0.050*
H13	-0.1022	0.1498	0.0272	0.050*
N2	0.4215 (4)	0.07383 (16)	0.0772 (3)	0.0334 (7)

supplementary materials

H21	0.5087	0.0757	0.0575	0.050*
H22	0.4077	0.0352	0.1003	0.050*
H23	0.4179	0.1019	0.1264	0.050*
N3	0.8546 (4)	0.1186 (2)	0.5827 (3)	0.0539 (11)
H31	0.7883	0.1354	0.6161	0.081*
H32	0.8346	0.0778	0.5713	0.081*
H33	0.8522	0.1384	0.5236	0.081*
N4	1.3989 (4)	0.08791 (18)	0.5938 (3)	0.0445 (9)
H41	1.4856	0.0974	0.6304	0.067*
H42	1.3901	0.1068	0.5334	0.067*
H43	1.3920	0.0463	0.5851	0.067*
C1	0.1122 (4)	0.14262 (19)	0.0843 (3)	0.0308 (8)
H1A	0.1699	0.1367	0.1544	0.037*
H1B	0.1430	0.1836	0.0563	0.037*
C2	0.1438 (4)	0.0867 (2)	0.0140 (3)	0.0322 (8)
H2A	0.1307	0.0457	0.0493	0.039*
H2B	0.0703	0.0879	-0.0505	0.039*
C3	0.3013 (4)	0.0886 (2)	-0.0138 (3)	0.0348 (9)
H3A	0.3073	0.0572	-0.0694	0.042*
H3B	0.3193	0.1317	-0.0407	0.042*
C4	1.0041 (5)	0.1253 (2)	0.6454 (4)	0.0438 (11)
H4A	1.0057	0.1038	0.7128	0.053*
H4B	1.0260	0.1713	0.6587	0.053*
C5	1.1240 (5)	0.0961 (2)	0.5913 (4)	0.0379 (9)
H5A	1.1098	0.0491	0.5862	0.045*
H5B	1.1141	0.1134	0.5204	0.045*
C6	1.2785 (5)	0.1104 (3)	0.6490 (4)	0.0488 (12)
H6A	1.2885	0.1573	0.6600	0.059*
H6B	1.2907	0.0897	0.7177	0.059*

Atomic displacement parameters (\AA^2)

	U^{11}	U^{22}	U^{33}	U^{12}	U^{13}	U^{23}
Ce1	0.01607 (9)	0.01389 (10)	0.01262 (9)	-0.00095 (6)	0.00347 (6)	-0.00052 (6)
S1	0.0263 (4)	0.0165 (4)	0.0378 (5)	-0.0020 (3)	0.0068 (4)	0.0034 (4)
S2	0.0183 (4)	0.0248 (4)	0.0135 (3)	-0.0014 (3)	0.0034 (3)	-0.0013 (3)
S3	0.0178 (4)	0.0231 (4)	0.0246 (4)	0.0031 (3)	0.0048 (3)	-0.0003 (3)
S4	0.0238 (4)	0.0151 (4)	0.0346 (5)	0.0002 (3)	0.0049 (3)	-0.0016 (3)
O1	0.0269 (13)	0.0162 (12)	0.0301 (13)	-0.0039 (10)	0.0017 (10)	0.0035 (10)
O2	0.0221 (14)	0.0242 (16)	0.129 (4)	-0.0024 (12)	0.0085 (17)	-0.0031 (18)
O3	0.0468 (17)	0.0337 (16)	0.0485 (19)	-0.0058 (13)	0.0090 (14)	-0.0141 (14)
O4	0.084 (3)	0.0323 (17)	0.0453 (19)	-0.0127 (16)	0.0176 (18)	0.0116 (14)
O5	0.0414 (15)	0.0198 (12)	0.0197 (12)	-0.0089 (11)	0.0079 (10)	-0.0012 (10)
O6	0.0318 (13)	0.0166 (12)	0.0197 (12)	0.0016 (10)	0.0045 (9)	-0.0005 (9)
O7	0.0197 (13)	0.082 (3)	0.0286 (15)	0.0005 (14)	0.0004 (11)	-0.0020 (15)
O8	0.0264 (12)	0.0317 (14)	0.0173 (12)	0.0022 (10)	0.0085 (10)	0.0005 (10)
O9	0.0202 (11)	0.0301 (14)	0.0204 (12)	0.0001 (10)	0.0031 (9)	-0.0020 (10)
O10	0.0210 (12)	0.0319 (14)	0.0230 (13)	0.0031 (10)	0.0039 (9)	0.0037 (10)

O11	0.0404 (16)	0.0291 (15)	0.0460 (18)	0.0150 (12)	0.0087 (13)	0.0000 (13)
O12	0.0199 (12)	0.0370 (16)	0.0395 (16)	-0.0061 (11)	0.0037 (11)	-0.0016 (12)
O13	0.0197 (11)	0.0184 (12)	0.0326 (14)	-0.0008 (9)	0.0041 (10)	-0.0020 (10)
O14	0.0205 (11)	0.0198 (12)	0.0294 (13)	-0.0002 (9)	0.0049 (10)	-0.0038 (10)
O15	0.0407 (17)	0.0327 (17)	0.069 (2)	0.0003 (13)	0.0012 (15)	0.0261 (16)
O16	0.0383 (16)	0.0337 (16)	0.060 (2)	-0.0041 (13)	0.0110 (14)	-0.0263 (15)
O1w	0.0227 (12)	0.0217 (12)	0.0302 (14)	-0.0015 (10)	0.0076 (10)	-0.0023 (10)
O2w	0.049 (2)	0.054 (2)	0.086 (3)	-0.0027 (17)	0.0191 (19)	0.005 (2)
O3w	0.053 (2)	0.064 (2)	0.056 (2)	-0.0085 (18)	0.0129 (17)	-0.0049 (18)
N1	0.0322 (17)	0.0359 (19)	0.0298 (17)	0.0104 (14)	0.0005 (13)	-0.0055 (14)
N2	0.0296 (16)	0.0261 (17)	0.044 (2)	-0.0017 (13)	0.0034 (14)	-0.0044 (15)
N3	0.037 (2)	0.076 (3)	0.047 (2)	0.022 (2)	0.0011 (17)	-0.008 (2)
N4	0.0301 (18)	0.033 (2)	0.073 (3)	-0.0063 (15)	0.0152 (18)	-0.0067 (18)
C1	0.033 (2)	0.029 (2)	0.030 (2)	0.0023 (16)	0.0053 (15)	-0.0011 (16)
C2	0.0249 (18)	0.035 (2)	0.037 (2)	-0.0001 (16)	0.0035 (15)	-0.0076 (17)
C3	0.033 (2)	0.041 (2)	0.031 (2)	0.0027 (17)	0.0079 (16)	-0.0011 (17)
C4	0.035 (2)	0.046 (3)	0.049 (3)	0.0014 (19)	0.0043 (19)	-0.012 (2)
C5	0.037 (2)	0.035 (2)	0.042 (2)	0.0055 (18)	0.0066 (18)	-0.0005 (18)
C6	0.037 (2)	0.049 (3)	0.062 (3)	-0.007 (2)	0.012 (2)	-0.015 (2)

Geometric parameters (Å, °)

Ce1—O1	2.252 (2)	N1—C1	1.479 (5)
Ce1—O8 ⁱ	2.352 (2)	N1—H11	0.8600
Ce1—O9	2.362 (2)	N1—H12	0.8600
Ce1—O1w	2.385 (2)	N1—H13	0.8600
Ce1—O13	2.399 (2)	N2—C3	1.484 (5)
Ce1—O14	2.413 (2)	N2—H21	0.8600
Ce1—O10	2.431 (2)	N2—H22	0.8600
Ce1—O6	2.433 (2)	N2—H23	0.8600
Ce1—O5	2.467 (2)	N3—C4	1.450 (6)
S1—O3	1.444 (3)	N3—H31	0.8600
S1—O4	1.448 (3)	N3—H32	0.8600
S1—O2	1.459 (3)	N3—H33	0.8600
S1—O1	1.515 (2)	N4—C6	1.460 (6)
S2—O7	1.427 (3)	N4—H41	0.8600
S2—O8	1.477 (2)	N4—H42	0.8600
S2—O5	1.485 (3)	N4—H43	0.8600
S2—O6	1.503 (2)	C1—C2	1.514 (5)
S3—O11	1.437 (3)	C1—H1A	0.9700
S3—O12	1.441 (3)	C1—H1B	0.9700
S3—O10	1.508 (3)	C2—C3	1.510 (5)
S3—O9	1.524 (3)	C2—H2A	0.9700
S4—O15	1.448 (3)	C2—H2B	0.9700
S4—O16	1.449 (3)	C3—H3A	0.9700
S4—O14	1.492 (2)	C3—H3B	0.9700
S4—O13	1.505 (2)	C4—C5	1.496 (6)
O8—Ce1 ⁱⁱ	2.352 (2)	C4—H4A	0.9700
O1w—H1w1	0.8368	C4—H4B	0.9700

supplementary materials

O1w—H1w2	0.8351	C5—C6	1.486 (6)
O2w—H2w1	0.8401	C5—H5A	0.9700
O2w—H2w2	0.8390	C5—H5B	0.9700
O3w—H3w1	0.8444	C6—H6A	0.9700
O3w—H3w2	0.8478	C6—H6B	0.9700
O1—Ce1—O8 ⁱ	79.66 (9)	S4—O14—Ce1	99.66 (12)
O1—Ce1—O9	91.34 (9)	Ce1—O1w—H1w1	110.1
O8 ⁱ —Ce1—O9	126.76 (9)	Ce1—O1w—H1w2	132.2
O1—Ce1—O1w	77.33 (9)	H1w1—O1w—H1w2	109.9
O8 ⁱ —Ce1—O1w	83.33 (9)	H2w1—O2w—H2w2	109.4
O9—Ce1—O1w	145.79 (8)	H3w1—O3w—H3w2	107.6
O1—Ce1—O13	148.37 (9)	C1—N1—H11	109.5
O8 ⁱ —Ce1—O13	81.78 (9)	C1—N1—H12	109.5
O9—Ce1—O13	79.59 (8)	H11—N1—H12	109.5
O1w—Ce1—O13	125.54 (8)	C1—N1—H13	109.5
O1—Ce1—O14	138.98 (8)	H11—N1—H13	109.5
O8 ⁱ —Ce1—O14	75.94 (9)	H12—N1—H13	109.5
O9—Ce1—O14	129.67 (8)	C3—N2—H21	109.5
O1w—Ce1—O14	67.56 (8)	C3—N2—H22	109.5
O13—Ce1—O14	58.03 (8)	H21—N2—H22	109.5
O1—Ce1—O10	76.20 (9)	C3—N2—H23	109.5
O8 ⁱ —Ce1—O10	68.77 (8)	H21—N2—H23	109.5
O9—Ce1—O10	58.17 (8)	H22—N2—H23	109.5
O1w—Ce1—O10	144.34 (9)	C4—N3—H31	109.5
O13—Ce1—O10	73.27 (8)	C4—N3—H32	109.5
O14—Ce1—O10	122.81 (8)	H31—N3—H32	109.5
O1—Ce1—O6	129.18 (9)	C4—N3—H33	109.5
O8 ⁱ —Ce1—O6	146.41 (8)	H31—N3—H33	109.5
O9—Ce1—O6	75.37 (8)	H32—N3—H33	109.5
O1w—Ce1—O6	86.94 (8)	C6—N4—H41	109.5
O13—Ce1—O6	78.00 (8)	C6—N4—H42	109.5
O14—Ce1—O6	70.63 (8)	H41—N4—H42	109.5
O10—Ce1—O6	128.51 (8)	C6—N4—H43	109.5
O1—Ce1—O5	72.31 (9)	H41—N4—H43	109.5
O8 ⁱ —Ce1—O5	145.65 (8)	H42—N4—H43	109.5
O9—Ce1—O5	74.19 (8)	N1—C1—C2	109.4 (3)
O1w—Ce1—O5	71.61 (9)	N1—C1—H1A	109.8
O13—Ce1—O5	131.94 (8)	C2—C1—H1A	109.8
O14—Ce1—O5	113.52 (9)	N1—C1—H1B	109.8
O10—Ce1—O5	121.19 (9)	C2—C1—H1B	109.8
O6—Ce1—O5	56.88 (8)	H1A—C1—H1B	108.2
O3—S1—O4	111.3 (2)	C3—C2—C1	113.3 (3)
O3—S1—O2	110.1 (2)	C3—C2—H2A	108.9
O4—S1—O2	111.7 (2)	C1—C2—H2A	108.9
O3—S1—O1	109.16 (17)	C3—C2—H2B	108.9
O4—S1—O1	106.74 (18)	C1—C2—H2B	108.9
O2—S1—O1	107.59 (16)	H2A—C2—H2B	107.7

O7—S2—O8	112.34 (16)	N2—C3—C2	113.0 (3)
O7—S2—O5	113.34 (18)	N2—C3—H3A	109.0
O8—S2—O5	108.60 (15)	C2—C3—H3A	109.0
O7—S2—O6	111.47 (17)	N2—C3—H3B	109.0
O8—S2—O6	107.82 (14)	C2—C3—H3B	109.0
O5—S2—O6	102.71 (14)	H3A—C3—H3B	107.8
O11—S3—O12	113.18 (17)	N3—C4—C5	111.7 (4)
O11—S3—O10	111.37 (17)	N3—C4—H4A	109.3
O12—S3—O10	111.43 (16)	C5—C4—H4A	109.3
O11—S3—O9	109.67 (17)	N3—C4—H4B	109.3
O12—S3—O9	110.01 (16)	C5—C4—H4B	109.3
O10—S3—O9	100.44 (14)	H4A—C4—H4B	107.9
O15—S4—O16	111.8 (2)	C6—C5—C4	111.5 (4)
O15—S4—O14	110.90 (17)	C6—C5—H5A	109.3
O16—S4—O14	110.85 (16)	C4—C5—H5A	109.3
O15—S4—O13	110.46 (17)	C6—C5—H5B	109.3
O16—S4—O13	110.17 (17)	C4—C5—H5B	109.3
O14—S4—O13	102.30 (14)	H5A—C5—H5B	108.0
S1—O1—Ce1	145.18 (15)	N4—C6—C5	113.2 (4)
S2—O5—Ce1	99.61 (12)	N4—C6—H6A	108.9
S2—O6—Ce1	100.49 (12)	C5—C6—H6A	108.9
S2—O8—Ce1 ⁱⁱ	144.13 (15)	N4—C6—H6B	108.9
S3—O9—Ce1	101.91 (12)	C5—C6—H6B	108.9
S3—O10—Ce1	99.48 (12)	H6A—C6—H6B	107.7
S4—O13—Ce1	99.86 (12)		
O3—S1—O1—Ce1	96.8 (3)	O13—Ce1—O9—S3	76.58 (13)
O4—S1—O1—Ce1	-142.7 (3)	O14—Ce1—O9—S3	108.19 (13)
O2—S1—O1—Ce1	-22.7 (4)	O10—Ce1—O9—S3	-0.09 (11)
O8 ⁱ —Ce1—O1—S1	112.7 (3)	O6—Ce1—O9—S3	156.76 (14)
O9—Ce1—O1—S1	-120.1 (3)	O5—Ce1—O9—S3	-144.11 (14)
O1w—Ce1—O1—S1	27.3 (3)	O11—S3—O10—Ce1	-116.21 (16)
O13—Ce1—O1—S1	167.8 (2)	O12—S3—O10—Ce1	116.37 (15)
O14—Ce1—O1—S1	58.6 (3)	O9—S3—O10—Ce1	-0.13 (15)
O10—Ce1—O1—S1	-176.8 (3)	O1—Ce1—O10—S3	100.51 (14)
O6—Ce1—O1—S1	-48.0 (3)	O8 ⁱ —Ce1—O10—S3	-175.39 (15)
O5—Ce1—O1—S1	-47.2 (3)	O9—Ce1—O10—S3	0.09 (11)
O7—S2—O5—Ce1	-115.29 (16)	O1w—Ce1—O10—S3	143.65 (13)
O8—S2—O5—Ce1	119.11 (13)	O13—Ce1—O10—S3	-87.88 (13)
O6—S2—O5—Ce1	5.10 (15)	O14—Ce1—O10—S3	-119.49 (13)
O1—Ce1—O5—S2	177.08 (15)	O6—Ce1—O10—S3	-28.99 (18)
O8 ⁱ —Ce1—O5—S2	140.24 (13)	O5—Ce1—O10—S3	41.46 (16)
O9—Ce1—O5—S2	-86.27 (13)	O15—S4—O13—Ce1	-114.54 (17)
O1w—Ce1—O5—S2	94.93 (14)	O16—S4—O13—Ce1	121.47 (16)
O13—Ce1—O5—S2	-26.73 (18)	O14—S4—O13—Ce1	3.56 (15)
O14—Ce1—O5—S2	40.62 (15)	O1—Ce1—O13—S4	-135.59 (15)
O10—Ce1—O5—S2	-121.98 (13)	O8 ⁱ —Ce1—O13—S4	-81.04 (13)
O6—Ce1—O5—S2	-3.67 (11)	O9—Ce1—O13—S4	149.06 (14)
O7—S2—O6—Ce1	116.49 (16)	O1w—Ce1—O13—S4	-5.37 (17)

supplementary materials

O8—S2—O6—Ce1	−119.76 (13)	O14—Ce1—O13—S4	−2.54 (11)
O5—S2—O6—Ce1	−5.19 (15)	O10—Ce1—O13—S4	−151.25 (14)
O1—Ce1—O6—S2	4.56 (17)	O6—Ce1—O13—S4	71.98 (13)
O8 ⁱ —Ce1—O6—S2	−139.44 (13)	O5—Ce1—O13—S4	91.57 (15)
O9—Ce1—O6—S2	84.09 (12)	O15—S4—O14—Ce1	114.25 (18)
O1w—Ce1—O6—S2	−66.35 (12)	O16—S4—O14—Ce1	−120.97 (16)
O13—Ce1—O6—S2	166.31 (13)	O13—S4—O14—Ce1	−3.54 (15)
O14—Ce1—O6—S2	−133.63 (14)	O1—Ce1—O14—S4	146.82 (12)
O10—Ce1—O6—S2	109.35 (13)	O8 ⁱ —Ce1—O14—S4	91.56 (13)
O5—Ce1—O6—S2	3.63 (11)	O9—Ce1—O14—S4	−34.86 (17)
O7—S2—O8—Ce1 ⁱⁱ	9.8 (3)	O1w—Ce1—O14—S4	−179.93 (15)
O5—S2—O8—Ce1 ⁱⁱ	136.0 (2)	O13—Ce1—O14—S4	2.56 (11)
O6—S2—O8—Ce1 ⁱⁱ	−113.4 (3)	O10—Ce1—O14—S4	38.84 (16)
O11—S3—O9—Ce1	117.48 (15)	O6—Ce1—O14—S4	−85.13 (13)
O12—S3—O9—Ce1	−117.42 (15)	O5—Ce1—O14—S4	−123.43 (12)
O10—S3—O9—Ce1	0.13 (15)	N1—C1—C2—C3	168.6 (3)
O1—Ce1—O9—S3	−72.92 (13)	C1—C2—C3—N2	70.0 (5)
O8 ⁱ —Ce1—O9—S3	5.16 (17)	N3—C4—C5—C6	172.7 (4)
O1w—Ce1—O9—S3	−142.07 (13)	C4—C5—C6—N4	−174.8 (4)


Symmetry codes: (i) $x, -y+1/2, z+1/2$; (ii) $x, -y+1/2, z-1/2$.

Hydrogen-bond geometry (\AA , $^\circ$)

$D\cdots H$	$D—H$	$H\cdots A$	$D\cdots A$	$D—H\cdots A$
O1w—H1w2 \cdots O12 ⁱⁱⁱ	0.84	2.05	2.854 (4)	162
O2w—H2w1 \cdots O11 ⁱⁱⁱ	0.84	2.24	2.935 (5)	140
O2w—H2w2 \cdots O15 ^{iv}	0.84	2.08	2.902 (5)	167
O3w—H3w1 \cdots O3 ⁱⁱⁱ	0.84	1.90	2.729 (5)	169
O3w—H3w2 \cdots O2w	0.85	1.98	2.802 (5)	163
N1—H11 \cdots O9 ⁱⁱⁱ	0.86	2.04	2.881 (4)	167
N1—H12 \cdots O3w	0.86	1.95	2.806 (5)	171
N1—H13 \cdots O13 ^v	0.86	2.19	3.036 (4)	166
N2—H21 \cdots O16 ⁱⁱ	0.86	2.09	2.904 (5)	157
N2—H22 \cdots O16 ^{iv}	0.86	2.25	2.978 (5)	142
N2—H23 \cdots O2	0.86	2.49	3.180 (6)	137
N2—H23 \cdots O5	0.86	2.44	2.989 (4)	122
N3—H31 \cdots O6 ⁱ	0.86	2.04	2.866 (4)	162
N3—H32 \cdots O4	0.86	2.37	2.882 (6)	119
N3—H32 \cdots O2w ^{vi}	0.86	2.37	3.085 (6)	141
N3—H33 \cdots O1	0.86	2.33	3.004 (5)	135
N3—H33 \cdots O10	0.86	2.43	3.191 (6)	147
N4—H41 \cdots O6 ^{vii}	0.86	2.34	2.986 (4)	132
N4—H41 \cdots O15 ^{vii}	0.86	2.25	2.951 (5)	138
N4—H43 \cdots O4 ^{viii}	0.86	2.00	2.851 (5)	173

Symmetry codes: (iii) $x-1, y, z$; (iv) $-x+1, y-1/2, -z+1/2$; (v) $x-1, -y+1/2, z-1/2$; (ii) $x, -y+1/2, z-1/2$; (i) $x, -y+1/2, z+1/2$; (vi) $-x+1, -y, -z+1$; (vii) $x+1, -y+1/2, z+1/2$; (viii) $-x+2, -y, -z+1$.

Fig. 1



supplementary materials

Fig. 2

