

***trans*-Diaquabis(cyclohexane-1,2-diamine)zinc(II) dichloride**

Kamelia Karimnejad, Hamid Khaledi* and Hapipah Mohd Ali

Department of Chemistry, University of Malaya, 50603 Kuala Lumpur, Malaysia
Correspondence e-mail: khaledi@siswa.um.edu.my


Received 28 February 2011; accepted 3 March 2011

Key indicators: single-crystal X-ray study; $T = 100$ K; mean $\sigma(\text{C}-\text{C}) = 0.002$ Å; R factor = 0.021; wR factor = 0.053; data-to-parameter ratio = 16.4.

In the title compound, $[\text{Zn}(\text{C}_6\text{H}_{14}\text{N}_2)(\text{H}_2\text{O})_2]\text{Cl}_2$, the Zn(II) atom resides on a special position with site symmetry $2/m$ and is octahedrally coordinated by four N atoms from two *trans* 1,2-diaminocyclohexane ligands and two water O atoms. In the crystal, $\text{N}-\text{H}\cdots\text{Cl}$ and $\text{O}-\text{H}\cdots\text{Cl}$ hydrogen bonds link the molecules into a two-dimensional network parallel to the bc plane.

Related literature

For an isotopic nickel(II) complex, see: Capilla *et al.* (1980) and for an analogous copper(II) complex, see: Pariya *et al.* (1998).

**Experimental***Crystal data*

$[\text{Zn}(\text{C}_6\text{H}_{14}\text{N}_2)_2(\text{H}_2\text{O})_2]\text{Cl}_2$
 $M_r = 400.69$
Orthorhombic, $Cmce$

$a = 24.6478$ (4) Å
 $b = 9.5107$ (2) Å
 $c = 7.6723$ (2) Å

$V = 1798.52$ (7) Å³
 $Z = 4$
Mo $K\alpha$ radiation

$\mu = 1.67$ mm⁻¹
 $T = 100$ K
 $0.26 \times 0.21 \times 0.04$ mm

Data collection

Bruker APEXII CCD diffractometer
Absorption correction: multi-scan (*SADABS*; Sheldrick, 1996)
 $T_{\min} = 0.670$, $T_{\max} = 0.936$

4324 measured reflections
999 independent reflections
805 reflections with $I > 2\sigma(I)$
 $R_{\text{int}} = 0.019$

Refinement

$R[F^2 > 2\sigma(F^2)] = 0.021$
 $wR(F^2) = 0.053$
 $S = 1.04$
999 reflections
61 parameters
3 restraints

H atoms treated by a mixture of independent and constrained refinement
 $\Delta\rho_{\max} = 0.36$ e Å⁻³
 $\Delta\rho_{\min} = -0.24$ e Å⁻³

Table 1
Hydrogen-bond geometry (Å, °).

| $D-\text{H}\cdots A$ | $D-\text{H}$ | $\text{H}\cdots A$ | $D\cdots A$ | $D-\text{H}\cdots A$ |
|--|--------------|--------------------|-------------|----------------------|
| $\text{N1}-\text{H1A}\cdots\text{Cl1}^i$ | 0.87 (1) | 2.80 (1) | 3.6139 (13) | 156 (1) |
| $\text{N1}-\text{H1B}\cdots\text{Cl1}$ | 0.87 (1) | 2.70 (1) | 3.5206 (14) | 157 (1) |
| $\text{O1}-\text{H1O}\cdots\text{Cl1}$ | 0.83 (1) | 2.26 (1) | 3.0857 (12) | 173 (2) |

Symmetry code: (i) $x, y + \frac{1}{2}, -z + \frac{1}{2}$.

Data collection: *APEX2* (Bruker, 2007); cell refinement: *SAINT* (Bruker, 2007); data reduction: *SAINT*; program(s) used to solve structure: *SHELXS97* (Sheldrick, 2008); program(s) used to refine structure: *SHELXL97* (Sheldrick, 2008); molecular graphics: *X-SEED* (Barbour, 2001); software used to prepare material for publication: *SHELXL97* and *pubCIF* (Westrip, 2010).

The authors thank University of Malaya for funding this study (FRGS grant No. FP004/2010B).

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: PV2395).

References

- Barbour, L. J. (2001). *J. Supramol. Chem.*, **1**, 189–191.
Bruker (2007). *APEX2* and *SAINT*. Bruker AXS Inc., Madison, Wisconsin, USA.
Capilla, A. V., Aranda, R. A. & Gomez-Beltran, F. (1980). *Cryst. Struct. Commun.* **9**, 147–150.
Pariya, C., Liao, F.-L., Wang, S.-L. & Chung, C.-S. (1998). *Polyhedron*, **17**, 547–554.
Sheldrick, G. M. (1996). *SADABS*. University of Göttingen, Germany.
Sheldrick, G. M. (2008). *Acta Cryst.* **A64**, 112–122.
Westrip, S. P. (2010). *J. Appl. Cryst.* **43**, 920–925.

supplementary materials

Acta Cryst. (2011). E67, m421 [doi:10.1107/S1600536811008166]

***trans*-Diaquabis(cyclohexane-1,2-diamine)zinc(II) dichloride**

K. Karimnejad, H. Khaledi and H. Mohd Ali

Comment

The title Zn^{II} complex (Fig. 1) is isostructural with a previously reported Ni^{II} complex (Capilla *et al.*, 1980). The metal center in the title complex is located on a special position with site symmetry $2/m$, and is six-coordinated in a distorted octahedral geometry. The equatorial plane is defined by four N atoms from two *trans*-1,2-diaminocyclohexane, while the axial positions are occupied by two water molecule oxygen atoms. This arrangement is similar to what was observed in an analogous Cu^{II} complex (Pariya *et al.*, 1998). In the crystal, an O—H...Cl interaction connects the cationic complexes and chloride anions into infinite chains along the *c*-axis. The chains are further linked by N—H...Cl hydrogen bonds into layers parallel to the *bc* plane.

Experimental

The colorless crystals of the title compound were obtained upon slow evaporation of an ethanolic solution (20 ml) of *trans*-1,2-diaminocyclohexane (0.23 g, 2 mmol) and zinc(II) acetate dihydrate (0.22 g, 1 mmol) in the presence of a few drops of HCl (37%).

Refinement

The C-bound hydrogen atoms were placed at calculated positions with C—H = 0.99 and 1.00 Å for methylene and methyne type H-atoms, respectively. The hydrogen atoms bonded to N and water molecule were located from a difference map and included with restraints: O—H = 0.84 (2) and N—H = 0.88 (2) Å. For hydrogen atoms *U*_{iso}(H) were set to 1.2–1.5 times *U*_{eq}(carrier atom).

Figures




Fig. 1. Thermal ellipsoid plot of the title compound at the 50% probability level. Hydrogen atoms are drawn as spheres of arbitrary radius. Symmetry codes: ' = -x, -y, -z; '' = x, -y, -z; ''' = -x, y, z.

***trans*-Diaquabis(cyclohexane-1,2-diamine)zinc(II) dichloride**

Crystal data

[Zn(C₆H₁₄N₂)₂(H₂O)₂]Cl₂

M_r = 400.69

Orthorhombic, *Cmce*

F(000) = 848

D_x = 1.480 Mg m⁻³

Mo *K*α radiation, λ = 0.71073 Å

supplementary materials

Hall symbol: -C 2bc 2
 $a = 24.6478$ (4) Å
 $b = 9.5107$ (2) Å
 $c = 7.6723$ (2) Å
 $V = 1798.52$ (7) Å³
 $Z = 4$

Cell parameters from 2315 reflections
 $\theta = 3.3\text{--}30.5^\circ$
 $\mu = 1.67$ mm⁻¹
 $T = 100$ K
Prism, colorless
 $0.26 \times 0.21 \times 0.04$ mm

Data collection

Bruker APEXII CCD
diffractometer
Radiation source: fine-focus sealed tube
graphite
 φ and ω scans
Absorption correction: multi-scan
(*SADABS*; Sheldrick, 1996)
 $T_{\min} = 0.670$, $T_{\max} = 0.936$
4324 measured reflections

999 independent reflections
805 reflections with $I > 2\sigma(I)$
 $R_{\text{int}} = 0.019$
 $\theta_{\max} = 27.0^\circ$, $\theta_{\min} = 3.3^\circ$
 $h = -31 \rightarrow 31$
 $k = -11 \rightarrow 12$
 $l = -9 \rightarrow 8$

Refinement

Refinement on F^2
Least-squares matrix: full
 $R[F^2 > 2\sigma(F^2)] = 0.021$
 $wR(F^2) = 0.053$
 $S = 1.04$
999 reflections
61 parameters
3 restraints

Primary atom site location: structure-invariant direct methods
Secondary atom site location: difference Fourier map
Hydrogen site location: inferred from neighbouring sites
H atoms treated by a mixture of independent and constrained refinement
 $w = 1/[\sigma^2(F_o^2) + (0.0239P)^2 + 1.6417P]$
where $P = (F_o^2 + 2F_c^2)/3$
 $(\Delta/\sigma)_{\max} < 0.001$
 $\Delta\rho_{\max} = 0.36$ e Å⁻³
 $\Delta\rho_{\min} = -0.23$ e Å⁻³

Special details

Geometry. All e.s.d.'s (except the e.s.d. in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell e.s.d.'s are taken into account individually in the estimation of e.s.d.'s in distances, angles and torsion angles; correlations between e.s.d.'s in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell e.s.d.'s is used for estimating e.s.d.'s involving l.s. planes.

Refinement. Refinement of F^2 against ALL reflections. The weighted R -factor wR and goodness of fit S are based on F^2 , conventional R -factors R are based on F , with F set to zero for negative F^2 . The threshold expression of $F^2 > \sigma(F^2)$ is used only for calculating R -factors(gt) *etc.* and is not relevant to the choice of reflections for refinement. R -factors based on F^2 are statistically about twice as large as those based on F , and R -factors based on ALL data will be even larger.

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (\AA^2)

| | x | y | z | $U_{\text{iso}}^*/U_{\text{eq}}$ |
|-----|---------------|---------------|--------------|----------------------------------|
| Zn1 | 0.0000 | 0.0000 | 0.0000 | 0.01354 (11) |
| O1 | 0.0000 | -0.10232 (16) | 0.2580 (2) | 0.0194 (3) |
| H1O | 0.0266 (6) | -0.0795 (18) | 0.318 (2) | 0.029* |
| N1 | 0.06632 (5) | 0.13128 (13) | 0.07888 (18) | 0.0163 (3) |
| H1A | 0.0617 (6) | 0.2190 (15) | 0.050 (2) | 0.020* |
| H1B | 0.0698 (7) | 0.1249 (17) | 0.1918 (18) | 0.020* |
| C1 | 0.11721 (5) | 0.08041 (15) | -0.0016 (2) | 0.0168 (3) |
| H1 | 0.1174 | 0.1111 | -0.1263 | 0.020* |
| C2 | 0.16808 (5) | 0.13853 (17) | 0.0850 (2) | 0.0205 (3) |
| H2A | 0.1678 | 0.1136 | 0.2102 | 0.025* |
| H2B | 0.1681 | 0.2423 | 0.0757 | 0.025* |
| C3 | 0.21934 (6) | 0.08004 (18) | 0.0001 (2) | 0.0255 (4) |
| H3A | 0.2218 | 0.1146 | -0.1213 | 0.031* |
| H3B | 0.2516 | 0.1145 | 0.0643 | 0.031* |
| Cl1 | 0.091868 (19) | 0.0000 | 0.5000 | 0.02019 (14) |

Atomic displacement parameters (\AA^2)

| | U^{11} | U^{22} | U^{33} | U^{12} | U^{13} | U^{23} |
|-----|--------------|--------------|--------------|-------------|-------------|---------------|
| Zn1 | 0.01056 (16) | 0.01421 (18) | 0.01585 (18) | 0.000 | 0.000 | -0.00039 (15) |
| O1 | 0.0165 (7) | 0.0220 (8) | 0.0197 (8) | 0.000 | 0.000 | 0.0020 (7) |
| N1 | 0.0158 (6) | 0.0144 (6) | 0.0188 (6) | 0.0013 (5) | 0.0016 (5) | -0.0005 (6) |
| C1 | 0.0132 (6) | 0.0175 (8) | 0.0197 (7) | 0.0003 (6) | 0.0009 (7) | 0.0003 (7) |
| C2 | 0.0159 (7) | 0.0209 (8) | 0.0247 (8) | -0.0033 (6) | -0.0007 (6) | -0.0013 (7) |
| C3 | 0.0148 (7) | 0.0274 (9) | 0.0342 (9) | -0.0031 (6) | -0.0007 (7) | 0.0014 (9) |
| Cl1 | 0.0188 (3) | 0.0243 (3) | 0.0174 (3) | 0.000 | 0.000 | -0.0004 (2) |

Geometric parameters (\AA , $^\circ$)

| | | | |
|---|-------------|-------------------------|-------------|
| Zn1—N1 ⁱ | 2.1440 (13) | C1—C2 | 1.5227 (19) |
| Zn1—N1 ⁱⁱ | 2.1440 (13) | C1—C1 ⁱⁱⁱ | 1.530 (3) |
| Zn1—N1 ⁱⁱⁱ | 2.1440 (13) | C1—H1 | 1.0000 |
| Zn1—N1 | 2.1441 (13) | C2—C3 | 1.526 (2) |
| Zn1—O1 | 2.2057 (16) | C2—H2A | 0.9900 |
| Zn1—O1 ⁱ | 2.2057 (16) | C2—H2B | 0.9900 |
| O1—H1O | 0.831 (13) | C3—C3 ⁱⁱⁱ | 1.522 (3) |
| N1—C1 | 1.4795 (18) | C3—H3A | 0.9900 |
| N1—H1A | 0.871 (14) | C3—H3B | 0.9900 |
| N1—H1B | 0.873 (13) | | |
| N1 ⁱ —Zn1—N1 ⁱⁱ | 80.65 (7) | Zn1—N1—H1B | 108.4 (11) |
| N1 ⁱ —Zn1—N1 ⁱⁱⁱ | 99.35 (7) | H1A—N1—H1B | 109.5 (16) |
| N1 ⁱⁱ —Zn1—N1 ⁱⁱⁱ | 180.00 (8) | N1—C1—C2 | 113.42 (12) |
| N1 ⁱ —Zn1—N1 | 180.0 | N1—C1—C1 ⁱⁱⁱ | 108.68 (10) |

supplementary materials

| | | | |
|--|------------|---------------------------|-------------|
| N1 ⁱⁱ —Zn1—N1 | 99.35 (7) | C2—C1—C1 ⁱⁱⁱ | 110.85 (10) |
| N1 ⁱⁱⁱ —Zn1—N1 | 80.65 (7) | N1—C1—H1 | 107.9 |
| N1 ⁱ —Zn1—O1 | 89.79 (5) | C2—C1—H1 | 107.9 |
| N1 ⁱⁱ —Zn1—O1 | 90.21 (5) | C1 ⁱⁱⁱ —C1—H1 | 107.9 |
| N1 ⁱⁱⁱ —Zn1—O1 | 89.79 (5) | C1—C2—C3 | 111.30 (13) |
| N1—Zn1—O1 | 90.21 (5) | C1—C2—H2A | 109.4 |
| N1 ⁱ —Zn1—O1 ⁱ | 90.21 (5) | C3—C2—H2A | 109.4 |
| N1 ⁱⁱ —Zn1—O1 ⁱ | 89.79 (5) | C1—C2—H2B | 109.4 |
| N1 ⁱⁱⁱ —Zn1—O1 ⁱ | 90.21 (5) | C3—C2—H2B | 109.4 |
| N1—Zn1—O1 ⁱ | 89.79 (5) | H2A—C2—H2B | 108.0 |
| O1—Zn1—O1 ⁱ | 180.00 (7) | C3 ⁱⁱⁱ —C3—C2 | 111.41 (11) |
| Zn1—O1—H1O | 112.5 (13) | C3 ⁱⁱⁱ —C3—H3A | 109.3 |
| C1—N1—Zn1 | 109.77 (9) | C2—C3—H3A | 109.3 |
| C1—N1—H1A | 108.5 (11) | C3 ⁱⁱⁱ —C3—H3B | 109.3 |
| Zn1—N1—H1A | 112.7 (11) | C2—C3—H3B | 109.3 |
| C1—N1—H1B | 107.9 (11) | H3A—C3—H3B | 108.0 |

Symmetry codes: (i) $-x, -y, -z$; (ii) $-x, y, z$; (iii) $x, -y, -z$.

Hydrogen-bond geometry (\AA , $^\circ$)

| $D-H\cdots A$ | $D-H$ | $H\cdots A$ | $D\cdots A$ | $D-H\cdots A$ |
|-----------------------------------|----------|-------------|-------------|---------------|
| N1—H1A \cdots C11 ^{iv} | 0.87 (1) | 2.80 (1) | 3.6139 (13) | 156 (1) |
| N1—H1B \cdots C11 | 0.87 (1) | 2.70 (1) | 3.5206 (14) | 157 (1) |
| O1—H1O \cdots C11 | 0.83 (1) | 2.26 (1) | 3.0857 (12) | 173 (2) |

Symmetry codes: (iv) $x, y+1/2, -z+1/2$.

Fig. 1

