

Bis[2-methoxy-6-(phenyliminomethyl)-phenolate- $\kappa^2 O,O'$]bis(thiocyanato- κN)-manganese(II)

Jin-Bei Shen,^a Guo-Di Ge^a and Guo-Liang Zhao^{a,b*}

^aCollege of Chemistry and Life Science, Zhejiang Normal University, Jinhua, Zhejiang 321004, People's Republic of China, and ^bXingzhi College, Zhejiang Normal University, Jinhua, Zhejiang 321004, People's Republic of China
Correspondence e-mail: sky53@zjnu.cn


Received 10 March 2011; accepted 11 March 2011

Key indicators: single-crystal X-ray study; $T = 296\text{ K}$; mean $\sigma(\text{C}-\text{C}) = 0.003\text{ \AA}$; R factor = 0.028; wR factor = 0.077; data-to-parameter ratio = 13.3.

The Mn^{II} atom in the title complex, $[\text{Mn}(\text{NCS})_2(\text{C}_{14}\text{H}_{13}\text{NO}_2)_2]$, lies on a center of inversion in a MnO_4N_2 octahedral geometry. The Schiff base is present in its zwitterionic form and is O,O' -chelated to the metal atom. The imino N atom is protonated and is involved in an intramolecular hydrogen bond with the phenolate O atom.

Related literature

For Schiff base ligands derived from *o*-vanillin and aniline and their rare earth complexes, see: Li *et al.* (2008); Liu *et al.* (2009); Xian *et al.* (2008); Zhao *et al.* (2005, 2007).



Experimental

Crystal data

| | |
|--|--|
| $[\text{Mn}(\text{NCS})_2(\text{C}_{14}\text{H}_{13}\text{NO}_2)_2]$ | $\gamma = 65.693 (1)^\circ$ |
| $M_r = 625.61$ | $V = 712.81 (3)\text{ \AA}^3$ |
| Triclinic, $P\bar{1}$ | $Z = 1$ |
| $a = 9.0204 (2)\text{ \AA}$ | Mo $K\alpha$ radiation |
| $b = 9.3070 (2)\text{ \AA}$ | $\mu = 0.65\text{ mm}^{-1}$ |
| $c = 9.4087 (2)\text{ \AA}$ | $T = 296\text{ K}$ |
| $\alpha = 87.417 (1)^\circ$ | $0.29 \times 0.17 \times 0.05\text{ mm}$ |
| $\beta = 82.010 (1)^\circ$ | |

Data collection

| | |
|--|--|
| Bruker APEXII area-detector diffractometer | 9483 measured reflections |
| Absorption correction: multi-scan (<i>SADABS</i> ; Sheldrick, 1996) | 2509 independent reflections |
| $R_{\text{int}} = 0.022$ | 2233 reflections with $I > 2\sigma(I)$ |
| $R_{\text{min}} = 0.877$, $T_{\text{max}} = 0.970$ | |

Refinement

| | |
|---------------------------------|---|
| $R[F^2 > 2\sigma(F^2)] = 0.028$ | 2 restraints |
| $wR(F^2) = 0.077$ | H-atom parameters constrained |
| $S = 1.06$ | $\Delta\rho_{\text{max}} = 0.17\text{ e \AA}^{-3}$ |
| 2509 reflections | $\Delta\rho_{\text{min}} = -0.24\text{ e \AA}^{-3}$ |
| 188 parameters | |

Table 1
Hydrogen-bond geometry (\AA , $^\circ$).

| $D-\text{H} \cdots A$ | $D-\text{H}$ | $\text{H} \cdots A$ | $D \cdots A$ | $D-\text{H} \cdots A$ |
|-----------------------|--------------|---------------------|--------------|-----------------------|
| N1—H1A \cdots O1 | 0.86 | 1.97 | 2.6501 (16) | 135 |

Data collection: *APEX2* (Bruker, 2006); cell refinement: *SAINT* (Bruker, 2006); data reduction: *SAINT*; program(s) used to solve structure: *SHELXS97* (Sheldrick, 2008); program(s) used to refine structure: *SHELXL97* (Sheldrick, 2008); molecular graphics: *SHELXTL* (Sheldrick, 2008); software used to prepare material for publication: *SHELXL97*.

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: NG5130).

References

- Bruker (2006). *APEX2* and *SAINT*. Bruker AXS Inc., Madison, Wisconsin, USA.
- Li, H.-Q., Xian, H.-D., Liu, J.-F. & Zhao, G.-L. (2008). *Acta Cryst. E* **64**, m1593–m1594.
- Liu, J.-F., Liu, J.-L. & Zhao, G.-L. (2009). *Acta Cryst. E* **65**, m1385–m1386.
- Sheldrick, G. M. (1996). *SADABS*. University of Göttingen, Germany.
- Sheldrick, G. M. (2008). *Acta Cryst. A* **64**, 112–122.
- Xian, H.-D., Liu, J.-F., Li, H.-Q. & Zhao, G.-L. (2008). *Acta Cryst. E* **64**, m1422.
- Zhao, G.-L., Shi, X. & Ng, S. W. (2007). *Acta Cryst. E* **63**, m267–m268.
- Zhao, G.-L., Zhang, P.-H. & Feng, Y.-L. (2005). *Chin. J. Inorg. Chem.* **21**, 421–424.

supplementary materials

Acta Cryst. (2011). E67, m463 [doi:10.1107/S1600536811009330]

Bis[2-methoxy-6-(phenyliminomethyl)phenolate- $\kappa^2 O,O'$]bis(thiocyanato- κN)manganese(II)

J.-B. Shen, G.-D. Ge and G.-L. Zhao

Comment

For many years, there has been considerable interest in the study of Schiff base compounds due to their biological activity (Zhao *et al.*, 2005). Interested in this field, we have been synthesized several analogous Schiff bases derived from *o*-vanillin and prepared their transitional and rare metal complexes further. In a few of articles we have reported our partial research results (Zhao *et al.*, 2007; Xian *et al.* 2008; Li *et al.* 2008; Liu *et al.* 2009). Herein, we describe a new Mn(II) complex.

The structure of complex (1) was shown in Fig. 1 and the coordination environment of Mn(II) was shown in Fig. 2. In this complex the Mn(II) is six-coordinated by two N atoms from thiocyanate ions and four O atoms from the Schiff bases (HL), which can be described as a distorted octahedral geometry. There thiocyanate anions coordinate to Mn(II) ion with N atoms occupying the apices and two HL ligands chelate the Mn(II) ion with four O atoms from deprotonated phenol groups and methoxyl groups occupying the equatorial positions. The Mn—O and Mn—N bond distances were listed in Table 1. The distances between Mn(II) and methoxyl O atoms are obvious longer than Mn—O(phenolic) bond distances, which are similar to the analogous complexes (Zhao *et al.*, 2007; Li *et al.*, 2008, Liu *et al.*, 2009).

The hydrogen bonds and $\pi-\pi$ weak non-covalent interactions lend stability to the structure. The stacking plot of this compound was shown in Fig. 3. In HL ligand, two protons of phenolic hydroxyl groups considered to have transferred to imine N atoms involve in forming intramolecular hydrogen bonds. The $\pi-\pi$ interactions exist both intra and extra molecules between the approximate paralleled participating benzene rings, which may be the primary forces keep the complex molecules packing together.

Experimental

Reagents and solvents used were of commercially available quality and without purified before using. The Schiff base ligand 2-(phenyliminomethyl)-6-methoxyphenol was synthesized from condensation of *o*-vanillin and aniline. The title compound was synthesized by traditional method. 1 mmol HL ligand was dissolved in ethanol, then 0.5 mmol Mn(NO₃)₂.6H₂O (in ethanol) was added to the upper solution. The mixture solution was stirred for 2 h at room temperature. Furthermore, 1 mmol NH₄SCN (dissolved in ethanol) was added. The mixture was stirred again for 8 h at room temperature. At last, deposit was filtered out and the reddish-brown solution was kept in the open air. The red crystal was obtained after several days.

Refinement

The structure was solved by direct methods and successive Fourier difference synthesis. The H atoms bonded to C and N atoms were positioned geometrically and refined using a riding model [aliphatic C—H = 0.96 Å ($U_{\text{iso}}(\text{H}) = 1.5U_{\text{eq}}(\text{C})$), aromatic C—H = 0.93 Å ($U_{\text{iso}}(\text{H}) = 1.2U_{\text{eq}}(\text{C})$) and N—H = 0.86 Å with $U_{\text{iso}}(\text{H}) = 1.2U_{\text{eq}}(\text{N})$].

supplementary materials

Figures




Fig. 1. The molecular structure of the title complex, showing the atom-labelling scheme. Displacement ellipsoids are drawn at the 30% probability level.




Fig. 2. The coordination environment of the Mn(II).




Fig. 3. The stacking plot of the title compound, showing H-bond interactions (dashed lines) and π - π stacking interactions.

Bis[2-methoxy-6-(phenyliminomethyl)phenolate- $\kappa^2 O,O'$]bis(thiocyanato- κN)manganese(II)

Crystal data

| | |
|--|---|
| $[\text{Mn}(\text{NCS})_2(\text{C}_{14}\text{H}_{13}\text{NO}_2)_2]$ | $Z = 1$ |
| $M_r = 625.61$ | $F(000) = 323$ |
| Triclinic, $P\bar{1}$ | $D_x = 1.457 \text{ Mg m}^{-3}$ |
| Hall symbol: -P 1 | Mo $K\alpha$ radiation, $\lambda = 0.71073 \text{ \AA}$ |
| $a = 9.0204 (2) \text{ \AA}$ | Cell parameters from 4271 reflections |
| $b = 9.3070 (2) \text{ \AA}$ | $\theta = 2.2\text{--}25.0^\circ$ |
| $c = 9.4087 (2) \text{ \AA}$ | $\mu = 0.65 \text{ mm}^{-1}$ |
| $\alpha = 87.417 (1)^\circ$ | $T = 296 \text{ K}$ |
| $\beta = 82.010 (1)^\circ$ | Block, red |
| $\gamma = 65.693 (1)^\circ$ | $0.29 \times 0.17 \times 0.05 \text{ mm}$ |
| $V = 712.81 (3) \text{ \AA}^3$ | |

Data collection

| | |
|---|--|
| Bruker APEXII area-detector diffractometer | 2509 independent reflections |
| Radiation source: fine-focus sealed tube graphite | 2233 reflections with $I > 2\sigma(I)$ $R_{\text{int}} = 0.022$ |

| | |
|--|---|
| φ and ω scans | $\theta_{\max} = 25.0^\circ, \theta_{\min} = 2.2^\circ$ |
| Absorption correction: multi-scan (SADABS; Sheldrick, 1996) | $h = -10 \rightarrow 10$ |
| $T_{\min} = 0.877, T_{\max} = 0.970$ | $k = -11 \rightarrow 11$ |
| 9483 measured reflections | $l = -11 \rightarrow 11$ |

Refinement

| | |
|---------------------------------|---|
| Refinement on F^2 | Primary atom site location: structure-invariant direct methods |
| Least-squares matrix: full | Secondary atom site location: difference Fourier map |
| $R[F^2 > 2\sigma(F^2)] = 0.028$ | Hydrogen site location: inferred from neighbouring sites |
| $wR(F^2) = 0.077$ | H-atom parameters constrained |
| $S = 1.06$ | $w = 1/[\sigma^2(F_o^2) + (0.0423P)^2 + 0.1242P]$ where $P = (F_o^2 + 2F_c^2)/3$ |
| 2509 reflections | $(\Delta/\sigma)_{\max} = 0.001$ |
| 188 parameters | $\Delta\rho_{\max} = 0.17 \text{ e \AA}^{-3}$ |
| 2 restraints | $\Delta\rho_{\min} = -0.24 \text{ e \AA}^{-3}$ |

Special details

Geometry. All e.s.d.'s (except the e.s.d. in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell e.s.d.'s are taken into account individually in the estimation of e.s.d.'s in distances, angles and torsion angles; correlations between e.s.d.'s in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell e.s.d.'s is used for estimating e.s.d.'s involving l.s. planes.

Refinement. Refinement of F^2 against ALL reflections. The weighted R -factor wR and goodness of fit S are based on F^2 , conventional R -factors R are based on F , with F set to zero for negative F^2 . The threshold expression of $F^2 > \sigma(F^2)$ is used only for calculating R -factors(gt) etc. and is not relevant to the choice of reflections for refinement. R -factors based on F^2 are statistically about twice as large as those based on F , and R -factors based on ALL data will be even larger.

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (\AA^2)

| | x | y | z | $U_{\text{iso}}^*/U_{\text{eq}}$ |
|-----|--------------|--------------|--------------|----------------------------------|
| Mn1 | 0.0000 | 0.5000 | 0.5000 | 0.03697 (13) |
| S1 | 0.00911 (6) | 0.11697 (6) | 0.14672 (6) | 0.05653 (16) |
| O1 | 0.24701 (13) | 0.46530 (13) | 0.41948 (12) | 0.0385 (3) |
| N1 | 0.48105 (16) | 0.51194 (15) | 0.24678 (14) | 0.0354 (3) |
| H1A | 0.3779 | 0.5397 | 0.2737 | 0.043* |
| O2 | 0.16157 (14) | 0.30666 (15) | 0.63303 (13) | 0.0465 (3) |
| C7 | 0.36297 (19) | 0.35112 (18) | 0.47429 (16) | 0.0320 (3) |
| C13 | 0.32723 (19) | 0.25949 (19) | 0.58901 (17) | 0.0361 (4) |
| C9 | 0.53201 (19) | 0.31088 (19) | 0.42564 (17) | 0.0354 (4) |
| C15 | 0.01407 (19) | 0.2348 (2) | 0.26679 (19) | 0.0401 (4) |
| C8 | 0.58045 (19) | 0.3952 (2) | 0.31424 (18) | 0.0389 (4) |
| H8A | 0.6924 | 0.3648 | 0.2870 | 0.047* |
| C4 | 0.5244 (2) | 0.59880 (19) | 0.13313 (17) | 0.0358 (4) |

supplementary materials

| | | | | |
|------|--------------|------------|--------------|------------|
| C3 | 0.6852 (2) | 0.5520 (2) | 0.06908 (19) | 0.0451 (4) |
| H3A | 0.7678 | 0.4615 | 0.0990 | 0.054* |
| C12 | 0.4481 (2) | 0.1394 (2) | 0.64754 (19) | 0.0452 (4) |
| H12A | 0.4207 | 0.0812 | 0.7213 | 0.054* |
| C5 | 0.4016 (2) | 0.7315 (2) | 0.08693 (19) | 0.0441 (4) |
| H5A | 0.2935 | 0.7612 | 0.1290 | 0.053* |
| C2 | 0.7210 (2) | 0.6418 (2) | -0.0399 (2) | 0.0537 (5) |
| H2A | 0.8288 | 0.6114 | -0.0832 | 0.064* |
| N2 | 0.01391 (19) | 0.3211 (2) | 0.35072 (18) | 0.0554 (4) |
| C10 | 0.6546 (2) | 0.1866 (2) | 0.4902 (2) | 0.0475 (4) |
| H10A | 0.7647 | 0.1625 | 0.4585 | 0.057* |
| C1 | 0.5992 (3) | 0.7757 (2) | -0.0853 (2) | 0.0551 (5) |
| H1B | 0.6248 | 0.8357 | -0.1583 | 0.066* |
| C11 | 0.6141 (2) | 0.1025 (2) | 0.5974 (2) | 0.0514 (5) |
| H11A | 0.6960 | 0.0202 | 0.6380 | 0.062* |
| C6 | 0.4396 (2) | 0.8202 (2) | -0.0220 (2) | 0.0529 (5) |
| H6A | 0.3570 | 0.9103 | -0.0526 | 0.063* |
| C14 | 0.1088 (3) | 0.2163 (3) | 0.7377 (3) | 0.0723 (7) |
| H14A | 0.1511 | 0.2175 | 0.8256 | 0.108* |
| H14B | 0.1487 | 0.1096 | 0.7035 | 0.108* |
| H14C | -0.0089 | 0.2609 | 0.7546 | 0.108* |

Atomic displacement parameters (\AA^2)

| | U^{11} | U^{22} | U^{33} | U^{12} | U^{13} | U^{23} |
|-----|--------------|-------------|-------------|---------------|---------------|--------------|
| Mn1 | 0.02590 (19) | 0.0477 (2) | 0.0377 (2) | -0.01664 (16) | -0.00189 (14) | 0.00535 (15) |
| S1 | 0.0503 (3) | 0.0525 (3) | 0.0652 (3) | -0.0210 (2) | 0.0012 (2) | -0.0120 (2) |
| O1 | 0.0265 (6) | 0.0450 (7) | 0.0419 (6) | -0.0142 (5) | -0.0032 (5) | 0.0135 (5) |
| N1 | 0.0272 (7) | 0.0401 (8) | 0.0386 (7) | -0.0153 (6) | 0.0018 (6) | 0.0017 (6) |
| O2 | 0.0352 (6) | 0.0583 (8) | 0.0466 (7) | -0.0229 (6) | -0.0025 (5) | 0.0208 (6) |
| C7 | 0.0309 (8) | 0.0344 (8) | 0.0321 (8) | -0.0152 (7) | -0.0033 (6) | 0.0011 (6) |
| C13 | 0.0345 (9) | 0.0419 (9) | 0.0347 (8) | -0.0188 (7) | -0.0038 (7) | 0.0027 (7) |
| C9 | 0.0306 (8) | 0.0381 (9) | 0.0355 (8) | -0.0132 (7) | -0.0018 (6) | 0.0021 (7) |
| C15 | 0.0272 (8) | 0.0426 (10) | 0.0477 (10) | -0.0136 (7) | 0.0009 (7) | 0.0048 (7) |
| C8 | 0.0263 (8) | 0.0460 (10) | 0.0418 (9) | -0.0139 (7) | 0.0010 (7) | 0.0012 (7) |
| C4 | 0.0367 (9) | 0.0379 (9) | 0.0359 (8) | -0.0199 (7) | 0.0000 (7) | -0.0004 (7) |
| C3 | 0.0359 (9) | 0.0495 (11) | 0.0469 (10) | -0.0175 (8) | 0.0016 (8) | 0.0083 (8) |
| C12 | 0.0488 (11) | 0.0445 (10) | 0.0409 (9) | -0.0184 (8) | -0.0078 (8) | 0.0120 (8) |
| C5 | 0.0382 (10) | 0.0441 (10) | 0.0468 (10) | -0.0154 (8) | -0.0008 (8) | 0.0022 (8) |
| C2 | 0.0426 (10) | 0.0636 (12) | 0.0546 (11) | -0.0263 (10) | 0.0068 (9) | 0.0085 (9) |
| N2 | 0.0465 (9) | 0.0663 (11) | 0.0562 (10) | -0.0280 (8) | 0.0035 (8) | -0.0098 (8) |
| C10 | 0.0306 (9) | 0.0516 (11) | 0.0512 (11) | -0.0089 (8) | -0.0038 (8) | 0.0063 (8) |
| C1 | 0.0620 (13) | 0.0575 (12) | 0.0500 (11) | -0.0318 (11) | -0.0014 (9) | 0.0142 (9) |
| C11 | 0.0422 (10) | 0.0481 (11) | 0.0531 (11) | -0.0073 (8) | -0.0115 (8) | 0.0139 (9) |
| C6 | 0.0541 (12) | 0.0450 (11) | 0.0542 (11) | -0.0160 (9) | -0.0076 (9) | 0.0124 (9) |
| C14 | 0.0556 (13) | 0.0807 (16) | 0.0782 (15) | -0.0333 (12) | 0.0049 (11) | 0.0363 (12) |

Geometric parameters (\AA , $^\circ$)

| | | | |
|--------------------------------------|-------------|--------------|-------------|
| Mn1—O1 | 2.1455 (10) | C4—C5 | 1.379 (2) |
| Mn1—O1 ⁱ | 2.1455 (10) | C4—C3 | 1.384 (2) |
| Mn1—N2 | 2.1794 (16) | C3—C2 | 1.381 (2) |
| Mn1—N2 ^j | 2.1794 (16) | C3—H3A | 0.9300 |
| Mn1—O2 | 2.2525 (12) | C12—C11 | 1.406 (3) |
| Mn1—O2 ⁱ | 2.2525 (12) | C12—H12A | 0.9300 |
| S1—C15 | 1.6270 (19) | C5—C6 | 1.381 (3) |
| O1—C7 | 1.2942 (19) | C5—H5A | 0.9300 |
| N1—C8 | 1.297 (2) | C2—C1 | 1.378 (3) |
| N1—C4 | 1.420 (2) | C2—H2A | 0.9300 |
| N1—H1A | 0.8600 | C10—C11 | 1.351 (3) |
| O2—C13 | 1.3790 (19) | C10—H10A | 0.9300 |
| O2—C14 | 1.424 (2) | C1—C6 | 1.376 (3) |
| C7—C9 | 1.423 (2) | C1—H1B | 0.9300 |
| C7—C13 | 1.429 (2) | C11—H11A | 0.9300 |
| C13—C12 | 1.360 (2) | C6—H6A | 0.9300 |
| C9—C8 | 1.409 (2) | C14—H14A | 0.9600 |
| C9—C10 | 1.414 (2) | C14—H14B | 0.9600 |
| C15—N2 | 1.151 (2) | C14—H14C | 0.9600 |
| C8—H8A | 0.9300 | | |
| O1—Mn1—O1 ⁱ | 180.0 | C5—C4—C3 | 120.42 (16) |
| O1—Mn1—N2 | 90.41 (5) | C5—C4—N1 | 118.22 (14) |
| O1 ⁱ —Mn1—N2 | 89.59 (5) | C3—C4—N1 | 121.36 (15) |
| O1—Mn1—N2 ^j | 89.59 (5) | C2—C3—C4 | 119.02 (17) |
| O1 ⁱ —Mn1—N2 ^j | 90.41 (5) | C2—C3—H3A | 120.5 |
| N2—Mn1—N2 ^j | 180.00 (7) | C4—C3—H3A | 120.5 |
| O1—Mn1—O2 | 74.23 (4) | C13—C12—C11 | 120.51 (16) |
| O1 ⁱ —Mn1—O2 | 105.77 (4) | C13—C12—H12A | 119.7 |
| N2—Mn1—O2 | 88.99 (6) | C11—C12—H12A | 119.7 |
| N2 ^j —Mn1—O2 | 91.01 (6) | C4—C5—C6 | 119.83 (16) |
| O1—Mn1—O2 ⁱ | 105.77 (4) | C4—C5—H5A | 120.1 |
| O1 ⁱ —Mn1—O2 ⁱ | 74.23 (4) | C6—C5—H5A | 120.1 |
| N2—Mn1—O2 ⁱ | 91.01 (6) | C1—C2—C3 | 120.86 (17) |
| N2 ^j —Mn1—O2 ⁱ | 88.99 (6) | C1—C2—H2A | 119.6 |
| O2—Mn1—O2 ⁱ | 180.00 (5) | C3—C2—H2A | 119.6 |
| C7—O1—Mn1 | 116.71 (9) | C15—N2—Mn1 | 175.35 (17) |
| C8—N1—C4 | 126.92 (14) | C11—C10—C9 | 120.88 (16) |
| C8—N1—H1A | 116.5 | C11—C10—H10A | 119.6 |
| C4—N1—H1A | 116.5 | C9—C10—H10A | 119.6 |
| C13—O2—C14 | 119.03 (14) | C6—C1—C2 | 119.66 (18) |
| C13—O2—Mn1 | 114.02 (9) | C6—C1—H1B | 120.2 |
| C14—O2—Mn1 | 126.09 (12) | C2—C1—H1B | 120.2 |
| O1—C7—C9 | 122.31 (14) | C10—C11—C12 | 120.05 (17) |

supplementary materials


| | | | |
|-----------------------------|--------------|-----------------|--------------|
| O1—C7—C13 | 121.33 (14) | C10—C11—H11A | 120.0 |
| C9—C7—C13 | 116.36 (14) | C12—C11—H11A | 120.0 |
| C12—C13—O2 | 124.73 (15) | C1—C6—C5 | 120.20 (18) |
| C12—C13—C7 | 121.84 (15) | C1—C6—H6A | 119.9 |
| O2—C13—C7 | 113.42 (14) | C5—C6—H6A | 119.9 |
| C8—C9—C10 | 118.78 (15) | O2—C14—H14A | 109.5 |
| C8—C9—C7 | 120.86 (15) | O2—C14—H14B | 109.5 |
| C10—C9—C7 | 120.35 (15) | H14A—C14—H14B | 109.5 |
| N2—C15—S1 | 178.24 (18) | O2—C14—H14C | 109.5 |
| N1—C8—C9 | 125.08 (14) | H14A—C14—H14C | 109.5 |
| N1—C8—H8A | 117.5 | H14B—C14—H14C | 109.5 |
| C9—C8—H8A | 117.5 | | |
| N2—Mn1—O1—C7 | −84.38 (11) | C13—C7—C9—C8 | −179.21 (15) |
| N2 ⁱ —Mn1—O1—C7 | 95.62 (11) | O1—C7—C9—C10 | 179.05 (15) |
| O2—Mn1—O1—C7 | 4.46 (10) | C13—C7—C9—C10 | −0.2 (2) |
| O2 ⁱ —Mn1—O1—C7 | −175.54 (10) | C4—N1—C8—C9 | −179.22 (15) |
| O1—Mn1—O2—C13 | −4.72 (10) | C10—C9—C8—N1 | −179.53 (16) |
| O1 ⁱ —Mn1—O2—C13 | 175.28 (10) | C7—C9—C8—N1 | −0.5 (3) |
| N2—Mn1—O2—C13 | 85.99 (11) | C8—N1—C4—C5 | −172.09 (16) |
| N2 ⁱ —Mn1—O2—C13 | −94.01 (11) | C8—N1—C4—C3 | 8.3 (3) |
| O1—Mn1—O2—C14 | −173.90 (17) | C5—C4—C3—C2 | 1.1 (3) |
| O1 ⁱ —Mn1—O2—C14 | 6.10 (17) | N1—C4—C3—C2 | −179.25 (16) |
| N2—Mn1—O2—C14 | −83.19 (17) | O2—C13—C12—C11 | −178.46 (16) |
| N2 ⁱ —Mn1—O2—C14 | 96.81 (17) | C7—C13—C12—C11 | 0.8 (3) |
| Mn1—O1—C7—C9 | 176.96 (11) | C3—C4—C5—C6 | −1.3 (3) |
| Mn1—O1—C7—C13 | −3.79 (19) | N1—C4—C5—C6 | 179.06 (16) |
| C14—O2—C13—C12 | −6.3 (3) | C4—C3—C2—C1 | −0.3 (3) |
| Mn1—O2—C13—C12 | −176.32 (13) | C8—C9—C10—C11 | 179.98 (17) |
| C14—O2—C13—C7 | 174.40 (17) | C7—C9—C10—C11 | 1.0 (3) |
| Mn1—O2—C13—C7 | 4.39 (17) | C3—C2—C1—C6 | −0.5 (3) |
| O1—C7—C13—C12 | −179.92 (15) | C9—C10—C11—C12 | −0.9 (3) |
| C9—C7—C13—C12 | −0.6 (2) | C13—C12—C11—C10 | 0.0 (3) |
| O1—C7—C13—O2 | −0.6 (2) | C2—C1—C6—C5 | 0.3 (3) |
| C9—C7—C13—O2 | 178.68 (13) | C4—C5—C6—C1 | 0.6 (3) |
| O1—C7—C9—C8 | 0.1 (2) | | |

Symmetry codes: (i) $-x, -y+1, -z+1$.

Hydrogen-bond geometry (\AA , $^\circ$)

| $D\text{—H}\cdots A$ | $D\text{—H}$ | $\text{H}\cdots A$ | $D\cdots A$ | $D\text{—H}\cdots A$ |
|----------------------|--------------|--------------------|-------------|----------------------|
| N1—H1A—O1 | 0.86 | 1.97 | 2.6501 (16) | 135. |

Fig. 1



supplementary materials

Fig. 2




Fig. 3

