

N-Ethyl-3,5-dinitrobenzamide

Jia-Ying Xu^{a*} and Wei-Hua Cheng^b

^aCollege of Chemical and Biological Engineering, Yancheng Institute of Technology, Yinbing Road No. 9 Yancheng, Yancheng 224051, People's Republic of China, and

^bDepartment of Chemical Engineering, Yancheng College of Textile Technology, Yancheng 224051, People's Republic of China

Correspondence e-mail: xuyiying1984@163.com


Received 24 February 2011; accepted 3 March 2011

Key indicators: single-crystal X-ray study; $T = 293\text{ K}$; mean $\sigma(\text{C}-\text{C}) = 0.003\text{ \AA}$; R factor = 0.051; wR factor = 0.161; data-to-parameter ratio = 12.7.

In the title molecule, $\text{C}_9\text{H}_9\text{N}_3\text{O}_5$, the dihedral angle between the mean planes of the amide group and the benzene ring is $31.24(14)^\circ$. In the crystal, $\text{N}-\text{H}\cdots\text{O}$ hydrogen bonds link the molecules to form one-dimensional chains propagating in [100].

Related literature

For the synthesis of the title compound, see: Lee *et al.* (2009). For standard bond-length data, see: Allen *et al.* (1987).



Experimental

Crystal data

$\text{C}_9\text{H}_9\text{N}_3\text{O}_5$
 $M_r = 239.19$
 Triclinic, $P\bar{1}$
 $a = 4.854(1)\text{ \AA}$
 $b = 10.488(2)\text{ \AA}$
 $c = 10.851(2)\text{ \AA}$

$\alpha = 101.49(3)^\circ$
 $\beta = 97.84(3)^\circ$
 $\gamma = 95.25(3)^\circ$
 $V = 532.26(19)\text{ \AA}^3$
 $Z = 2$
 Mo $K\alpha$ radiation

$\mu = 0.12\text{ mm}^{-1}$
 $T = 293\text{ K}$

$0.20 \times 0.10 \times 0.10\text{ mm}$

Data collection

Enraf–Nonius CAD-4
 diffractometer
 Absorption correction: ψ scan
 (North *et al.*, 1968)
 $T_{\min} = 0.976$, $T_{\max} = 0.988$
 2203 measured reflections

1955 independent reflections
 1321 reflections with $I > 2\sigma(I)$
 $R_{\text{int}} = 0.021$
 3 standard reflections every 200
 reflections
 intensity decay: 1%

Refinement

$R[F^2 > 2\sigma(F^2)] = 0.051$
 $wR(F^2) = 0.161$
 $S = 1.01$
 1955 reflections

154 parameters
 H-atom parameters constrained
 $\Delta\rho_{\max} = 0.15\text{ e \AA}^{-3}$
 $\Delta\rho_{\min} = -0.23\text{ e \AA}^{-3}$

Table 1
 Hydrogen-bond geometry (\AA , $^\circ$).

$D-\text{H}\cdots A$	$D-\text{H}$	$\text{H}\cdots A$	$D\cdots A$	$D-\text{H}\cdots A$
$\text{N}3-\text{H}3\text{A}\cdots\text{O}5^i$	0.86	2.13	2.886 (3)	146

Symmetry code: (i) $x + 1, y, z$.

Data collection: *CAD-4 Software* (Enraf–Nonius, 1985); cell refinement: *CAD-4 Software*; data reduction: *XCAD4* (Harms & Wocadlo, 1995); program(s) used to solve structure: *SHELXS97* (Sheldrick, 2008); program(s) used to refine structure: *SHELXL97* (Sheldrick, 2008); molecular graphics: *SHELXTL* (Sheldrick, 2008); software used to prepare material for publication: *SHELXTL*.

The authors thank the Center of Testing and Analysis, Nanjing University, for data collection.

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: SU2259).

References

- Allen, F. H., Kennard, O., Watson, D. G., Brammer, L., Orpen, A. G. & Taylor, R. (1987). *J. Chem. Soc. Perkin Trans. 2*, pp. S1–19.
- Enraf–Nonius (1985). *CAD-4 Software*. Enraf–Nonius, Delft, The Netherlands.
- Harms, K. & Wocadlo, S. (1995). *XCAD4*. University of Marburg, Germany.
- Lee, S., Song, K. H., Choe, J., Ju, J. & Jo, Y. (2009). *J. Org. Chem.* **74**, 6358–6361.
- North, A. C. T., Phillips, D. C. & Mathews, F. S. (1968). *Acta Cryst. A* **24**, 351–359.
- Sheldrick, G. M. (2008). *Acta Cryst. A* **64**, 112–122.

supplementary materials

Acta Cryst. (2011). E67, o822 [doi:10.1107/S160053681100804X]

N-Ethyl-3,5-dinitrobenzamide

J.-Y. Xu and W.-H. Cheng

Comment

The title compound is an important organic intermediate, and such amide derivatives exhibit biological activities, such as antibacterial and antifungal effects (Lee *et al.*, 2009). Herein we report on the crystal structure of the title substituted benzamide compound.

The molecular structure of the title compound is illustrated in Fig. 1. The bond lengths and angles are within normal ranges (Allen *et al.*, 1987). The dihedral angle between the amide group [atoms O5,N3,C7,C8, planar to within 0.012 Å] and the benzene ring (C1-C6) is 31.24 (14) °.

In the crystal of the title compound molecules are connected *via* N—H···O intermolecular hydrogen bonds (Table 1), to form a one-dimensional polymer propagating in [100]. These chains stack along the *c* axis direction.

Experimental

The title compound was prepared following a literature procedure (Lee *et al.*, 2009). Crystals, suitable for X-ray diffraction analysis, were obtained by slow evaporation, over a period of 5 days, of a solution of the title compound in ethanol [0.2 g, 0.84 mmol in 25 ml ethanol].

Refinement

All the H-atoms were positioned geometrically and constrained to ride on their parent atom: N—H = 0.86 Å, C—H = 0.93, 0.96 and 0.97 Å for CH(aromatic), CH₂ and CH₃ H-atoms, respectively, with $U_{\text{iso}}(\text{H}) = k \times U_{\text{eq}}(\text{C or N})$, where $k = 1.5$ for CH₃ H-atoms, and $k = 1.2$ for all other H-atoms.

Figures




Fig. 1. The molecular structure of the title molecule, with displacement ellipsoids drawn at the 50% probability level.

supplementary materials




Fig. 2. A view of the crystal packing of the title compound. The N—H···O hydrogen bonds are shown as dashed lines.

N-Ethyl-3,5-dinitrobenzamide

Crystal data

C ₉ H ₉ N ₃ O ₅	Z = 2
M _r = 239.19	F(000) = 248
Triclinic, P $\bar{1}$	D _x = 1.492 Mg m ⁻³
Hall symbol: -P 1	Mo K α radiation, λ = 0.71073 Å
a = 4.854 (1) Å	Cell parameters from 25 reflections
b = 10.488 (2) Å	θ = 10–13°
c = 10.851 (2) Å	μ = 0.12 mm ⁻¹
α = 101.49 (3)°	T = 293 K
β = 97.84 (3)°	Block, colourless
γ = 95.25 (3)°	0.20 × 0.10 × 0.10 mm
V = 532.26 (19) Å ³	

Data collection

Enraf–Nonius CAD-4 diffractometer	1321 reflections with $I > 2\sigma(I)$
Radiation source: fine-focus sealed tube graphite	$R_{\text{int}} = 0.021$
$\omega/2\theta$ scans	$\theta_{\text{max}} = 25.4^\circ$, $\theta_{\text{min}} = 1.9^\circ$
Absorption correction: ψ scan (North <i>et al.</i> , 1968)	$h = 0 \rightarrow 5$
$T_{\text{min}} = 0.976$, $T_{\text{max}} = 0.988$	$k = -12 \rightarrow 12$
2203 measured reflections	$l = -13 \rightarrow 12$
1955 independent reflections	3 standard reflections every 200 reflections intensity decay: 1%

Refinement

Refinement on F^2	Primary atom site location: structure-invariant direct methods
Least-squares matrix: full	Secondary atom site location: difference Fourier map

$R[F^2 > 2\sigma(F^2)] = 0.051$	Hydrogen site location: inferred from neighbouring sites
$wR(F^2) = 0.161$	H-atom parameters constrained
$S = 1.01$	$w = 1/[\sigma^2(F_o^2) + (0.090P)^2]$ where $P = (F_o^2 + 2F_c^2)/3$
1955 reflections	$(\Delta/\sigma)_{\max} < 0.001$
154 parameters	$\Delta\rho_{\max} = 0.15 \text{ e } \text{\AA}^{-3}$
0 restraints	$\Delta\rho_{\min} = -0.23 \text{ e } \text{\AA}^{-3}$

Special details

Geometry. All e.s.d.'s (except the e.s.d. in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell e.s.d.'s are taken into account individually in the estimation of e.s.d.'s in distances, angles and torsion angles; correlations between e.s.d.'s in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell e.s.d.'s is used for estimating e.s.d.'s involving l.s. planes.

Refinement. Refinement of F^2 against ALL reflections. The weighted R -factor wR and goodness of fit S are based on F^2 , conventional R -factors R are based on F , with F set to zero for negative F^2 . The threshold expression of $F^2 > \sigma(F^2)$ is used only for calculating R -factors(gt) etc. and is not relevant to the choice of reflections for refinement. R -factors based on F^2 are statistically about twice as large as those based on F , and R -factors based on ALL data will be even larger.

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (\AA^2)

	x	y	z	$U_{\text{iso}}^*/U_{\text{eq}}$
C1	0.1490 (5)	0.7075 (2)	0.2465 (2)	0.0428 (6)
H1A	-0.0017	0.6473	0.2494	0.051*
N1	0.0031 (5)	0.6713 (2)	0.0152 (2)	0.0617 (6)
O1	0.0675 (6)	0.6862 (3)	-0.0856 (2)	0.0985 (9)
N2	0.8248 (5)	0.9842 (2)	0.2289 (3)	0.0576 (6)
C2	0.1956 (5)	0.7377 (2)	0.1327 (2)	0.0454 (6)
O2	-0.2072 (5)	0.6034 (2)	0.02613 (19)	0.0795 (7)
N3	0.4868 (4)	0.7427 (2)	0.57021 (18)	0.0453 (5)
H3A	0.6513	0.7677	0.5563	0.054*
C3	0.4127 (5)	0.8280 (2)	0.1236 (2)	0.0492 (6)
H3B	0.4392	0.8480	0.0459	0.059*
O3	0.8469 (5)	1.0119 (2)	0.1269 (2)	0.0911 (8)
C4	0.5890 (5)	0.8874 (2)	0.2352 (2)	0.0449 (6)
O4	0.9911 (4)	1.0292 (2)	0.3254 (2)	0.0743 (6)
O5	0.0241 (3)	0.70717 (19)	0.49151 (17)	0.0574 (5)
C5	0.5555 (5)	0.8591 (2)	0.3510 (2)	0.0414 (6)
H5A	0.6803	0.8997	0.4241	0.050*
C6	0.3318 (4)	0.7688 (2)	0.3573 (2)	0.0388 (5)
C7	0.2693 (4)	0.7366 (2)	0.4798 (2)	0.0408 (6)
C8	0.4582 (5)	0.7087 (2)	0.6918 (2)	0.0497 (6)
H8A	0.2783	0.7293	0.7144	0.060*
H8B	0.6035	0.7612	0.7572	0.060*
C9	0.4794 (7)	0.5668 (3)	0.6881 (3)	0.0712 (9)
H9A	0.4582	0.5484	0.7697	0.107*

supplementary materials

H9B	0.6592	0.5465	0.6680	0.107*
H9C	0.3345	0.5145	0.6242	0.107*

Atomic displacement parameters (\AA^2)

	U^{11}	U^{22}	U^{33}	U^{12}	U^{13}	U^{23}
C1	0.0323 (12)	0.0467 (13)	0.0504 (14)	0.0078 (10)	0.0086 (10)	0.0101 (11)
N1	0.0608 (16)	0.0709 (16)	0.0496 (14)	0.0068 (13)	0.0017 (11)	0.0094 (11)
O1	0.113 (2)	0.131 (2)	0.0458 (13)	-0.0138 (16)	0.0034 (12)	0.0245 (13)
N2	0.0512 (14)	0.0510 (13)	0.0797 (17)	0.0095 (11)	0.0225 (13)	0.0255 (12)
C2	0.0424 (14)	0.0518 (15)	0.0425 (13)	0.0125 (11)	0.0041 (10)	0.0101 (11)
O2	0.0575 (13)	0.1052 (18)	0.0627 (13)	-0.0106 (12)	-0.0020 (10)	0.0039 (12)
N3	0.0292 (10)	0.0650 (13)	0.0446 (11)	0.0026 (9)	0.0098 (8)	0.0172 (9)
C3	0.0528 (15)	0.0544 (15)	0.0489 (14)	0.0174 (12)	0.0161 (12)	0.0207 (12)
O3	0.0941 (18)	0.1021 (18)	0.0914 (17)	-0.0079 (14)	0.0313 (14)	0.0522 (14)
C4	0.0396 (13)	0.0423 (13)	0.0586 (15)	0.0087 (10)	0.0153 (11)	0.0176 (11)
O4	0.0597 (13)	0.0651 (13)	0.0940 (16)	-0.0137 (10)	0.0088 (12)	0.0193 (11)
O5	0.0287 (9)	0.0861 (14)	0.0596 (11)	0.0045 (8)	0.0149 (8)	0.0164 (9)
C5	0.0330 (12)	0.0439 (13)	0.0489 (13)	0.0095 (10)	0.0090 (10)	0.0098 (11)
C6	0.0287 (11)	0.0446 (13)	0.0458 (13)	0.0100 (10)	0.0090 (9)	0.0114 (10)
C7	0.0270 (12)	0.0501 (14)	0.0468 (13)	0.0074 (10)	0.0115 (10)	0.0089 (10)
C8	0.0454 (14)	0.0633 (17)	0.0415 (14)	0.0050 (12)	0.0099 (11)	0.0125 (12)
C9	0.089 (2)	0.0671 (19)	0.0588 (18)	0.0082 (17)	0.0087 (16)	0.0185 (15)

Geometric parameters (\AA , $^\circ$)

C1—C2	1.376 (3)	C3—C4	1.380 (4)
C1—C6	1.394 (3)	C3—H3B	0.9300
C1—H1A	0.9300	C4—C5	1.375 (3)
N1—O1	1.212 (3)	O5—C7	1.232 (3)
N1—O2	1.224 (3)	C5—C6	1.392 (3)
N1—C2	1.479 (3)	C5—H5A	0.9300
N2—O4	1.214 (3)	C6—C7	1.498 (3)
N2—O3	1.214 (3)	C8—C9	1.494 (4)
N2—C4	1.477 (3)	C8—H8A	0.9700
C2—C3	1.379 (4)	C8—H8B	0.9700
N3—C7	1.327 (3)	C9—H9A	0.9600
N3—C8	1.454 (3)	C9—H9B	0.9600
N3—H3A	0.8600	C9—H9C	0.9600
C2—C1—C6	118.8 (2)	C4—C5—C6	119.0 (2)
C2—C1—H1A	120.6	C4—C5—H5A	120.5
C6—C1—H1A	120.6	C6—C5—H5A	120.5
O1—N1—O2	124.6 (2)	C5—C6—C1	119.7 (2)
O1—N1—C2	117.6 (2)	C5—C6—C7	123.0 (2)
O2—N1—C2	117.8 (2)	C1—C6—C7	117.3 (2)
O4—N2—O3	123.4 (2)	O5—C7—N3	124.0 (2)
O4—N2—C4	118.2 (2)	O5—C7—C6	119.2 (2)
O3—N2—C4	118.4 (3)	N3—C7—C6	116.78 (19)

C1—C2—C3	122.9 (2)	N3—C8—C9	112.1 (2)
C1—C2—N1	118.7 (2)	N3—C8—H8A	109.2
C3—C2—N1	118.5 (2)	C9—C8—H8A	109.2
C7—N3—C8	122.68 (19)	N3—C8—H8B	109.2
C7—N3—H3A	118.7	C9—C8—H8B	109.2
C8—N3—H3A	118.7	H8A—C8—H8B	107.9
C2—C3—C4	116.8 (2)	C8—C9—H9A	109.5
C2—C3—H3B	121.6	C8—C9—H9B	109.5
C4—C3—H3B	121.6	H9A—C9—H9B	109.5
C5—C4—C3	122.7 (2)	C8—C9—H9C	109.5
C5—C4—N2	118.9 (2)	H9A—C9—H9C	109.5
C3—C4—N2	118.3 (2)	H9B—C9—H9C	109.5
C6—C1—C2—C3	-1.3 (4)	C3—C4—C5—C6	-1.3 (4)
C6—C1—C2—N1	179.3 (2)	N2—C4—C5—C6	179.42 (19)
O1—N1—C2—C1	-170.8 (3)	C4—C5—C6—C1	0.9 (3)
O2—N1—C2—C1	8.0 (4)	C4—C5—C6—C7	-176.8 (2)
O1—N1—C2—C3	9.8 (4)	C2—C1—C6—C5	0.3 (3)
O2—N1—C2—C3	-171.4 (2)	C2—C1—C6—C7	178.2 (2)
C1—C2—C3—C4	0.9 (4)	C8—N3—C7—O5	3.0 (4)
N1—C2—C3—C4	-179.6 (2)	C8—N3—C7—C6	-177.4 (2)
C2—C3—C4—C5	0.4 (4)	C5—C6—C7—O5	147.8 (2)
C2—C3—C4—N2	179.7 (2)	C1—C6—C7—O5	-30.0 (3)
O4—N2—C4—C5	4.5 (3)	C5—C6—C7—N3	-31.9 (3)
O3—N2—C4—C5	-177.4 (2)	C1—C6—C7—N3	150.3 (2)
O4—N2—C4—C3	-174.8 (2)	C7—N3—C8—C9	90.0 (3)
O3—N2—C4—C3	3.2 (3)		

Hydrogen-bond geometry (Å, °)

D—H···A	D—H	H···A	D···A	D—H···A
N3—H3A···O5 ⁱ	0.86	2.13	2.886 (3)	146

Symmetry codes: (i) $x+1, y, z$.

supplementary materials

Fig. 1




Fig. 2

