

2-(4-Chlorophenyl)-1,5-diphenyl-3-tosyl-imidazolidin-4-one

S. Ranjith,^a A. Subbiah Pandi,^{a*} K. Namitharan^b and K. Pitchumani^b

^aDepartment of Physics, Presidency College (Autonomous), Chennai 600 005, India, and ^bSchool of Chemistry, Madurai Kamaraj University, Madurai 625 021, India

Correspondence e-mail: a_sp59@yahoo.in


Received 13 January 2011; accepted 5 March 2011

Key indicators: single-crystal X-ray study; $T = 293\text{ K}$; mean $\sigma(\text{C}-\text{C}) = 0.002\text{ \AA}$; R factor = 0.042; wR factor = 0.115; data-to-parameter ratio = 23.1.

In the title compound, $\text{C}_{28}\text{H}_{23}\text{ClN}_2\text{O}_3\text{S}$, the central imidazolidine ring adopts a twisted conformation and the S atom has distorted tetrahedral geometry. The crystal packing is stabilized by $\text{C}-\text{H}\cdots\text{O}$, $\text{C}-\text{H}\cdots\pi$ and $\pi-\pi$ interactions [centroid–centroid distance = $3.8302(10)\text{ \AA}$].

Related literature

For the biological activity of sulfonamides, see: Zareef *et al.* (2007); Chohan *et al.* (2007); Pomarnacka & Kozlarska-Kedra (2003); Nieto *et al.* (2005); Wang *et al.* (1995). For a related structure, see: Chakkaravarthi *et al.* (2008). For puckering parameters, see: Cremer & Pople (1975). For asymmetry parameters, see: Nardelli *et al.* (1983). For graph-set notation, see: Bernstein *et al.* (1995).



Experimental

Crystal data

$\text{C}_{28}\text{H}_{23}\text{ClN}_2\text{O}_3\text{S}$

$M_r = 503.00$

Monoclinic, $P2_1/n$

$a = 10.8458(3)\text{ \AA}$

$b = 13.0191(4)\text{ \AA}$

$c = 17.6720(5)\text{ \AA}$

$\beta = 103.757(2)^\circ$

$V = 2423.75(12)\text{ \AA}^3$

$Z = 4$

Mo $K\alpha$ radiation

$\mu = 0.28\text{ mm}^{-1}$

$T = 293\text{ K}$

$0.25 \times 0.22 \times 0.19\text{ mm}$

Data collection

Bruker APEXII CCD area-detector diffractometer
Absorption correction: multi-scan (*SADABS*; Sheldrick, 1996)
 $T_{\min} = 0.981$, $T_{\max} = 0.985$

32315 measured reflections
7313 independent reflections
5098 reflections with $I > 2\sigma(I)$
 $R_{\text{int}} = 0.030$

Refinement

$R[F^2 > 2\sigma(F^2)] = 0.042$
 $wR(F^2) = 0.115$
 $S = 1.05$
7313 reflections

317 parameters
H-atom parameters constrained
 $\Delta\rho_{\max} = 0.28\text{ e \AA}^{-3}$
 $\Delta\rho_{\min} = -0.35\text{ e \AA}^{-3}$

Table 1

Hydrogen-bond geometry (\AA , $^\circ$).

$Cg2$ and $Cg4$ are the centroids of the C2–C7 and C15–C20 rings, respectively.

$D-\text{H}\cdots A$	$D-\text{H}$	$\text{H}\cdots A$	$D\cdots A$	$D-\text{H}\cdots A$
$\text{C8}-\text{H8}\cdots\text{O3}^{\text{i}}$	0.98	2.44	3.3659 (17)	158
$\text{C14}-\text{H14}\cdots\text{O3}^{\text{i}}$	0.93	2.57	3.3855 (19)	146
$\text{C24}-\text{H24}\cdots\text{O2}^{\text{ii}}$	0.93	2.59	3.289 (2)	132
$\text{Cl}-\text{H1C}\cdots\text{Cg4}^{\text{iii}}$	0.96	2.90	3.484 (2)	120
$\text{C11}-\text{H11}\cdots\text{Cg2}^{\text{iii}}$	0.93	2.88	3.619 (17)	138

Symmetry codes: (i) $-x + \frac{3}{2}, y - \frac{1}{2}, -z + \frac{1}{2}$; (ii) $-x + \frac{3}{2}, y + \frac{1}{2}, -z + \frac{1}{2}$; (iii) $-x + 1, -y + 1, -z$.

Data collection: *APEX2* (Bruker, 2004); cell refinement: *SAINT* (Bruker, 2004); data reduction: *SAINT*; program(s) used to solve structure: *SHELXS97* (Sheldrick, 2008); program(s) used to refine structure: *SHELXL97* (Sheldrick, 2008); molecular graphics: *ORTEP-3* (Farrugia, 1997); software used to prepare material for publication: *SHELXL97* and *PLATON* (Spek, 2009).

SR and ASP thank Dr Babu Varghese, SAIF, IIT, Chennai, India, for the data collection.

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: PK2300).

References

- Bernstein, J., Davis, R. E., Shimoni, L. & Chang, N.-L. (1995). *Angew. Chem. Int. Ed. Engl.* **34**, 1555–1573.
- Bruker (2004). *APEX2* and *SAINT*. Bruker AXS Inc., Madison Wisconsin, USA.
- Chakkaravarthi, G., Dhayalan, V., Mohanakrishnan, A. K. & Manivannan, V. (2008). *Acta Cryst. E64*, o542.
- Chohan, Z. H. & Shah, H. A. (2007). *J. Enz. Inhib. Med. Chem.* **23**, 369–379.
- Cremer, D. & Pople, J. A. (1975). *J. Am. Chem. Soc.* **97**, 1354–1358.
- Farrugia, L. J. (1997). *J. Appl. Cryst.* **30**, 565.
- Nardelli, M. (1983). *Acta Cryst. C39*, 1141–1142.
- Nieto, M. J., Aloverio, F. L., Manzo, R. H. & Mazzieri, M. R. (2005). *Eur. J. Med. Chem.* **40**, 361–369.
- Pomarnacka, E. & Kozlarska-Kedra, I. (2003). *Farmaco*, **58**, 423–429.
- Sheldrick, G. M. (1996). *SADABS*. University of Göttingen, Germany.
- Sheldrick, G. M. (2008). *Acta Cryst. A64*, 112–122.
- Spek, A. L. (2009). *Acta Cryst. D65*, 148–155.
- Wang, W., Liang, T. C., Zheng, M. & Gao, X. (1995). *Tetrahedron Lett.* **36**, 1181–1184.
- Zareef, M., Iqbal, R., De Dominguez, N. G., Rodrigues, J., Zaidi, J. H., Arfan, M. & Supuran, C. T. (2007). *J. Enz. Inhib. Med. Chem.* **22**, 301–308.

supplementary materials

Acta Cryst. (2011). E67, o843 [doi:10.1107/S1600536811008427]

2-(4-Chlorophenyl)-1,5-diphenyl-3-tosylimidazolidin-4-one

S. Ranjith, A. SubbiahPandi, K. Namitharan and K. Pitchumani

Comment

Sulfonamides have widely been recognized for their wide variety of pharmacological activities such as antibacterial, anti-tumor, anti-carbonic anhydrase, diuretic, hypoglycaemic, antithyroid and protease inhibitory activity. Sulfonamides have also been used clinically as antimalarial agents (Zareef *et al.*, 2007), particularly sulfadiazine and sulfadoxine. Due to their significant pharmacology applications and widespread use in medicine, these compounds have also gained attention in bioinorganic and metal-based (Chohan *et al.*, 2007) drug chemistry. Sulfonamide derivatives are well known drugs and are used to control diseases caused by bacterial infections. Benzene sulfonamide derivatives are known to exhibit anticancer and HIV activities (Pomarnacka & Kozlarska-Kedra, 2003) and antibacterial activities (Nieto *et al.*, 2005). Imidazolidine compounds are important intermediates and building blocks in the construction of various biologically active compounds (Wang *et al.*, 1995). Against this background, and in order to obtain detailed information on molecular conformations in the solid state, an X-ray study of the title compound was carried out.

X-ray analysis confirms the molecular structure and atom connectivity as illustrated in Fig. 1. The geometry around the S atom is distorted tetrahedral, comprising two O atoms of the sulfonyl group, a C atom of a benzene ring and the imidazolidine N atom. The S–O, S–C, and S–N distances are 1.418 (1), 1.747 (1) and 1.677 (1) Å, respectively. These are comparable to those in similar structures (Chakkaravarthi *et al.*, 2008). The atom C11 deviates by 0.130 (1) Å from the least-squares plane of the ring C9–C14. The S atom exhibits significant deviation from that of a regular tetrahedron, with the largest deviations for the O–S–O [O1–S1–O2 120.9 (7)°] and O–S–N angles [O1–S1–N1 106.5 (6)°]. The widening of the angles may be due to repulsive interactions between the two short S=O bonds, similar to what is observed in related structures (Chakkaravarthi *et al.*, 2008). The chlorobenzene ring makes the dihedral angles of 39.4 (8), 85.1 (8) and 1.9 (9)° with respect to the C2–C7, C15–C20 and C23–C28 benzene rings.

The imidazolidine ring adopts a twisted conformation, with puckering parameters q_2 and ϕ (Cremer & Pople, 1975) and the smallest displacement asymmetric parameters, Δ , (Nardelli *et al.*, 1983) as follows: $q_2 = 0.1300$ (14) Å, $\phi = 21.9$ (6)°, $\Delta_s(C8) = 4.70$ (15). The intramolecular C6–H6···O2 hydrogen bond completes a five-membered ring, which generates an S(5) motif (Bernstein *et al.*, 1995). Atoms C8 and C14 act as donors to form bifurcated hydrogen bonds with atom O3 as an acceptor, results in the formation of $R^2_1(6)$ bifurcated ring. In addition to van der Waals interactions, the crystal packing is stabilized by C–H..O, C–H···π and π···π interactions (Table. 1).

Experimental

4-Toluenesulfonyl azide (1.3 mmol), phenylacetylene (1.2 mmol), 4-chlorophenyl *N*-phenylnitrone (1.0 mmol) and triethylamine (2 mmol) were successively added to Cu¹—Y zeolite (30 mg) in dichloromethane under N₂ atmosphere. After stirring at room temperature for 6 h, the mixture was diluted with dichloromethane. After removing the catalyst by filtration, followed by solvent evaporation, the resulting crude product was finally purified by column chromatography (silica gel).

supplementary materials

Single crystals suitable for X-ray diffraction were obtained by slow evaporation of a solution of the title compound in acetone at room temperature.

Refinement

All H atoms were fixed geometrically and allowed to ride on their parent C atoms, with C—H distances fixed in the range 0.93–0.97 Å with $U_{\text{iso}}(\text{H}) = 1.5U_{\text{eq}}(\text{C})$ for methyl H 1.2 $U_{\text{eq}}(\text{C})$ for other H atoms.

Figures




Fig. 1. The structure of the title compound showing the atom-numbering scheme. The displacement ellipsoids are drawn at the 30% probability level.




Fig. 2. π — π , C—H··· π interactions (dotted lines) in the title compound and also bifurcated hydrogen bonds formed by C8—H8···O3 and C14—H14···O3 at $3/2 - x, -1/2 + y, 1/2 - z$, results in the formation of $R^2_1(6)$ [O3,H8,C8,C9,C14,H14] bifurcated ring. Cg denotes ring centroid.[Symmetry code: (i) $2 - x, 1 - y, -z$; (ii) $1 - x, 1 - y, -z$; (iii) $3/2 - x, -1/2 + y, 1/2 - z$.]

2-(4-Chlorophenyl)-1,5-diphenyl-3-tosylimidazolidin-4-one

Crystal data

$C_{28}H_{23}ClN_2O_3S$	$F(000) = 1048$
$M_r = 503.00$	$D_x = 1.378 \text{ Mg m}^{-3}$
Monoclinic, $P2_1/n$	Mo $K\alpha$ radiation, $\lambda = 0.71073 \text{ \AA}$
Hall symbol: -P 2yn	Cell parameters from 7313 reflections
$a = 10.8458 (3) \text{ \AA}$	$\theta = 2.4\text{--}30.4^\circ$
$b = 13.0191 (4) \text{ \AA}$	$\mu = 0.28 \text{ mm}^{-1}$
$c = 17.6720 (5) \text{ \AA}$	$T = 293 \text{ K}$
$\beta = 103.757 (2)^\circ$	Block, colourless
$V = 2423.75 (12) \text{ \AA}^3$	$0.25 \times 0.22 \times 0.19 \text{ mm}$
$Z = 4$	

Data collection

Bruker APEXII CCD area-detector diffractometer	7313 independent reflections
Radiation source: fine-focus sealed tube graphite	5098 reflections with $I > 2\sigma(I)$
ω and ϕ scans	$R_{\text{int}} = 0.030$ $\theta_{\text{max}} = 30.4^\circ, \theta_{\text{min}} = 2.4^\circ$

Absorption correction: multi-scan
(*SADABS*; Sheldrick, 1996)
 $T_{\min} = 0.981$, $T_{\max} = 0.985$
32315 measured reflections

$h = -15 \rightarrow 15$
 $k = -17 \rightarrow 18$
 $l = -22 \rightarrow 25$

Refinement

Refinement on F^2
Least-squares matrix: full
 $R[F^2 > 2\sigma(F^2)] = 0.042$
 $wR(F^2) = 0.115$
 $S = 1.05$
7313 reflections
317 parameters
0 restraints

Primary atom site location: structure-invariant direct methods
Secondary atom site location: difference Fourier map
Hydrogen site location: inferred from neighbouring sites
H-atom parameters constrained
 $w = 1/\sigma^2(F_o^2) + (0.0495P)^2 + 0.5408P]$
where $P = (F_o^2 + 2F_c^2)/3$
 $(\Delta/\sigma)_{\max} = 0.001$
 $\Delta\rho_{\max} = 0.28 \text{ e } \text{\AA}^{-3}$
 $\Delta\rho_{\min} = -0.35 \text{ e } \text{\AA}^{-3}$

Special details

Geometry. All e.s.d.'s (except the e.s.d. in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell e.s.d.'s are taken into account individually in the estimation of e.s.d.'s in distances, angles and torsion angles; correlations between e.s.d.'s in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell e.s.d.'s is used for estimating e.s.d.'s involving l.s. planes.

Refinement. Refinement of F^2 against ALL reflections. The weighted R -factor wR and goodness of fit S are based on F^2 , conventional R -factors R are based on F , with F set to zero for negative F^2 . The threshold expression of $F^2 > \sigma(F^2)$ is used only for calculating R -factors(gt) etc. and is not relevant to the choice of reflections for refinement. R -factors based on F^2 are statistically about twice as large as those based on F , and R -factors based on ALL data will be even larger.

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (\AA^2)

	x	y	z	$U_{\text{iso}}^*/U_{\text{eq}}$
C1	0.8204 (2)	0.6211 (2)	-0.10561 (13)	0.0944 (10)
H1A	0.8293	0.5757	-0.1468	0.142*
H1B	0.8843	0.6735	-0.0987	0.142*
H1C	0.7378	0.6523	-0.1188	0.142*
C2	0.83550 (16)	0.5610 (2)	-0.03116 (10)	0.0615 (6)
C3	0.87311 (18)	0.60888 (16)	0.03995 (11)	0.0592 (5)
H3	0.8903	0.6789	0.0417	0.071*
C4	0.88600 (16)	0.55548 (13)	0.10884 (10)	0.0477 (4)
H4	0.9107	0.5892	0.1565	0.057*
C5	0.86169 (13)	0.45127 (12)	0.10598 (8)	0.0370 (3)
C6	0.82418 (16)	0.40127 (16)	0.03542 (10)	0.0524 (4)
H6	0.8079	0.3311	0.0334	0.063*
C7	0.81134 (17)	0.4579 (2)	-0.03242 (10)	0.0649 (6)
H7	0.7855	0.4248	-0.0802	0.078*

supplementary materials

C8	0.65428 (12)	0.31035 (10)	0.20980 (7)	0.0301 (3)
H8	0.6929	0.2445	0.2291	0.036*
C9	0.58939 (12)	0.30175 (10)	0.12399 (8)	0.0298 (3)
C10	0.51246 (13)	0.37975 (11)	0.08615 (8)	0.0359 (3)
H10	0.4942	0.4355	0.1145	0.043*
C11	0.46242 (14)	0.37599 (12)	0.00672 (9)	0.0396 (3)
H11	0.4102	0.4284	-0.0184	0.048*
C12	0.49087 (14)	0.29372 (12)	-0.03472 (8)	0.0398 (3)
C13	0.56268 (16)	0.21323 (13)	0.00174 (9)	0.0455 (4)
H13	0.5782	0.1566	-0.0268	0.055*
C14	0.61182 (14)	0.21732 (11)	0.08156 (9)	0.0389 (3)
H14	0.6602	0.1629	0.1068	0.047*
C15	0.46241 (13)	0.29610 (11)	0.26339 (8)	0.0336 (3)
C16	0.43733 (14)	0.19730 (12)	0.23351 (9)	0.0388 (3)
H16	0.4952	0.1644	0.2104	0.047*
C17	0.32626 (16)	0.14799 (14)	0.23816 (10)	0.0491 (4)
H17	0.3105	0.0818	0.2185	0.059*
C18	0.23885 (16)	0.19555 (16)	0.27141 (11)	0.0551 (5)
H18	0.1638	0.1623	0.2735	0.066*
C19	0.26360 (15)	0.29236 (15)	0.30142 (10)	0.0516 (4)
H19	0.2050	0.3246	0.3242	0.062*
C20	0.37404 (14)	0.34271 (13)	0.29837 (9)	0.0429 (3)
H20	0.3898	0.4081	0.3197	0.052*
C21	0.60120 (14)	0.45092 (11)	0.28719 (8)	0.0353 (3)
H21	0.5306	0.4961	0.2630	0.042*
C23	0.63258 (14)	0.46406 (11)	0.37521 (8)	0.0364 (3)
C28	0.70564 (19)	0.39176 (15)	0.42268 (10)	0.0578 (5)
H28	0.7302	0.3321	0.4013	0.069*
C27	0.7424 (2)	0.40811 (19)	0.50230 (11)	0.0731 (6)
H27	0.7914	0.3593	0.5344	0.088*
C26	0.7067 (2)	0.49587 (19)	0.53377 (11)	0.0694 (6)
H26	0.7324	0.5070	0.5872	0.083*
C25	0.6339 (2)	0.56698 (16)	0.48732 (11)	0.0628 (5)
H25	0.6089	0.6261	0.5092	0.075*
C24	0.59660 (18)	0.55168 (13)	0.40753 (10)	0.0491 (4)
H24	0.5472	0.6007	0.3759	0.059*
N1	0.75179 (11)	0.39178 (9)	0.22299 (6)	0.0318 (2)
N2	0.57393 (11)	0.34697 (10)	0.25969 (7)	0.0368 (3)
O1	0.98232 (10)	0.43180 (9)	0.25004 (6)	0.0459 (3)
O2	0.89588 (10)	0.27584 (9)	0.17703 (7)	0.0479 (3)
C22	0.71777 (14)	0.47825 (11)	0.25818 (8)	0.0350 (3)
S1	0.88693 (3)	0.38196 (3)	0.19302 (2)	0.03485 (10)
Cl1	0.43640 (5)	0.29404 (4)	-0.13549 (2)	0.06360 (15)
O3	0.77172 (11)	0.55951 (8)	0.26593 (7)	0.0480 (3)

Atomic displacement parameters (\AA^2)

U^{11}	U^{22}	U^{33}	U^{12}	U^{13}	U^{23}
----------	----------	----------	----------	----------	----------

C1	0.0684 (13)	0.165 (3)	0.0561 (13)	0.0392 (16)	0.0268 (11)	0.0549 (15)
C2	0.0390 (9)	0.1076 (17)	0.0414 (10)	0.0192 (10)	0.0163 (7)	0.0250 (10)
C3	0.0596 (11)	0.0658 (12)	0.0559 (11)	0.0111 (9)	0.0209 (9)	0.0236 (9)
C4	0.0540 (9)	0.0509 (9)	0.0390 (8)	-0.0006 (8)	0.0131 (7)	0.0051 (7)
C5	0.0329 (7)	0.0504 (9)	0.0292 (7)	-0.0002 (6)	0.0099 (5)	0.0016 (6)
C6	0.0496 (9)	0.0714 (12)	0.0382 (8)	-0.0063 (8)	0.0148 (7)	-0.0099 (8)
C7	0.0507 (10)	0.1166 (19)	0.0283 (8)	0.0025 (11)	0.0110 (7)	-0.0052 (10)
C8	0.0323 (6)	0.0316 (7)	0.0269 (6)	-0.0013 (5)	0.0078 (5)	-0.0011 (5)
C9	0.0304 (6)	0.0323 (7)	0.0273 (6)	-0.0012 (5)	0.0081 (5)	-0.0016 (5)
C10	0.0387 (7)	0.0351 (7)	0.0338 (7)	0.0041 (6)	0.0087 (6)	-0.0029 (6)
C11	0.0372 (7)	0.0431 (8)	0.0359 (7)	0.0017 (6)	0.0033 (6)	0.0054 (6)
C12	0.0383 (7)	0.0527 (9)	0.0267 (7)	-0.0074 (7)	0.0041 (6)	-0.0027 (6)
C13	0.0533 (9)	0.0458 (9)	0.0369 (8)	0.0005 (7)	0.0097 (7)	-0.0129 (7)
C14	0.0440 (8)	0.0370 (8)	0.0343 (7)	0.0054 (6)	0.0067 (6)	-0.0031 (6)
C15	0.0331 (7)	0.0423 (8)	0.0247 (6)	-0.0014 (6)	0.0057 (5)	0.0036 (5)
C16	0.0390 (7)	0.0424 (8)	0.0338 (7)	-0.0032 (6)	0.0067 (6)	0.0011 (6)
C17	0.0489 (9)	0.0514 (9)	0.0452 (9)	-0.0129 (8)	0.0076 (7)	0.0015 (7)
C18	0.0379 (8)	0.0745 (13)	0.0525 (10)	-0.0143 (8)	0.0098 (7)	0.0071 (9)
C19	0.0374 (8)	0.0731 (12)	0.0467 (9)	0.0013 (8)	0.0149 (7)	0.0025 (8)
C20	0.0408 (8)	0.0524 (9)	0.0372 (8)	0.0006 (7)	0.0123 (6)	-0.0024 (7)
C21	0.0404 (7)	0.0353 (7)	0.0313 (7)	0.0007 (6)	0.0106 (6)	-0.0028 (6)
C23	0.0420 (7)	0.0387 (7)	0.0314 (7)	-0.0048 (6)	0.0144 (6)	-0.0054 (6)
C28	0.0711 (12)	0.0630 (11)	0.0390 (9)	0.0181 (9)	0.0122 (8)	-0.0029 (8)
C27	0.0816 (14)	0.0961 (16)	0.0380 (10)	0.0180 (13)	0.0067 (10)	0.0061 (10)
C26	0.0871 (15)	0.0886 (15)	0.0351 (9)	-0.0171 (12)	0.0198 (10)	-0.0148 (10)
C25	0.0907 (14)	0.0578 (11)	0.0477 (10)	-0.0143 (11)	0.0320 (10)	-0.0208 (9)
C24	0.0671 (11)	0.0402 (8)	0.0462 (9)	-0.0046 (8)	0.0255 (8)	-0.0058 (7)
N1	0.0336 (6)	0.0333 (6)	0.0298 (6)	-0.0039 (5)	0.0101 (5)	-0.0039 (5)
N2	0.0402 (6)	0.0400 (6)	0.0341 (6)	-0.0072 (5)	0.0165 (5)	-0.0096 (5)
O1	0.0361 (5)	0.0592 (7)	0.0382 (6)	-0.0085 (5)	0.0003 (4)	0.0051 (5)
O2	0.0454 (6)	0.0413 (6)	0.0606 (7)	0.0060 (5)	0.0200 (6)	-0.0006 (5)
C22	0.0427 (7)	0.0357 (7)	0.0269 (6)	-0.0005 (6)	0.0088 (6)	-0.0004 (5)
S1	0.03133 (17)	0.0400 (2)	0.03336 (18)	0.00022 (14)	0.00796 (13)	0.00157 (14)
Cl1	0.0700 (3)	0.0859 (4)	0.02902 (19)	-0.0131 (3)	0.00031 (19)	-0.0035 (2)
O3	0.0615 (7)	0.0359 (6)	0.0511 (7)	-0.0115 (5)	0.0225 (6)	-0.0069 (5)

Geometric parameters (\AA , $^\circ$)

C1—C2	1.506 (3)	C15—N2	1.3941 (18)
C1—H1A	0.9600	C15—C20	1.397 (2)
C1—H1B	0.9600	C16—C17	1.385 (2)
C1—H1C	0.9600	C16—H16	0.9300
C2—C7	1.366 (3)	C17—C18	1.376 (3)
C2—C3	1.375 (3)	C17—H17	0.9300
C3—C4	1.380 (2)	C18—C19	1.369 (3)
C3—H3	0.9300	C18—H18	0.9300
C4—C5	1.381 (2)	C19—C20	1.378 (2)
C4—H4	0.9300	C19—H19	0.9300
C5—C6	1.380 (2)	C20—H20	0.9300

supplementary materials

C5—S1	1.7478 (15)	C21—N2	1.4446 (19)
C6—C7	1.386 (3)	C21—C22	1.5154 (19)
C6—H6	0.9300	C21—C23	1.5207 (19)
C7—H7	0.9300	C21—H21	0.9800
C8—N2	1.4592 (17)	C23—C24	1.373 (2)
C8—N1	1.4764 (17)	C23—C28	1.379 (2)
C8—C9	1.5165 (18)	C28—C27	1.385 (3)
C8—H8	0.9800	C28—H28	0.9300
C9—C10	1.3817 (19)	C27—C26	1.366 (3)
C9—C14	1.3844 (19)	C27—H27	0.9300
C10—C11	1.380 (2)	C26—C25	1.359 (3)
C10—H10	0.9300	C26—H26	0.9300
C11—C12	1.373 (2)	C25—C24	1.386 (2)
C11—H11	0.9300	C25—H25	0.9300
C12—C13	1.372 (2)	C24—H24	0.9300
C12—Cl1	1.7377 (15)	N1—C22	1.3782 (18)
C13—C14	1.385 (2)	N1—S1	1.6776 (11)
C13—H13	0.9300	O1—S1	1.4181 (11)
C14—H14	0.9300	O2—S1	1.4181 (12)
C15—C16	1.393 (2)	C22—O3	1.2008 (17)
C2—C1—H1A	109.5	C15—C16—H16	120.0
C2—C1—H1B	109.5	C18—C17—C16	120.95 (17)
H1A—C1—H1B	109.5	C18—C17—H17	119.5
C2—C1—H1C	109.5	C16—C17—H17	119.5
H1A—C1—H1C	109.5	C19—C18—C17	119.23 (16)
H1B—C1—H1C	109.5	C19—C18—H18	120.4
C7—C2—C3	118.31 (17)	C17—C18—H18	120.4
C7—C2—C1	121.0 (2)	C18—C19—C20	121.00 (16)
C3—C2—C1	120.7 (2)	C18—C19—H19	119.5
C2—C3—C4	121.58 (19)	C20—C19—H19	119.5
C2—C3—H3	119.2	C19—C20—C15	120.34 (16)
C4—C3—H3	119.2	C19—C20—H20	119.8
C3—C4—C5	118.97 (17)	C15—C20—H20	119.8
C3—C4—H4	120.5	N2—C21—C22	103.10 (11)
C5—C4—H4	120.5	N2—C21—C23	115.32 (12)
C6—C5—C4	120.63 (15)	C22—C21—C23	108.53 (11)
C6—C5—S1	120.18 (13)	N2—C21—H21	109.9
C4—C5—S1	119.11 (12)	C22—C21—H21	109.9
C5—C6—C7	118.55 (19)	C23—C21—H21	109.9
C5—C6—H6	120.7	C24—C23—C28	119.54 (15)
C7—C6—H6	120.7	C24—C23—C21	120.15 (14)
C2—C7—C6	121.95 (18)	C28—C23—C21	120.13 (13)
C2—C7—H7	119.0	C23—C28—C27	119.88 (17)
C6—C7—H7	119.0	C23—C28—H28	120.1
N2—C8—N1	100.27 (10)	C27—C28—H28	120.1
N2—C8—C9	115.24 (11)	C26—C27—C28	120.1 (2)
N1—C8—C9	110.82 (10)	C26—C27—H27	120.0
N2—C8—H8	110.0	C28—C27—H27	120.0
N1—C8—H8	110.0	C25—C26—C27	120.27 (18)

C9—C8—H8	110.0	C25—C26—H26	119.9
C10—C9—C14	119.01 (13)	C27—C26—H26	119.9
C10—C9—C8	120.87 (12)	C26—C25—C24	120.25 (18)
C14—C9—C8	120.04 (12)	C26—C25—H25	119.9
C11—C10—C9	120.82 (13)	C24—C25—H25	119.9
C11—C10—H10	119.6	C23—C24—C25	119.98 (18)
C9—C10—H10	119.6	C23—C24—H24	120.0
C12—C11—C10	119.08 (14)	C25—C24—H24	120.0
C12—C11—H11	120.5	C22—N1—C8	113.53 (11)
C10—C11—H11	120.5	C22—N1—S1	123.54 (9)
C13—C12—C11	121.32 (14)	C8—N1—S1	122.84 (9)
C13—C12—Cl1	120.00 (12)	C15—N2—C21	122.66 (12)
C11—C12—Cl1	118.67 (12)	C15—N2—C8	121.51 (11)
C12—C13—C14	119.14 (14)	C21—N2—C8	113.92 (11)
C12—C13—H13	120.4	O3—C22—N1	126.55 (13)
C14—C13—H13	120.4	O3—C22—C21	126.26 (13)
C9—C14—C13	120.49 (14)	N1—C22—C21	107.17 (11)
C9—C14—H14	119.8	O1—S1—O2	120.96 (7)
C13—C14—H14	119.8	O1—S1—N1	106.58 (6)
C16—C15—N2	120.96 (13)	O2—S1—N1	104.10 (6)
C16—C15—C20	118.45 (14)	O1—S1—C5	108.92 (7)
N2—C15—C20	120.58 (14)	O2—S1—C5	109.35 (7)
C17—C16—C15	120.01 (15)	N1—S1—C5	105.82 (6)
C17—C16—H16	120.0		
C7—C2—C3—C4	-0.3 (3)	C27—C26—C25—C24	-0.8 (3)
C1—C2—C3—C4	179.17 (17)	C28—C23—C24—C25	0.1 (3)
C2—C3—C4—C5	0.7 (3)	C21—C23—C24—C25	-174.97 (15)
C3—C4—C5—C6	-0.5 (2)	C26—C25—C24—C23	0.4 (3)
C3—C4—C5—S1	176.25 (13)	N2—C8—N1—C22	-14.57 (14)
C4—C5—C6—C7	-0.1 (2)	C9—C8—N1—C22	107.61 (13)
S1—C5—C6—C7	-176.75 (13)	N2—C8—N1—S1	168.83 (9)
C3—C2—C7—C6	-0.2 (3)	C9—C8—N1—S1	-69.00 (14)
C1—C2—C7—C6	-179.74 (17)	C16—C15—N2—C21	177.49 (13)
C5—C6—C7—C2	0.4 (3)	C20—C15—N2—C21	-3.5 (2)
N2—C8—C9—C10	45.41 (17)	C16—C15—N2—C8	14.2 (2)
N1—C8—C9—C10	-67.56 (15)	C20—C15—N2—C8	-166.84 (13)
N2—C8—C9—C14	-137.83 (13)	C22—C21—N2—C15	-170.26 (12)
N1—C8—C9—C14	109.21 (14)	C23—C21—N2—C15	71.64 (17)
C14—C9—C10—C11	-2.5 (2)	C22—C21—N2—C8	-5.78 (16)
C8—C9—C10—C11	174.27 (13)	C23—C21—N2—C8	-123.89 (13)
C9—C10—C11—C12	-0.5 (2)	N1—C8—N2—C15	176.67 (12)
C10—C11—C12—C13	3.2 (2)	C9—C8—N2—C15	57.67 (17)
C10—C11—C12—Cl1	-175.62 (11)	N1—C8—N2—C21	11.99 (15)
C11—C12—C13—C14	-2.8 (2)	C9—C8—N2—C21	-107.01 (14)
Cl1—C12—C13—C14	176.03 (12)	C8—N1—C22—O3	-169.50 (14)
C10—C9—C14—C13	2.9 (2)	S1—N1—C22—O3	7.1 (2)
C8—C9—C14—C13	-173.88 (13)	C8—N1—C22—C21	11.90 (15)
C12—C13—C14—C9	-0.3 (2)	S1—N1—C22—C21	-171.52 (9)
N2—C15—C16—C17	179.67 (14)	N2—C21—C22—O3	177.75 (14)

supplementary materials

C20—C15—C16—C17	0.7 (2)	C23—C21—C22—O3	−59.49 (19)
C15—C16—C17—C18	0.6 (2)	N2—C21—C22—N1	−3.64 (15)
C16—C17—C18—C19	−1.1 (3)	C23—C21—C22—N1	119.12 (13)
C17—C18—C19—C20	0.4 (3)	C22—N1—S1—O1	38.67 (13)
C18—C19—C20—C15	0.9 (3)	C8—N1—S1—O1	−145.07 (10)
C16—C15—C20—C19	−1.4 (2)	C22—N1—S1—O2	167.58 (11)
N2—C15—C20—C19	179.58 (14)	C8—N1—S1—O2	−16.15 (12)
N2—C21—C23—C24	−145.95 (14)	C22—N1—S1—C5	−77.18 (12)
C22—C21—C23—C24	99.02 (16)	C8—N1—S1—C5	99.08 (11)
N2—C21—C23—C28	39.0 (2)	C6—C5—S1—O1	149.38 (12)
C22—C21—C23—C28	−76.01 (18)	C4—C5—S1—O1	−27.35 (14)
C24—C23—C28—C27	−0.2 (3)	C6—C5—S1—O2	15.21 (14)
C21—C23—C28—C27	174.86 (17)	C4—C5—S1—O2	−161.53 (12)
C23—C28—C27—C26	−0.2 (3)	C6—C5—S1—N1	−96.37 (13)
C28—C27—C26—C25	0.7 (4)	C4—C5—S1—N1	86.89 (13)


Hydrogen-bond geometry (Å, °)

Cg2 and Cg4 are the centroids of the C2—C7 and C15—C20 rings, respectively.

<i>D</i> —H··· <i>A</i>	<i>D</i> —H	H··· <i>A</i>	<i>D</i> ··· <i>A</i>	<i>D</i> —H··· <i>A</i>
C6—H6···O2	0.93	2.59	2.934 (2)	102
C8—H8···O3 ⁱ	0.98	2.44	3.3659 (17)	158
C14—H14···O3 ⁱ	0.93	2.57	3.3855 (19)	146
C24—H24···O2 ⁱⁱ	0.93	2.59	3.289 (2)	132
C1—H1C···Cg4 ⁱⁱⁱ	0.96	2.90	3.484 (2)	120
C11—H11···Cg2 ⁱⁱⁱ	0.93	2.88	3.619 (17)	138

Symmetry codes: (i) $-x+3/2, y-1/2, -z+1/2$; (ii) $-x+3/2, y+1/2, -z+1/2$; (iii) $-x+1, -y+1, -z$.

Fig. 1



supplementary materials

Fig. 2

