

3-Methyl-5-oxo-4-(2-phenylhydrazinylidene)-4,5-dihydro-1H-pyrazole-1-carbothioamide

Hoong-Kun Fun,^{a,*} Safra Izuan Jama Asik,^a
Ibrahim Abdul Razak,^a Shobhitha Shetty^b and Balakrishna Kalluraya^b

^aX-ray Crystallography Unit, School of Physics, Universiti Sains Malaysia, 11800 USM, Penang, Malaysia, and ^bDepartment of Studies in Chemistry, Mangalore University, Mangalagangothri, Mangalore 574 199, India
Correspondence e-mail: hkfun@usm.my


Received 14 March 2011; accepted 15 March 2011

Key indicators: single-crystal X-ray study; $T = 296$ K; mean $\sigma(\text{C}-\text{C}) = 0.002$ Å; R factor = 0.049; wR factor = 0.135; data-to-parameter ratio = 26.8.

In the title compound, $\text{C}_{11}\text{H}_{11}\text{N}_5\text{OS}$, the pyrazole ring is approximately planar, with a maximum deviation of 0.010 (2) Å. The dihedral angles between the benzene ring and the pyrazole and carbothioamide groups are 5.42 (9) and 10.61 (18)°, respectively. An intramolecular $\text{N}-\text{H}\cdots\text{O}$ hydrogen bond generates an $S(6)$ ring motif. In the crystal, molecules are connected by intermolecular $\text{N}-\text{H}\cdots\text{O}$ and $\text{C}-\text{H}\cdots\text{S}$ hydrogen bonds, forming $R_2^2(12)$ ring motifs. In addition, there is a $\pi-\pi$ stacking interaction [centroid-centroid distance = 3.5188 (11) Å] between the pyrazole and benzene rings. These interactions link the molecules into infinite chains along [001].

Related literature

For general background to and applications of pyrazole derivatives, see: Rai & Kalluraya (2006); Rai *et al.* (2008); Sridhar & Perumal (2003). For graph-set theory, see: Bernstein *et al.* (1995). For bond-length data, see: Allen *et al.* (1987).



* Thomson Reuters ResearcherID: A-3561-2009.

Experimental

Crystal data

$\text{C}_{11}\text{H}_{11}\text{N}_5\text{OS}$
 $M_r = 261.31$
Monoclinic, $P2_1/c$
 $a = 7.7388$ (1) Å
 $b = 16.1103$ (3) Å
 $c = 11.3575$ (2) Å
 $\beta = 121.058$ (1)°
 $V = 1213.00$ (3) Å³
 $Z = 4$
Mo $K\alpha$ radiation
 $\mu = 0.26$ mm⁻¹
 $T = 296$ K
 $0.53 \times 0.39 \times 0.13$ mm

Data collection

Bruker SMART APEXII CCD area-detector diffractometer
Absorption correction: multi-scan (SADABS; Bruker, 2009)
 $T_{\min} = 0.874$, $T_{\max} = 0.967$
16258 measured reflections
4393 independent reflections
3037 reflections with $I > 2\sigma(I)$
 $R_{\text{int}} = 0.026$

Refinement

$R[F^2 > 2\sigma(F^2)] = 0.049$
 $wR(F^2) = 0.135$
 $S = 1.05$
4393 reflections
164 parameters
H-atom parameters constrained
 $\Delta\rho_{\text{max}} = 0.33$ e Å⁻³
 $\Delta\rho_{\text{min}} = -0.39$ e Å⁻³

Table 1

Hydrogen-bond geometry (Å, °).

$D-\text{H}\cdots A$	$D-\text{H}$	$\text{H}\cdots A$	$D\cdots A$	$D-\text{H}\cdots A$
$\text{N1}-\text{H1B}\cdots\text{O1}$	0.86	2.16	2.8147 (16)	132
$\text{N5}-\text{H5C}\cdots\text{O1}^{\text{i}}$	0.86	2.03	2.8806 (15)	172
$\text{C1}-\text{H1A}\cdots\text{S1}^{\text{ii}}$	0.93	2.80	3.6838 (16)	159

Symmetry codes: (i) $x, -y + \frac{1}{2}, z - \frac{1}{2}$; (ii) $x, -y + \frac{1}{2}, z + \frac{1}{2}$.

Data collection: APEX2 (Bruker, 2009); cell refinement: SAINT (Bruker, 2009); data reduction: SAINT; program(s) used to solve structure: SHELXTL (Sheldrick, 2008); program(s) used to refine structure: SHELXTL; molecular graphics: SHELXTL; software used to prepare material for publication: SHELXTL and PLATON (Spek, 2009).

HKF, SIJA and IAR thank Universiti Sains Malaysia for the Research University Grants (Nos. 1001/PFIZIK/811160 and 1001/PFIZIK/811151).

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: SJ5118).

References

- Allen, F. H., Kennard, O., Watson, D. G., Brammer, L., Orpen, A. G. & Taylor, R. (1987). *J. Chem. Soc. Perkin Trans. 2*, pp. S1–19.
Bernstein, J., Davis, R. E., Shimon, L. & Chang, N.-L. (1995). *Angew. Chem. Int. Ed. Engl.* **34**, 1555–1573.
Bruker (2009). APEX2, SAINT and SADABS. Bruker AXS Inc., Madison, Wisconsin, USA.
Rai, N. S. & Kalluraya, B. (2006). *Indian J. Chem. Sect. B*, **46**, 375–378.
Rai, N. S., Kalluraya, B., Lingappa, B., Shenoy, S. & Puranic, V. G. (2008). *Eur. J. Med. Chem.* **43**, 1715–1720.
Sheldrick, G. M. (2008). *Acta Cryst.* **A64**, 112–122.
Spek, A. L. (2009). *Acta Cryst.* **D65**, 148–155.
Sridhar, R. & Perumal, P. T. (2003). *Synth. Commun.* **33**, 1483–1488.

supplementary materials

Acta Cryst. (2011). E67, o928 [doi:10.1107/S1600536811009779]

3-Methyl-5-oxo-4-(2-phenylhydrazinylidene)-4,5-dihydro-1H-pyrazole-1-carbothioamide

H.-K. Fun, S. I. J. Asik, I. A. Razak, S. Shetty and B. Kalluraya

Comment

Pyrazole derivatives are in general well-known nitrogen-containing heterocyclic compounds and various procedures have been developed for their synthesis (Rai & Kalluraya, 2006). The chemistry of pyrazole derivatives has been the subject of much interest due to their importance for various applications and their widespread potential and proven biological and pharmacological activities (Rai *et al.*, 2008). Steroids containing a pyrazole moiety are of interest as psychopharmacological agents. Some alkyl- and aryl-substituted pyrazoles have a sharply pronounced sedative action on the central nervous system. Furthermore, certain alkyl pyrazoles show significant bacteriostatic, bacteriocidal, fungicidal, analgesic and anti-pyretic activities (Sridhar & Perumal, 2003).

Fig.1 shows the molecular structure of (I). The pyrazole (C7–C9/N3/N4) ring is approximately planar with a maximum deviation of 0.010 (2) Å for atom C8. The dihedral angle between benzene and pyrazole rings is 5.42 (9)°. The carbothioamide group (S1/C11/N5) is twisted at a dihedral angle 10.61 (18)° from the pyrazole ring. The bond lengths (Allen *et al.*, 1987) in (I) show normal values. An intramolecular N1—H1B···O1 hydrogen bond (Table 1) generates an S(6) ring motif (Bernstein *et al.*, 1995).

In the crystal packing of (I) (Fig. 2), molecules are connected by N5—H5C···O1 and C1—H1A···S1 intermolecular hydrogen bonds to form R²₂(12) ring motifs. These interactions also link the molecules into infinite one-dimensional chains along [0 0 1]. In addition, there is a π – π stacking interaction between pyrazole (C7–C9/N3/N4; centroid Cg1) and benzene (C1–C6; centroid Cg2) rings with a Cg1···Cg2 separation of 3.5188 (11) Å.

Experimental

To a solution of ethyl-3-oxo-2-(2-phenylhydrazinylidene) butanoate (0.01 mol) dissolved in glacial acetic acid (20 ml), a solution of thiosemicarbazide (0.02 mol) in glacial acetic acid (25 ml) was added and the mixture was refluxed for 4 h. It is cooled and allowed to stand overnight. The solid product that separated out was filtered and dried. It was then recrystallized from ethanol. Crystals suitable for X-ray analysis were obtained from 1:2 mixtures of DMF and ethanol by slow evaporation.

Refinement

All the H atoms were placed in calculated positions with N—H = 0.86 Å, C—H = 0.93 Å, and for C—H₃ = 0.96 Å. The U_{iso} values were constrained to be 1.5 U_{eq} of the carrier atom for methyl H atoms and 1.2 U_{eq} for the remaining H atoms. A rotating group model was used for the methyl groups.

Figures




Fig. 1. The structure of (I), showing 50% probability displacement ellipsoids and the atom-numbering scheme.




Fig. 2. The crystal packing of (I) viewed along the *a* axis, showing infinite one-dimensional chains along [0 0 1]. Hydrogen bonds are shown as dashed lines.

3-Methyl-5-oxo-4-(2-phenylhydrazinylidene)-4,5-dihydro-1H-pyrazole-1-carbothioamide

Crystal data

$C_{11}H_{11}N_5OS$

$M_r = 261.31$

Monoclinic, $P2_1/c$

Hall symbol: -P 2ybc

$a = 7.7388$ (1) Å

$b = 16.1103$ (3) Å

$c = 11.3575$ (2) Å

$\beta = 121.058$ (1)°

$V = 1213.00$ (3) Å³

$Z = 4$

$F(000) = 544$

$D_x = 1.431$ Mg m⁻³

Mo $K\alpha$ radiation, $\lambda = 0.71073$ Å

Cell parameters from 5770 reflections

$\theta = 2.5$ – 31.9 °

$\mu = 0.26$ mm⁻¹

$T = 296$ K

Plate, orange

$0.53 \times 0.39 \times 0.13$ mm

Data collection

Bruker SMART APEXII CCD area-detector diffractometer

Radiation source: sealed tube graphite

φ and ω scans

Absorption correction: multi-scan (SADABS; Bruker, 2009)

$T_{\min} = 0.874$, $T_{\max} = 0.967$

16258 measured reflections

4393 independent reflections

3037 reflections with $I > 2\sigma(I)$

$R_{\text{int}} = 0.026$

$\theta_{\max} = 32.6$ °, $\theta_{\min} = 2.5$ °

$h = -11 \rightarrow 11$

$k = -24 \rightarrow 23$

$l = -17 \rightarrow 16$

Refinement

Refinement on F^2

Least-squares matrix: full

Primary atom site location: structure-invariant direct methods

Secondary atom site location: difference Fourier map

$R[F^2 > 2\sigma(F^2)] = 0.049$	Hydrogen site location: inferred from neighbouring sites
$wR(F^2) = 0.135$	H-atom parameters constrained
$S = 1.05$	$w = 1/[\sigma^2(F_o^2) + (0.0587P)^2 + 0.2537P]$
4393 reflections	where $P = (F_o^2 + 2F_c^2)/3$
164 parameters	$(\Delta/\sigma)_{\max} < 0.001$
0 restraints	$\Delta\rho_{\max} = 0.33 \text{ e } \text{\AA}^{-3}$
	$\Delta\rho_{\min} = -0.39 \text{ e } \text{\AA}^{-3}$

Special details

Geometry. All esds (except the esd in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell esds are taken into account individually in the estimation of esds in distances, angles and torsion angles; correlations between esds in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell esds is used for estimating esds involving l.s. planes.

Refinement. Refinement of F^2 against ALL reflections. The weighted R-factor wR and goodness of fit S are based on F^2 , conventional R-factors R are based on F , with F set to zero for negative F^2 . The threshold expression of $F^2 > 2\sigma(F^2)$ is used only for calculating R-factors(gt) etc. and is not relevant to the choice of reflections for refinement. R-factors based on F^2 are statistically about twice as large as those based on F , and R-factors based on ALL data will be even larger.

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (\AA^2)

	<i>x</i>	<i>y</i>	<i>z</i>	$U_{\text{iso}}^*/U_{\text{eq}}$
S1	0.35472 (10)	0.29034 (2)	0.30227 (5)	0.07026 (19)
O1	0.36818 (17)	0.14963 (6)	0.49606 (10)	0.0434 (2)
N1	0.26541 (17)	0.00165 (7)	0.57723 (11)	0.0346 (2)
H1B	0.3036	0.0523	0.5989	0.041*
N2	0.23279 (17)	-0.02868 (7)	0.46071 (11)	0.0340 (2)
N3	0.25183 (18)	0.05536 (7)	0.18342 (11)	0.0340 (2)
N4	0.30783 (17)	0.12552 (6)	0.27173 (11)	0.0324 (2)
N5	0.3303 (2)	0.19488 (8)	0.10601 (13)	0.0478 (3)
H5B	0.3161	0.1470	0.0685	0.057*
H5C	0.3439	0.2384	0.0678	0.057*
C1	0.2876 (2)	-0.01706 (10)	0.79423 (14)	0.0413 (3)
H1A	0.3376	0.0367	0.8182	0.050*
C2	0.2625 (3)	-0.06542 (11)	0.88499 (15)	0.0494 (4)
H2A	0.2961	-0.0441	0.9703	0.059*
C3	0.1883 (3)	-0.14496 (11)	0.84978 (17)	0.0509 (4)
H3A	0.1732	-0.1775	0.9115	0.061*
C4	0.1363 (3)	-0.17618 (10)	0.72246 (19)	0.0523 (4)
H4A	0.0839	-0.2296	0.6983	0.063*
C5	0.1611 (2)	-0.12903 (9)	0.62989 (16)	0.0433 (3)
H5A	0.1272	-0.1505	0.5446	0.052*
C6	0.23757 (19)	-0.04923 (8)	0.66727 (13)	0.0337 (3)
C7	0.25637 (19)	0.01943 (7)	0.37735 (12)	0.0308 (2)
C8	0.31689 (19)	0.10634 (8)	0.39492 (12)	0.0313 (3)
C9	0.2218 (2)	-0.00556 (8)	0.24546 (13)	0.0328 (3)

supplementary materials

C10	0.1592 (3)	-0.08981 (9)	0.18412 (16)	0.0481 (4)
H10A	0.0995	-0.0862	0.0863	0.072*
H10B	0.0625	-0.1121	0.2046	0.072*
H10C	0.2751	-0.1255	0.2219	0.072*
C11	0.3313 (2)	0.20147 (8)	0.22208 (14)	0.0356 (3)

Atomic displacement parameters (\AA^2)

	U^{11}	U^{22}	U^{33}	U^{12}	U^{13}	U^{23}
S1	0.1363 (5)	0.0316 (2)	0.0759 (3)	-0.0167 (2)	0.0783 (4)	-0.01097 (18)
O1	0.0638 (7)	0.0371 (5)	0.0345 (5)	-0.0008 (4)	0.0291 (5)	-0.0053 (4)
N1	0.0439 (6)	0.0335 (5)	0.0316 (5)	0.0008 (4)	0.0233 (5)	0.0044 (4)
N2	0.0390 (6)	0.0347 (5)	0.0320 (5)	0.0033 (4)	0.0210 (5)	0.0043 (4)
N3	0.0454 (6)	0.0316 (5)	0.0304 (5)	-0.0027 (4)	0.0233 (5)	-0.0037 (4)
N4	0.0459 (6)	0.0276 (5)	0.0302 (5)	-0.0019 (4)	0.0242 (5)	-0.0015 (4)
N5	0.0785 (10)	0.0355 (6)	0.0431 (6)	-0.0039 (6)	0.0411 (7)	0.0037 (5)
C1	0.0478 (8)	0.0442 (7)	0.0362 (6)	-0.0027 (6)	0.0247 (6)	0.0024 (6)
C2	0.0543 (9)	0.0645 (10)	0.0351 (7)	0.0010 (7)	0.0271 (7)	0.0084 (7)
C3	0.0547 (9)	0.0569 (9)	0.0507 (8)	0.0075 (7)	0.0339 (7)	0.0219 (7)
C4	0.0645 (10)	0.0377 (7)	0.0662 (10)	0.0024 (7)	0.0420 (9)	0.0121 (7)
C5	0.0561 (9)	0.0370 (7)	0.0448 (8)	0.0019 (6)	0.0318 (7)	0.0042 (6)
C6	0.0345 (6)	0.0375 (6)	0.0330 (6)	0.0050 (5)	0.0202 (5)	0.0091 (5)
C7	0.0368 (6)	0.0298 (5)	0.0298 (5)	0.0014 (5)	0.0201 (5)	0.0018 (4)
C8	0.0381 (6)	0.0309 (5)	0.0297 (5)	0.0024 (5)	0.0208 (5)	0.0004 (4)
C9	0.0397 (7)	0.0310 (6)	0.0314 (6)	-0.0003 (5)	0.0210 (5)	-0.0015 (4)
C10	0.0700 (10)	0.0342 (7)	0.0471 (8)	-0.0090 (6)	0.0351 (8)	-0.0083 (6)
C11	0.0441 (7)	0.0307 (6)	0.0377 (6)	-0.0003 (5)	0.0252 (6)	0.0024 (5)

Geometric parameters (\AA , $^\circ$)

S1—C11	1.6560 (13)	C1—H1A	0.9300
O1—C8	1.2218 (15)	C2—C3	1.377 (3)
N1—N2	1.3073 (15)	C2—H2A	0.9300
N1—C6	1.4111 (16)	C3—C4	1.380 (3)
N1—H1B	0.8600	C3—H3A	0.9300
N2—C7	1.3072 (16)	C4—C5	1.389 (2)
N3—C9	1.2972 (17)	C4—H4A	0.9300
N3—N4	1.4222 (14)	C5—C6	1.387 (2)
N4—C11	1.3979 (16)	C5—H5A	0.9300
N4—C8	1.3988 (16)	C7—C9	1.4366 (17)
N5—C11	1.3185 (18)	C7—C8	1.4572 (17)
N5—H5B	0.8602	C9—C10	1.4880 (19)
N5—H5C	0.8600	C10—H10A	0.9600
C1—C2	1.383 (2)	C10—H10B	0.9600
C1—C6	1.3863 (19)	C10—H10C	0.9600
N2—N1—C6	119.64 (11)	C6—C5—H5A	120.7
N2—N1—H1B	120.2	C4—C5—H5A	120.7
C6—N1—H1B	120.2	C1—C6—C5	120.73 (13)

C7—N2—N1	119.04 (11)	C1—C6—N1	118.13 (12)
C9—N3—N4	107.04 (10)	C5—C6—N1	121.13 (12)
C11—N4—C8	130.31 (11)	N2—C7—C9	124.65 (12)
C11—N4—N3	117.83 (10)	N2—C7—C8	128.73 (12)
C8—N4—N3	111.70 (10)	C9—C7—C8	106.61 (10)
C11—N5—H5B	120.0	O1—C8—N4	129.69 (12)
C11—N5—H5C	120.0	O1—C8—C7	127.07 (12)
H5B—N5—H5C	120.0	N4—C8—C7	103.20 (10)
C2—C1—C6	119.56 (14)	N3—C9—C7	111.42 (11)
C2—C1—H1A	120.2	N3—C9—C10	122.83 (12)
C6—C1—H1A	120.2	C7—C9—C10	125.75 (12)
C3—C2—C1	120.38 (15)	C9—C10—H10A	109.5
C3—C2—H2A	119.8	C9—C10—H10B	109.5
C1—C2—H2A	119.8	H10A—C10—H10B	109.5
C2—C3—C4	119.73 (14)	C9—C10—H10C	109.5
C2—C3—H3A	120.1	H10A—C10—H10C	109.5
C4—C3—H3A	120.1	H10B—C10—H10C	109.5
C3—C4—C5	120.96 (16)	N5—C11—N4	113.45 (11)
C3—C4—H4A	119.5	N5—C11—S1	124.20 (10)
C5—C4—H4A	119.5	N4—C11—S1	122.34 (10)
C6—C5—C4	118.63 (15)		
C6—N1—N2—C7	179.04 (11)	C11—N4—C8—C7	173.49 (13)
C9—N3—N4—C11	-174.52 (12)	N3—N4—C8—C7	-1.73 (14)
C9—N3—N4—C8	1.36 (15)	N2—C7—C8—O1	4.4 (2)
C6—C1—C2—C3	-0.1 (2)	C9—C7—C8—O1	-176.61 (13)
C1—C2—C3—C4	-0.8 (3)	N2—C7—C8—N4	-177.54 (13)
C2—C3—C4—C5	1.2 (3)	C9—C7—C8—N4	1.46 (13)
C3—C4—C5—C6	-0.7 (2)	N4—N3—C9—C7	-0.33 (15)
C2—C1—C6—C5	0.6 (2)	N4—N3—C9—C10	179.58 (13)
C2—C1—C6—N1	-179.84 (13)	N2—C7—C9—N3	178.31 (12)
C4—C5—C6—C1	-0.2 (2)	C8—C7—C9—N3	-0.73 (15)
C4—C5—C6—N1	-179.78 (13)	N2—C7—C9—C10	-1.6 (2)
N2—N1—C6—C1	175.36 (12)	C8—C7—C9—C10	179.36 (14)
N2—N1—C6—C5	-5.05 (19)	C8—N4—C11—N5	174.38 (13)
N1—N2—C7—C9	-178.74 (12)	N3—N4—C11—N5	-10.64 (18)
N1—N2—C7—C8	0.1 (2)	C8—N4—C11—S1	-6.4 (2)
C11—N4—C8—O1	-8.5 (2)	N3—N4—C11—S1	168.56 (10)
N3—N4—C8—O1	176.27 (13)		

Hydrogen-bond geometry (Å, °)

<i>D</i> —H \cdots <i>A</i>	<i>D</i> —H	H \cdots <i>A</i>	<i>D</i> \cdots <i>A</i>	<i>D</i> —H \cdots <i>A</i>
N1—H1B \cdots O1	0.86	2.16	2.8147 (16)	132
N5—H5C \cdots O1 ⁱ	0.86	2.03	2.8806 (15)	172
C1—H1A \cdots S1 ⁱⁱ	0.93	2.80	3.6838 (16)	159

Symmetry codes: (i) *x*, $-y+1/2$, $z-1/2$; (ii) *x*, $-y+1/2$, $z+1/2$.

Fig. 1




Fig. 2

