

Diiiodido{2-(morpholin-4-yl)-N-[1-(2-pyridyl)ethylidene]ethanamine- $\kappa^3 N,N',N''$ }zinc

Nura Suleiman Gwaram, Hamid Khaledi* and Hapipah Mohd Ali

Department of Chemistry, University of Malaya, 50603 Kuala Lumpur, Malaysia
Correspondence e-mail: khaledi@siswa.um.edu.my


Received 14 April 2011; accepted 19 April 2011

Key indicators: single-crystal X-ray study; $T = 100\text{ K}$; mean $\sigma(\text{C}-\text{C}) = 0.004\text{ \AA}$; R factor = 0.018; wR factor = 0.045; data-to-parameter ratio = 19.3.

In the title compound, $[\text{ZnI}_2(\text{C}_{13}\text{H}_{19}\text{N}_3\text{O})]$, the Zn^{II} ion is five-coordinated in a distorted square-pyramidal geometry, in which the basal plane is defined by three N atoms from the Schiff base ligand and one iodide ion. A second iodide ligand, situated in the apical position, completes the coordination geometry. In the crystal structure, $\text{C}-\text{H}\cdots\text{O}$ hydrogen bonds link a pair of molecules around an inversion centre into a dimer.

Related literature

For the structure of an analogous ZnCl_2 complex, see: Ikmal Hisham *et al.* (2011). For square-pyramidal ZnI_2 complexes with N,N',N'' -tridentate ligands, see: Drew & Hollis (1978); Yousefi (2010). For a description of the geometry of complexes with five-coordinated metal ions, see: Addison *et al.* (1984).



Experimental

Crystal data

$[\text{ZnI}_2(\text{C}_{13}\text{H}_{19}\text{N}_3\text{O})]$

$M_r = 552.48$

Triclinic, $P\bar{1}$

$a = 8.8874(3)\text{ \AA}$

$b = 10.3117(4)\text{ \AA}$

$c = 10.3643(4)\text{ \AA}$

$\alpha = 68.8810(18)^\circ$

$\beta = 81.959(2)^\circ$

$\gamma = 66.3990(17)^\circ$

$V = 811.91(6)\text{ \AA}^3$

$Z = 2$

Mo $K\alpha$ radiation

$\mu = 5.31\text{ mm}^{-1}$

$T = 100\text{ K}$

$0.17 \times 0.13 \times 0.09\text{ mm}$

Data collection

Bruker APEXII CCD diffractometer

Absorption correction: multi-scan (*SADABS*; Sheldrick, 1996)

$T_{\min} = 0.465$, $T_{\max} = 0.646$

7292 measured reflections
3517 independent reflections
3260 reflections with $I > 2\sigma(I)$
 $R_{\text{int}} = 0.013$

Refinement

$R[F^2 > 2\sigma(F^2)] = 0.018$

$wR(F^2) = 0.045$

$S = 1.05$

3517 reflections

182 parameters
H-atom parameters constrained
 $\Delta\rho_{\max} = 0.85\text{ e \AA}^{-3}$
 $\Delta\rho_{\min} = -1.23\text{ e \AA}^{-3}$

Table 1
Selected bond lengths (\AA).

Zn1–N1	2.205 (2)	Zn1–I1	2.6018 (4)
Zn1–N2	2.093 (2)	Zn1–I2	2.6506 (4)
Zn1–N3	2.269 (2)		

Table 2
Hydrogen-bond geometry (\AA , $^\circ$).

$D-\text{H}\cdots A$	$D-\text{H}$	$\text{H}\cdots A$	$D\cdots A$	$D-\text{H}\cdots A$
C13–H13A \cdots O1 ⁱ	0.99	2.55	3.491 (3)	159

Symmetry code: (i) $-x + 1, -y + 2, -z + 2$.

Data collection: *APEX2* (Bruker, 2007); cell refinement: *SAINT* (Bruker, 2007); data reduction: *SAINT*; program(s) used to solve structure: *SHELXS97* (Sheldrick, 2008); program(s) used to refine structure: *SHELXL97* (Sheldrick, 2008); molecular graphics: *X-SEED* (Barbour, 2001); software used to prepare material for publication: *SHELXL97* and *publCIF* (Westrip, 2010).

The authors thank the University of Malaya for funding this study (FRGS grant No. FP004/2010B).

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: HY2422).

References

- Addison, A. W., Rao, T. N., Reedijk, J., Rijn, V. J. & Verschoor, G. C. (1984). *J. Chem. Soc. Dalton Trans.* pp. 1349–1356.
Barbour, L. J. (2001). *J. Supramol. Chem.* **1**, 189–191.
Bruker (2007). *APEX2* and *SAINT*. Bruker AXS Inc., Madison, Wisconsin, USA.
Drew, M. G. B. & Hollis, S. (1978). *J. Chem. Soc. Dalton Trans.* pp. 511–516.
Ikmal Hisham, N. A., Suleiman Gwaram, N., Khaledi, H. & Mohd Ali, H. (2011). *Acta Cryst. E67*, m55.
Sheldrick, G. M. (1996). *SADABS*. University of Göttingen, Germany.
Sheldrick, G. M. (2008). *Acta Cryst. A64*, 112–122.
Westrip, S. P. (2010). *J. Appl. Cryst.* **43**, 920–925.
Yousefi, M. (2010). *Acta Cryst. E66*, m1600–m1601.

supplementary materials

Acta Cryst. (2011). E67, m628 [doi:10.1107/S1600536811014656]

Diiodido{2-(morpholin-4-yl)-N-[1-(2-pyridyl)ethylidene]ethanamine- κ^3N,N',N'' }zinc

N. Suleiman Gwaram, H. Khaledi and H. Mohd Ali

Comment

The title compound (Fig. 1) was obtained *via* the complexation of Zn^{II} ion with the *in situ* prepared Schiff base, 2-morpholino-N-[1-(2-pyridyl)ethylidene]ethanamine, and two iodide ions. Similar to what was observed in an analogous ZnCl₂ complex (Ikmal Hisham *et al.*, 2011), the Schiff base acts as an *N,N',N''*-tridentate chelate ligand, along with two halide ligands, make a distorted square-pyramidal geometry around the metal ion ($\tau = 0.24$, Addison *et al.*, 1984). The Zn—I and Zn—N interatomic distances (Table 1) are comparable to the values reported for similar structures (Drew & Hollis, 1978; Yousefi, 2010). In the crystal, a pair of the molecules, related by a symmetry operation $-x + 1, -y + 2, -z + 2$, are linked through C—H···O hydrogen bonds into a centrosymmetric dimer (Table 2).

Experimental

A mixture of 2-acetylpyridine (0.20 g, 1.65 mmol) and 4-(2-aminoethyl)morpholine (0.21 g, 1.65 mmol) in ethanol (20 ml) was refluxed for 2 h, followed by addition of a solution of zinc(II) acetate dihydrate (0.36 g, 1.65 mmol) and potassium iodide (0.54 g, 3.3 mmol) in a minimum amount of water. The resulting solution was refluxed for 30 min and then left at room temperature. Brown crystals of the title complex were obtained in a few days.

Refinement

H atoms were placed at calculated positions and refined as riding atoms, with C—H = 0.95 (aryl), 0.98 (methyl) and 0.99 (methylene) Å and with $U_{\text{iso}}(\text{H}) = 1.2(1.5 \text{ for methyl})U_{\text{eq}}(\text{C})$.

Figures




Fig. 1. Molecular structure of the title compound. Displacement ellipsoids are drawn at the 50% probability level.

Diiodido{2-(morpholin-4-yl)-N-[1-(2-pyridyl)ethylidene]ethanamine- κ^3N,N',N'' }zinc

Crystal data

[ZnI ₂ (C ₁₃ H ₁₉ N ₃ O)]	Z = 2
$M_r = 552.48$	$F(000) = 524$
Triclinic, $P\bar{1}$	$D_x = 2.260 \text{ Mg m}^{-3}$

supplementary materials

Hall symbol: -P 1

$a = 8.8874(3)$ Å

$b = 10.3117(4)$ Å

$c = 10.3643(4)$ Å

$\alpha = 68.8810(18)^\circ$

$\beta = 81.959(2)^\circ$

$\gamma = 66.3990(17)^\circ$

$V = 811.91(6)$ Å³

Mo $K\alpha$ radiation, $\lambda = 0.71073$ Å

Cell parameters from 6030 reflections

$\theta = 2.3\text{--}30.4^\circ$

$\mu = 5.31$ mm⁻¹

$T = 100$ K

Block, brown

$0.17 \times 0.13 \times 0.09$ mm

Data collection

Bruker APEXII CCD diffractometer

3517 independent reflections

Radiation source: fine-focus sealed tube graphite

3260 reflections with $I > 2\sigma(I)$

φ and ω scans

$R_{\text{int}} = 0.013$

Absorption correction: multi-scan (*SADABS*; Sheldrick, 1996)

$\theta_{\text{max}} = 27.0^\circ$, $\theta_{\text{min}} = 2.1^\circ$

$T_{\text{min}} = 0.465$, $T_{\text{max}} = 0.646$

$h = -11\text{--}11$

7292 measured reflections

$k = -13\text{--}13$

$l = -13\text{--}13$

Refinement

Refinement on F^2

Primary atom site location: structure-invariant direct methods

Least-squares matrix: full

Secondary atom site location: difference Fourier map

$R[F^2 > 2\sigma(F^2)] = 0.018$

Hydrogen site location: inferred from neighbouring sites

$wR(F^2) = 0.045$

H-atom parameters constrained

$S = 1.05$

$w = 1/[\sigma^2(F_o^2) + (0.0197P)^2 + 1.3278P]$

where $P = (F_o^2 + 2F_c^2)/3$

3517 reflections

$(\Delta/\sigma)_{\text{max}} = 0.002$

182 parameters

$\Delta\rho_{\text{max}} = 0.85$ e Å⁻³

0 restraints

$\Delta\rho_{\text{min}} = -1.23$ e Å⁻³

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (Å²)

	x	y	z	$U_{\text{iso}}^*/U_{\text{eq}}$
Zn1	0.77476(4)	0.81170(3)	0.66161(3)	0.01267(7)
I1	0.64389(2)	1.103488(18)	0.584395(17)	0.01508(5)
I2	1.05998(2)	0.694938(19)	0.791314(17)	0.01566(5)
O1	0.3717(2)	0.8562(2)	1.0506(2)	0.0212(4)
N1	0.8699(3)	0.8016(2)	0.4558(2)	0.0135(4)
N2	0.7235(3)	0.6403(2)	0.6380(2)	0.0137(4)
N3	0.6122(3)	0.7619(2)	0.8469(2)	0.0137(4)
C1	0.9527(3)	0.8810(3)	0.3702(3)	0.0161(5)
H1	0.9697	0.9537	0.3959	0.019*
C2	1.0153(3)	0.8615(3)	0.2446(3)	0.0181(5)

H2	1.0767	0.9176	0.1868	0.022*
C3	0.9859 (3)	0.7587 (3)	0.2062 (3)	0.0190 (6)
H3	1.0239	0.7455	0.1197	0.023*
C4	0.9002 (3)	0.6744 (3)	0.2951 (3)	0.0165 (5)
H4	0.8797	0.6024	0.2709	0.020*
C5	0.8456 (3)	0.6981 (3)	0.4199 (3)	0.0145 (5)
C6	0.7574 (3)	0.6108 (3)	0.5253 (3)	0.0139 (5)
C7	0.7137 (3)	0.5006 (3)	0.4920 (3)	0.0181 (5)
H7A	0.6732	0.4412	0.5754	0.027*
H7B	0.8112	0.4336	0.4582	0.027*
H7C	0.6280	0.5547	0.4204	0.027*
C8	0.6366 (3)	0.5663 (3)	0.7513 (3)	0.0161 (5)
H8A	0.6837	0.4566	0.7693	0.019*
H8B	0.5189	0.6052	0.7275	0.019*
C9	0.6557 (3)	0.5993 (3)	0.8785 (3)	0.0150 (5)
H9A	0.5839	0.5643	0.9531	0.018*
H9B	0.7707	0.5435	0.9119	0.018*
C10	0.4339 (3)	0.8512 (3)	0.8159 (3)	0.0157 (5)
H10A	0.4008	0.8254	0.7440	0.019*
H10B	0.4152	0.9594	0.7782	0.019*
C11	0.3273 (3)	0.8231 (3)	0.9429 (3)	0.0177 (5)
H11A	0.2107	0.8864	0.9174	0.021*
H11B	0.3391	0.7166	0.9766	0.021*
C12	0.5400 (3)	0.7684 (3)	1.0885 (3)	0.0188 (6)
H12A	0.5561	0.6608	1.1276	0.023*
H12B	0.5690	0.7960	1.1610	0.023*
C13	0.6531 (3)	0.7918 (3)	0.9652 (3)	0.0168 (5)
H13A	0.6466	0.8967	0.9338	0.020*
H13B	0.7676	0.7247	0.9950	0.020*

Atomic displacement parameters (\AA^2)

	U^{11}	U^{22}	U^{33}	U^{12}	U^{13}	U^{23}
Zn1	0.01377 (14)	0.01207 (14)	0.01168 (14)	-0.00432 (11)	0.00086 (11)	-0.00440 (11)
I1	0.01607 (9)	0.01218 (8)	0.01448 (9)	-0.00349 (6)	0.00078 (6)	-0.00401 (6)
I2	0.01406 (9)	0.01676 (9)	0.01511 (9)	-0.00421 (7)	-0.00121 (6)	-0.00553 (7)
O1	0.0184 (10)	0.0264 (11)	0.0200 (10)	-0.0062 (8)	0.0039 (8)	-0.0130 (9)
N1	0.0131 (10)	0.0134 (10)	0.0120 (10)	-0.0034 (8)	0.0003 (8)	-0.0039 (8)
N2	0.0130 (10)	0.0110 (10)	0.0147 (11)	-0.0028 (8)	-0.0011 (8)	-0.0033 (8)
N3	0.0150 (11)	0.0130 (10)	0.0116 (10)	-0.0047 (9)	0.0001 (8)	-0.0032 (8)
C1	0.0148 (12)	0.0163 (12)	0.0164 (13)	-0.0051 (10)	-0.0007 (10)	-0.0051 (10)
C2	0.0142 (12)	0.0194 (13)	0.0178 (13)	-0.0053 (11)	0.0010 (10)	-0.0047 (11)
C3	0.0171 (13)	0.0214 (14)	0.0138 (13)	-0.0024 (11)	0.0017 (10)	-0.0067 (11)
C4	0.0153 (13)	0.0178 (13)	0.0160 (13)	-0.0034 (10)	-0.0007 (10)	-0.0084 (11)
C5	0.0108 (12)	0.0140 (12)	0.0160 (13)	-0.0014 (10)	-0.0011 (9)	-0.0053 (10)
C6	0.0116 (12)	0.0123 (12)	0.0159 (13)	-0.0012 (10)	-0.0012 (9)	-0.0057 (10)
C7	0.0187 (13)	0.0189 (13)	0.0185 (14)	-0.0070 (11)	0.0010 (10)	-0.0085 (11)
C8	0.0176 (13)	0.0155 (12)	0.0165 (13)	-0.0077 (10)	0.0038 (10)	-0.0064 (10)

supplementary materials

C9	0.0167 (13)	0.0123 (12)	0.0135 (13)	-0.0048 (10)	0.0009 (10)	-0.0028 (10)
C10	0.0145 (12)	0.0169 (12)	0.0148 (13)	-0.0042 (10)	0.0003 (10)	-0.0061 (10)
C11	0.0151 (13)	0.0192 (13)	0.0186 (14)	-0.0055 (11)	0.0020 (10)	-0.0079 (11)
C12	0.0183 (13)	0.0256 (14)	0.0123 (13)	-0.0071 (11)	0.0008 (10)	-0.0076 (11)
C13	0.0181 (13)	0.0197 (13)	0.0124 (13)	-0.0064 (11)	0.0018 (10)	-0.0067 (10)

Geometric parameters (\AA , $^{\circ}$)

Zn1—N1	2.205 (2)	C4—H4	0.9500
Zn1—N2	2.093 (2)	C5—C6	1.495 (4)
Zn1—N3	2.269 (2)	C6—C7	1.495 (4)
Zn1—I1	2.6018 (4)	C7—H7A	0.9800
Zn1—I2	2.6506 (4)	C7—H7B	0.9800
O1—C11	1.423 (3)	C7—H7C	0.9800
O1—C12	1.426 (3)	C8—C9	1.521 (4)
N1—C1	1.332 (3)	C8—H8A	0.9900
N1—C5	1.349 (3)	C8—H8B	0.9900
N2—C6	1.277 (3)	C9—H9A	0.9900
N2—C8	1.459 (3)	C9—H9B	0.9900
N3—C9	1.479 (3)	C10—C11	1.521 (4)
N3—C13	1.490 (3)	C10—H10A	0.9900
N3—C10	1.491 (3)	C10—H10B	0.9900
C1—C2	1.394 (4)	C11—H11A	0.9900
C1—H1	0.9500	C11—H11B	0.9900
C2—C3	1.380 (4)	C12—C13	1.521 (4)
C2—H2	0.9500	C12—H12A	0.9900
C3—C4	1.394 (4)	C12—H12B	0.9900
C3—H3	0.9500	C13—H13A	0.9900
C4—C5	1.387 (4)	C13—H13B	0.9900
N2—Zn1—N1	74.71 (8)	C6—C7—H7B	109.5
N2—Zn1—N3	78.31 (8)	H7A—C7—H7B	109.5
N1—Zn1—N3	151.54 (8)	C6—C7—H7C	109.5
N2—Zn1—I1	137.01 (6)	H7A—C7—H7C	109.5
N1—Zn1—I1	95.53 (6)	H7B—C7—H7C	109.5
N3—Zn1—I1	98.03 (6)	N2—C8—C9	107.5 (2)
N2—Zn1—I2	109.52 (6)	N2—C8—H8A	110.2
N1—Zn1—I2	98.21 (6)	C9—C8—H8A	110.2
N3—Zn1—I2	99.18 (6)	N2—C8—H8B	110.2
I1—Zn1—I2	113.325 (12)	C9—C8—H8B	110.2
C11—O1—C12	110.8 (2)	H8A—C8—H8B	108.5
C1—N1—C5	118.9 (2)	N3—C9—C8	111.1 (2)
C1—N1—Zn1	126.70 (18)	N3—C9—H9A	109.4
C5—N1—Zn1	114.34 (17)	C8—C9—H9A	109.4
C6—N2—C8	122.6 (2)	N3—C9—H9B	109.4
C6—N2—Zn1	120.65 (18)	C8—C9—H9B	109.4
C8—N2—Zn1	116.61 (16)	H9A—C9—H9B	108.0
C9—N3—C13	111.1 (2)	N3—C10—C11	112.5 (2)
C9—N3—C10	112.5 (2)	N3—C10—H10A	109.1
C13—N3—C10	107.4 (2)	C11—C10—H10A	109.1

C9—N3—Zn1	100.98 (15)	N3—C10—H10B	109.1
C13—N3—Zn1	111.80 (16)	C11—C10—H10B	109.1
C10—N3—Zn1	113.10 (15)	H10A—C10—H10B	107.8
N1—C1—C2	122.6 (3)	O1—C11—C10	111.4 (2)
N1—C1—H1	118.7	O1—C11—H11A	109.3
C2—C1—H1	118.7	C10—C11—H11A	109.3
C3—C2—C1	118.3 (3)	O1—C11—H11B	109.3
C3—C2—H2	120.8	C10—C11—H11B	109.3
C1—C2—H2	120.8	H11A—C11—H11B	108.0
C2—C3—C4	119.6 (3)	O1—C12—C13	111.7 (2)
C2—C3—H3	120.2	O1—C12—H12A	109.3
C4—C3—H3	120.2	C13—C12—H12A	109.3
C5—C4—C3	118.4 (3)	O1—C12—H12B	109.3
C5—C4—H4	120.8	C13—C12—H12B	109.3
C3—C4—H4	120.8	H12A—C12—H12B	107.9
N1—C5—C4	122.1 (2)	N3—C13—C12	113.3 (2)
N1—C5—C6	114.8 (2)	N3—C13—H13A	108.9
C4—C5—C6	123.1 (2)	C12—C13—H13A	108.9
N2—C6—C5	115.4 (2)	N3—C13—H13B	108.9
N2—C6—C7	125.4 (2)	C12—C13—H13B	108.9
C5—C6—C7	119.2 (2)	H13A—C13—H13B	107.7
C6—C7—H7A	109.5		

Hydrogen-bond geometry (\AA , $^\circ$)

$D\text{—H}\cdots A$	$D\text{—H}$	$H\cdots A$	$D\cdots A$	$D\text{—H}\cdots A$
C13—H13A \cdots O1 ⁱ	0.99	2.55	3.491 (3)	159

Symmetry codes: (i) $-x+1, -y+2, -z+2$.

supplementary materials

Fig. 1

