

## Dicarbonyl( $\eta^5$ -cyclopentadienyl)- [2-(phenylsulfanyl)ethyl]iron(II)

Dennis S. Mkhize, Vincent O. Nyamori and  
Muhammad D. Bala\*

School of Chemistry, University of KwaZulu-Natal, Westville Campus, Private Bag X54001, Durban 4000, South Africa

Correspondence e-mail: bala@ukzn.ac.za


Received 11 April 2011; accepted 21 April 2011

Key indicators: single-crystal X-ray study;  $T = 173$  K; mean  $\sigma(\text{C}-\text{C}) = 0.010$  Å;  $R$  factor = 0.075;  $wR$  factor = 0.153; data-to-parameter ratio = 16.0.

The title compound,  $[\text{Fe}(\text{C}_5\text{H}_5)(\text{C}_8\text{H}_9\text{S})(\text{CO})_2]$ , is a three-legged piano-stool iron(II) complex that is characterized by a thioethyl-linked phenyl ring and a cyclopentadienyl moiety that occupies the apical coordination site. The two aromatic rings are essentially planar with the same maximum deviation of 0.009 Å. The mean planes of the phenyl and cyclopentadienyl rings bisect at an acute angle of 50.08°.

### Related literature

For general background and related synthesis, see: King & Bisnette (1965a,b); Theys *et al.* (2009); Nyamori *et al.* (2008). For related structures, see: O'Connor *et al.* (1987).



### Experimental

#### Crystal data

$[\text{Fe}(\text{C}_5\text{H}_5)(\text{C}_8\text{H}_9\text{S})(\text{CO})_2]$	$V = 2814.0$ (2) Å <sup>3</sup>
$M_r = 314.17$	$Z = 8$
Orthorhombic, $Pbca$	Mo $K\alpha$ radiation
$a = 10.3992$ (5) Å	$\mu = 1.21$ mm <sup>-1</sup>
$b = 7.6402$ (4) Å	$T = 173$ K
$c = 35.4173$ (16) Å	$0.57 \times 0.27 \times 0.08$ mm

#### Data collection

Bruker APEXII CCD area-detector diffractometer	20490 measured reflections
Absorption correction: integration ( <i>SADABS</i> ; Bruker, 2009)	2752 independent reflections
$(SADABS$ ; Bruker, 2009)	2339 reflections with $I > 2\sigma(I)$
$T_{\min} = 0.545$ , $T_{\max} = 0.909$	$R_{\text{int}} = 0.068$

#### Refinement

$R[F^2 > 2\sigma(F^2)] = 0.075$	172 parameters
$wR(F^2) = 0.153$	H-atom parameters constrained
$S = 1.26$	$\Delta\rho_{\max} = 0.49$ e Å <sup>-3</sup>
2752 reflections	$\Delta\rho_{\min} = -0.75$ e Å <sup>-3</sup>

Data collection: *APEX2* (Bruker, 2009); cell refinement: *SAINT* (Bruker, 2009; data reduction: *SAINT*; program(s) used to solve structure: *SHELXS97* (Sheldrick, 2008); program(s) used to refine structure: *SHELXL97* (Sheldrick, 2008); molecular graphics: *ORTEP-3* (Farrugia, 1997); software used to prepare material for publication: *SHELXTL* (Sheldrick, 2008).

The authors thank Dr Manuel Fernandes for data collection, the NRF and the University of KwaZulu-Natal for financial support.

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: DN2676).

### References

- Bruker (2009). *APEX2, SADABS and SAINT*. Bruker AXS Inc., Madison, Wisconsin, USA.  
Farrugia, L. J. (1997). *J. Appl. Cryst.* **30**, 565.  
King, R. B. & Bisnette, M. B. (1965a). *Inorg. Chem.* **4**, 482–485.  
King, R. B. & Bisnette, M. B. (1965b). *Inorg. Chem.* **4**, 486–493.  
Nyamori, V. O., Mhlanga, S. D. & Coville, N. J. (2008). *J. Organomet. Chem.* **693**, 2205–2222.  
O'Connor, E. J., Brandt, S. & Helquist, P. (1987). *J. Am. Chem. Soc.* **109**, 3739–3747.  
Sheldrick, G. M. (2008). *Acta Cryst. A* **64**, 112–122.  
Theys, R. D., Dudely, M. E. & Hossain, M. M. (2009). *Coord. Chem. Rev.* **293**, 180–234.

## **supplementary materials**

*Acta Cryst.* (2011). E67, m644 [doi:10.1107/S160053681101511X]

## Dicarbonyl( $\eta^5$ -cyclopentadienyl)[2-(phenylsulfanyl)ethyl]iron(II)

**D. S. Mkhize, V. O. Nyamori and M. D. Bala**

### Comment

Organosulfur compounds of iron are valuable compounds that are used for the synthesis of organometallic catalysts and catalysts precursor salts. Amongst many applications this salts have been used for the cyclopropanation of alkenes (O'Connor *et al.*, 1987). The recent proliferation and interest in the chemistry of materials has also rekindled interest in Fe-based organometallics as catalysts for the synthesis of carbon nanotubes and nanomaterials in general (Nyamori *et al.*, 2008).

We are currently studying the application of (I) as a precursor catalyst in the synthesis of carbon nanomaterials. The presence of sulfur in this compound was believed to play the role of increasing the yield and affecting the morphology of the products obtained. In the title compound (I), the coordination of ligands around the Fe(II) ion is described as three-legged piano stool distorted octahedral geometry (Fig. 1). This is a typical geometry for complexes based on the dicarbonyl( $\eta^5$ -cyclopentadienyl)Fe(II) moiety (generally abbreviated as the Fp anion) and many compounds of iron with this geometry are well documented (Theye, *et al.*, 2009; O'Connor *et al.*, 1987).

### Experimental

A solution of cyclopentadienylirondicarbonyl dimer (1.501 g, 0.424 mmol) in dry tetrahydrofuran was treated with Na/Hg amalgam [sodium (0.2902 g, 12.70 mmol) and mercury (2.8 ml, 18.75 mmol)] and stirred at room temperature for two hours. The resulting yellowish-brown solution was transferred into another flask *via* a cannula and reacted with 2-chloroethyl phenyl sulfide (0.729 g, 4.242 mmol). The reaction mixture was allowed to stir for 16 h at room temperature. After concentrating the greyish brown mixture *in vacuo* it was extracted with dichloromethane ( $3 \times 30$  ml). Filtration of the extracted product through a celite pad afforded a clear brownish-orange filtrate which was further reduced *in vacuo*. The brown oil that resulted was purified by flash column chromatography on silica using hexane as mobile phase. A dark yellow band was collected and removal of volatiles afforded 0.8183 g of the final product as a yellow crystalline solid (yield, 61%).

$^1\text{H}$  NMR  $\delta_{\text{H}}$  (400 MHz,  $\text{CDCl}_3$ , p.p.m.): 1.58 (2*H*, t, *J* 8.56,  $\text{CH}_2$ ), 3.02 (2*H*, t, *J* 8.40,  $\text{CH}_2$ ), 4.68 (5*H*, s, Cp), 7.08 (1*H*, t, ArH), 7.20 (2*H*, m, *J* 16.4, ArH<sub>2</sub>), 7.26 (2*H*, m, *J* 16.5, ArH<sub>2</sub>).

$^{13}\text{C}$  NMR (400 MHz,  $\text{CDCl}_3$ , p.p.m.): 40.72, 84.03, 124.3, 127.6, 127.8, 136.1, 215.6.

IR ( $\nu_{\text{CO}}$ ,  $\text{cm}^{-1}$ ): 1993, 1937, 1708.

### Refinement

All H-atoms were refined using a riding model, with C—H = 0.93 Å and  $U_{\text{iso}}(\text{H}) = 1.2U_{\text{eq}}(\text{C})$  for aromatic and C—H = 0.97 Å and  $U_{\text{iso}}(\text{H}) = 1.2U_{\text{eq}}(\text{C})$  for  $\text{CH}_2$ .

# supplementary materials

---

## Figures




Fig. 1. Molecular structure of the title compound with the atom numbering scheme. Displacement ellipsoids are drawn at the 50% probability level. H atoms are presented as small spheres of arbitrary radii.

## Dicarbonyl( $\eta^5$ -cyclopentadienyl)[2-(phenylsulfanyl)ethyl]iron(II)

### Crystal data

[Fe(C <sub>5</sub> H <sub>5</sub> )(C <sub>8</sub> H <sub>9</sub> S)(CO) <sub>2</sub> ]	$F(000) = 1296$
$M_r = 314.17$	$D_x = 1.483 \text{ Mg m}^{-3}$
Orthorhombic, <i>Pbca</i>	Mo $K\alpha$ radiation, $\lambda = 0.71073 \text{ \AA}$
Hall symbol: -P 2ac 2ab	Cell parameters from 6363 reflections
$a = 10.3992 (5) \text{ \AA}$	$\theta = 2.3\text{--}27.3^\circ$
$b = 7.6402 (4) \text{ \AA}$	$\mu = 1.21 \text{ mm}^{-1}$
$c = 35.4173 (16) \text{ \AA}$	$T = 173 \text{ K}$
$V = 2814.0 (2) \text{ \AA}^3$	Plate, yellow
$Z = 8$	$0.57 \times 0.27 \times 0.08 \text{ mm}$

### Data collection

Bruker APEXII CCD area-detector diffractometer	2752 independent reflections
Radiation source: fine-focus sealed tube graphite	2339 reflections with $I > 2\sigma(I)$
$\phi$ and $\omega$ scans	$R_{\text{int}} = 0.068$
Absorption correction: integration ( <i>SADABS</i> ; Bruker, 2009)	$\theta_{\text{max}} = 26.0^\circ, \theta_{\text{min}} = 2.3^\circ$
$T_{\text{min}} = 0.545, T_{\text{max}} = 0.909$	$h = -12 \rightarrow 12$
20490 measured reflections	$k = -9 \rightarrow 9$
	$l = -41 \rightarrow 43$

### Refinement

Refinement on $F^2$	Primary atom site location: structure-invariant direct methods
Least-squares matrix: full	Secondary atom site location: difference Fourier map
$R[F^2 > 2\sigma(F^2)] = 0.075$	Hydrogen site location: inferred from neighbouring sites
$wR(F^2) = 0.153$	H-atom parameters constrained
$S = 1.26$	$w = 1/[\sigma^2(F_o^2) + (0.P)^2 + 19.462P]$
2752 reflections	where $P = (F_o^2 + 2F_c^2)/3$
172 parameters	$(\Delta/\sigma)_{\text{max}} = 0.004$
0 restraints	$\Delta\rho_{\text{max}} = 0.49 \text{ e \AA}^{-3}$
	$\Delta\rho_{\text{min}} = -0.75 \text{ e \AA}^{-3}$

*Special details*

**Geometry.** All e.s.d.'s (except the e.s.d. in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell e.s.d.'s are taken into account individually in the estimation of e.s.d.'s in distances, angles and torsion angles; correlations between e.s.d.'s in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell e.s.d.'s is used for estimating e.s.d.'s involving l.s. planes.

**Refinement.** Refinement of  $F^2$  against ALL reflections. The weighted  $R$ -factor  $wR$  and goodness of fit  $S$  are based on  $F^2$ , conventional  $R$ -factors  $R$  are based on  $F$ , with  $F$  set to zero for negative  $F^2$ . The threshold expression of  $F^2 > \sigma(F^2)$  is used only for calculating  $R$ -factors(gt) etc. and is not relevant to the choice of reflections for refinement.  $R$ -factors based on  $F^2$  are statistically about twice as large as those based on  $F$ , and  $R$ -factors based on ALL data will be even larger.

*Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters ( $\text{\AA}^2$ )*

	<i>x</i>	<i>y</i>	<i>z</i>	$U_{\text{iso}}^*/U_{\text{eq}}$
C1	0.5039 (8)	0.7838 (8)	0.40483 (18)	0.0425 (17)
H1	0.4331	0.8509	0.3961	0.051*
C2	0.5565 (8)	0.7851 (9)	0.44168 (19)	0.0459 (19)
H2	0.5266	0.8551	0.4620	0.055*
C3	0.6571 (7)	0.6697 (10)	0.4435 (2)	0.0461 (18)
H3	0.7090	0.6465	0.4650	0.055*
C4	0.6699 (7)	0.5910 (9)	0.4073 (2)	0.0430 (17)
H4	0.7308	0.5040	0.4004	0.052*
C5	0.5767 (7)	0.6644 (7)	0.38373 (17)	0.0348 (15)
H5	0.5648	0.6379	0.3578	0.042*
C6	0.5240 (6)	0.3405 (8)	0.45376 (15)	0.0279 (13)
C7	0.3395 (7)	0.5626 (8)	0.45266 (17)	0.0366 (15)
C8	0.3943 (6)	0.3817 (8)	0.39055 (16)	0.0333 (14)
H8A	0.3294	0.3098	0.4039	0.040*
H8B	0.3476	0.4590	0.3728	0.040*
C9	0.4800 (6)	0.2613 (8)	0.36798 (16)	0.0327 (14)
H9A	0.5282	0.1825	0.3851	0.039*
H9B	0.5424	0.3302	0.3530	0.039*
C10	0.4778 (7)	-0.0152 (8)	0.31408 (16)	0.0377 (15)
C11	0.6121 (7)	-0.0106 (7)	0.31470 (16)	0.0357 (15)
H11	0.6553	0.0756	0.3293	0.043*
C12	0.6823 (7)	-0.1315 (9)	0.29414 (17)	0.0392 (16)
H12	0.7736	-0.1261	0.2943	0.047*
C13	0.6207 (8)	-0.2602 (9)	0.27334 (18)	0.0471 (19)
H13	0.6691	-0.3431	0.2593	0.056*
C14	0.4893 (9)	-0.2665 (10)	0.27328 (18)	0.053 (2)
H14	0.4468	-0.3551	0.2592	0.063*
C15	0.4172 (8)	-0.1468 (9)	0.29330 (17)	0.0453 (18)
H15	0.3260	-0.1538	0.2930	0.054*
Fe1	0.48739 (8)	0.53522 (10)	0.42952 (2)	0.0264 (2)
O1	0.5463 (4)	0.2142 (5)	0.46915 (12)	0.0375 (11)
O2	0.2434 (5)	0.5784 (7)	0.46790 (14)	0.0558 (14)
S1	0.37701 (18)	0.1351 (2)	0.33694 (5)	0.0407 (4)

## supplementary materials

---

### *Atomic displacement parameters ( $\text{\AA}^2$ )*

	$U^{11}$	$U^{22}$	$U^{33}$	$U^{12}$	$U^{13}$	$U^{23}$
C1	0.059 (5)	0.025 (3)	0.043 (4)	-0.004 (3)	0.005 (4)	0.014 (3)
C2	0.066 (5)	0.035 (4)	0.037 (4)	-0.024 (4)	0.015 (3)	0.002 (3)
C3	0.043 (4)	0.050 (4)	0.045 (4)	-0.022 (4)	-0.010 (3)	0.017 (3)
C4	0.034 (4)	0.039 (4)	0.055 (4)	-0.002 (3)	0.012 (3)	0.018 (3)
C5	0.052 (4)	0.025 (3)	0.028 (3)	-0.008 (3)	0.008 (3)	0.006 (2)
C6	0.026 (3)	0.033 (3)	0.025 (3)	0.003 (3)	0.000 (2)	0.000 (2)
C7	0.046 (4)	0.033 (3)	0.031 (3)	0.004 (3)	-0.001 (3)	0.003 (3)
C8	0.035 (4)	0.032 (3)	0.034 (3)	-0.005 (3)	-0.003 (3)	0.006 (3)
C9	0.041 (4)	0.027 (3)	0.031 (3)	-0.006 (3)	0.001 (3)	-0.002 (2)
C10	0.058 (5)	0.033 (3)	0.022 (3)	-0.011 (3)	-0.008 (3)	0.006 (2)
C11	0.058 (4)	0.020 (3)	0.030 (3)	-0.002 (3)	-0.004 (3)	0.004 (2)
C12	0.049 (4)	0.036 (3)	0.033 (3)	0.002 (3)	0.003 (3)	0.008 (3)
C13	0.078 (6)	0.031 (3)	0.033 (4)	0.003 (4)	0.003 (4)	0.001 (3)
C14	0.079 (6)	0.047 (4)	0.033 (3)	-0.016 (4)	-0.002 (4)	-0.010 (3)
C15	0.061 (5)	0.042 (4)	0.032 (3)	-0.014 (4)	-0.009 (3)	0.003 (3)
Fe1	0.0295 (5)	0.0238 (4)	0.0259 (4)	-0.0012 (4)	0.0019 (4)	0.0043 (3)
O1	0.043 (3)	0.030 (2)	0.040 (2)	0.001 (2)	0.001 (2)	0.007 (2)
O2	0.044 (3)	0.063 (4)	0.060 (3)	0.009 (3)	0.016 (3)	0.006 (3)
S1	0.0453 (11)	0.0401 (9)	0.0367 (8)	-0.0059 (8)	-0.0100 (8)	-0.0045 (7)

### *Geometric parameters ( $\text{\AA}$ , $^\circ$ )*

C1—C5	1.401 (9)	C8—C9	1.510 (8)
C1—C2	1.415 (9)	C8—Fe1	2.054 (6)
C1—Fe1	2.098 (6)	C8—H8A	0.9900
C1—H1	0.9489	C8—H8B	0.9900
C2—C3	1.370 (11)	C9—S1	1.812 (6)
C2—Fe1	2.085 (6)	C9—H9A	0.9900
C2—H2	0.9490	C9—H9B	0.9900
C3—C4	1.422 (10)	C10—C15	1.396 (9)
C3—Fe1	2.101 (7)	C10—C11	1.397 (10)
C3—H3	0.9500	C10—S1	1.753 (7)
C4—C5	1.396 (9)	C11—C12	1.384 (9)
C4—Fe1	2.098 (7)	C11—H11	0.9500
C4—H4	0.9501	C12—C13	1.386 (9)
C5—Fe1	2.114 (6)	C12—H12	0.9500
C5—H5	0.9485	C13—C14	1.367 (11)
C6—O1	1.132 (7)	C13—H13	0.9500
C6—Fe1	1.759 (6)	C14—C15	1.379 (10)
C7—O2	1.142 (8)	C14—H14	0.9500
C7—Fe1	1.755 (7)	C15—H15	0.9500
C5—C1—C2	106.7 (6)	C11—C10—S1	125.1 (5)
C5—C1—Fe1	71.2 (3)	C12—C11—C10	120.2 (6)
C2—C1—Fe1	69.7 (4)	C12—C11—H11	119.9

C5—C1—H1	126.7	C10—C11—H11	119.9
C2—C1—H1	126.6	C11—C12—C13	120.6 (7)
Fe1—C1—H1	124.1	C11—C12—H12	119.7
C3—C2—C1	109.5 (7)	C13—C12—H12	119.7
C3—C2—Fe1	71.5 (4)	C14—C13—C12	119.2 (7)
C1—C2—Fe1	70.7 (4)	C14—C13—H13	120.4
C3—C2—H2	125.3	C12—C13—H13	120.4
C1—C2—H2	125.2	C13—C14—C15	121.3 (7)
Fe1—C2—H2	124.0	C13—C14—H14	119.4
C2—C3—C4	107.5 (6)	C15—C14—H14	119.4
C2—C3—Fe1	70.3 (4)	C14—C15—C10	120.2 (7)
C4—C3—Fe1	70.1 (4)	C14—C15—H15	119.9
C2—C3—H3	126.3	C10—C15—H15	119.9
C4—C3—H3	126.2	C7—Fe1—C6	93.6 (3)
Fe1—C3—H3	125.1	C7—Fe1—C8	88.2 (3)
C5—C4—C3	107.7 (6)	C6—Fe1—C8	87.0 (3)
C5—C4—Fe1	71.2 (4)	C7—Fe1—C2	95.5 (3)
C3—C4—Fe1	70.3 (4)	C6—Fe1—C2	126.8 (3)
C5—C4—H4	126.1	C8—Fe1—C2	145.5 (3)
C3—C4—H4	126.2	C7—Fe1—C4	160.7 (3)
Fe1—C4—H4	123.9	C6—Fe1—C4	99.1 (3)
C4—C5—C1	108.5 (6)	C8—Fe1—C4	106.9 (3)
C4—C5—Fe1	70.0 (3)	C2—Fe1—C4	65.2 (3)
C1—C5—Fe1	70.0 (3)	C7—Fe1—C1	99.1 (3)
C4—C5—H5	125.7	C6—Fe1—C1	162.0 (3)
C1—C5—H5	125.8	C8—Fe1—C1	106.0 (3)
Fe1—C5—H5	125.9	C2—Fe1—C1	39.6 (3)
O1—C6—Fe1	179.2 (6)	C4—Fe1—C1	65.5 (3)
O2—C7—Fe1	179.2 (6)	C7—Fe1—C3	124.6 (3)
C9—C8—Fe1	115.2 (4)	C6—Fe1—C3	96.7 (3)
C9—C8—H8A	108.5	C8—Fe1—C3	146.4 (3)
Fe1—C8—H8A	108.5	C2—Fe1—C3	38.2 (3)
C9—C8—H8B	108.5	C4—Fe1—C3	39.6 (3)
Fe1—C8—H8B	108.5	C1—Fe1—C3	65.6 (3)
H8A—C8—H8B	107.5	C7—Fe1—C5	133.5 (3)
C8—C9—S1	107.3 (4)	C6—Fe1—C5	132.4 (3)
C8—C9—H9A	110.3	C8—Fe1—C5	87.6 (2)
S1—C9—H9A	110.3	C2—Fe1—C5	65.1 (2)
C8—C9—H9B	110.3	C4—Fe1—C5	38.7 (3)
S1—C9—H9B	110.3	C1—Fe1—C5	38.9 (2)
H9A—C9—H9B	108.5	C3—Fe1—C5	65.4 (3)
C15—C10—C11	118.5 (7)	C10—S1—C9	106.0 (3)
C15—C10—S1	116.4 (6)		
C5—C1—C2—C3	-0.6 (7)	C5—C4—Fe1—C8	63.4 (4)
Fe1—C1—C2—C3	61.4 (5)	C3—C4—Fe1—C8	-179.1 (4)
C5—C1—C2—Fe1	-62.0 (4)	C5—C4—Fe1—C2	-80.6 (4)
C1—C2—C3—C4	-0.4 (7)	C3—C4—Fe1—C2	37.0 (4)
Fe1—C2—C3—C4	60.5 (4)	C5—C4—Fe1—C1	-36.9 (4)
C1—C2—C3—Fe1	-60.8 (5)	C3—C4—Fe1—C1	80.7 (4)

## supplementary materials

---

C2—C3—C4—C5	1.2 (7)	C5—C4—Fe1—C3	-117.6 (6)
Fe1—C3—C4—C5	61.8 (4)	C3—C4—Fe1—C5	117.6 (6)
C2—C3—C4—Fe1	-60.6 (5)	C5—C1—Fe1—C7	-155.6 (4)
C3—C4—C5—C1	-1.6 (7)	C2—C1—Fe1—C7	87.6 (5)
Fe1—C4—C5—C1	59.6 (4)	C5—C1—Fe1—C6	69.9 (9)
C3—C4—C5—Fe1	-61.2 (4)	C2—C1—Fe1—C6	-46.9 (10)
C2—C1—C5—C4	1.4 (7)	C5—C1—Fe1—C8	-64.9 (5)
Fe1—C1—C5—C4	-59.6 (4)	C2—C1—Fe1—C8	178.4 (4)
C2—C1—C5—Fe1	61.0 (4)	C5—C1—Fe1—C2	116.7 (6)
Fe1—C8—C9—S1	178.1 (3)	C5—C1—Fe1—C4	36.8 (4)
C15—C10—C11—C12	1.9 (9)	C2—C1—Fe1—C4	-80.0 (5)
S1—C10—C11—C12	-177.0 (4)	C5—C1—Fe1—C3	80.4 (5)
C10—C11—C12—C13	-1.3 (9)	C2—C1—Fe1—C3	-36.3 (4)
C11—C12—C13—C14	0.1 (10)	C2—C1—Fe1—C5	-116.7 (6)
C12—C13—C14—C15	0.4 (11)	C2—C3—Fe1—C7	-46.6 (5)
C13—C14—C15—C10	0.3 (11)	C4—C3—Fe1—C7	-164.7 (4)
C11—C10—C15—C14	-1.4 (9)	C2—C3—Fe1—C6	-145.7 (4)
S1—C10—C15—C14	177.6 (5)	C4—C3—Fe1—C6	96.2 (4)
C9—C8—Fe1—C7	-160.1 (5)	C2—C3—Fe1—C8	119.7 (5)
C9—C8—Fe1—C6	-66.4 (5)	C4—C3—Fe1—C8	1.6 (7)
C9—C8—Fe1—C2	102.8 (6)	C4—C3—Fe1—C2	-118.1 (6)
C9—C8—Fe1—C4	32.3 (5)	C2—C3—Fe1—C4	118.1 (6)
C9—C8—Fe1—C1	100.9 (5)	C2—C3—Fe1—C1	37.6 (4)
C9—C8—Fe1—C3	31.2 (7)	C4—C3—Fe1—C1	-80.5 (4)
C9—C8—Fe1—C5	66.3 (4)	C2—C3—Fe1—C5	80.5 (4)
C3—C2—Fe1—C7	143.1 (4)	C4—C3—Fe1—C5	-37.6 (4)
C1—C2—Fe1—C7	-97.6 (5)	C4—C5—Fe1—C7	153.7 (5)
C3—C2—Fe1—C6	44.4 (5)	C1—C5—Fe1—C7	34.2 (6)
C1—C2—Fe1—C6	163.7 (4)	C4—C5—Fe1—C6	-37.4 (5)
C3—C2—Fe1—C8	-122.0 (6)	C1—C5—Fe1—C6	-156.9 (4)
C1—C2—Fe1—C8	-2.7 (8)	C4—C5—Fe1—C8	-121.1 (4)
C3—C2—Fe1—C4	-38.3 (4)	C1—C5—Fe1—C8	119.4 (4)
C1—C2—Fe1—C4	81.0 (5)	C4—C5—Fe1—C2	80.7 (5)
C3—C2—Fe1—C1	-119.3 (6)	C1—C5—Fe1—C2	-38.8 (4)
C1—C2—Fe1—C3	119.3 (6)	C1—C5—Fe1—C4	-119.5 (6)
C3—C2—Fe1—C5	-81.1 (4)	C4—C5—Fe1—C1	119.5 (6)
C1—C2—Fe1—C5	38.1 (4)	C4—C5—Fe1—C3	38.4 (4)
C5—C4—Fe1—C7	-76.5 (10)	C1—C5—Fe1—C3	-81.0 (5)
C3—C4—Fe1—C7	41.1 (10)	C15—C10—S1—C9	169.7 (5)
C5—C4—Fe1—C6	153.0 (4)	C11—C10—S1—C9	-11.5 (6)
C3—C4—Fe1—C6	-89.5 (4)	C8—C9—S1—C10	-175.3 (4)

Fig. 1

